



# PAVILION<sup>TM</sup>

## 3.3 EC HERBICIDE

GROUP	3	HERBICIDE
-------	---	-----------

FOR USE IN SELECTED CROPS

**ACTIVE INGREDIENT:**

Pendimethalin: N-(1-ethylpropyl)-3, 4-dimethyl-2, 6-dinitrobenzenamine..... **% BY WT.** 37.4%

**OTHER INGREDIENTS\*:** ..... 62.6%

**TOTAL:** ..... 100.0%

(1 gallon contains 3.3 pounds of pendimethalin)

\*Contains aromatic naphtha

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION / PRECAUCIÓN**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

See inside for complete **First Aid, Precautionary Statements, Directions for Use, Conditions of Sale and Warranty**, and state-specific crop and/or use site restrictions.

In case of an emergency endangering life or property involving this product, call Chemtrec **1-800-424-9300**.

EPA Reg. No.: 89167-29-89391

# HERBICIDE



Distributed By:  
**INNVICTIS<sup>®</sup> CROP CARE, LLC**  
1880 Fall River Drive, Suite 100  
Loveland, CO 80538

051917R063017A

**FOR USE IN ALFALFA (FORAGE, HAY OR SEED PRODUCTION) BEARING CITRUS FRUIT TREES, BEARING NUT TREES, BEARING POME FRUIT TREES, BEARING STONE FRUIT TREES, CARROTS, CORN (FIELD, POP, SEED, SWEET), COTTON, EDIBLE BEANS, FORAGE LEGUMES, FRUITING VEGETABLES (PEPPER, TOMATO), GARLIC, GRAIN SORGHUM, LEEK, LENTILS AND PEAS, MINT, NONBEARING FRUIT TREE AND NUT TREE CROPS, NONBEARING VINEYARDS, ONIONS AND SHALLOTS (DRY BULB, GREEN), PEANUTS, POTATOES, RICE, SOYBEANS, SUGARCANE, SUNFLOWERS, TOBACCO AND WHEAT.**

<b>FIRST AID</b>	
<b>If Swallowed:</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advise.</li> <li>• Do not give any liquid to person.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>If in Eyes:</b>	<ul style="list-style-type: none"> <li>• Hold eyes open and rinse slowly and gently with water for 15 to 20 minutes.</li> <li>• Remove contact lenses, if present, after first 5 minutes; then continue rinsing eyes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If on Skin or Clothing:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>NOTE TO PHYSICIAN</b> - Because of increased risk of chemical pneumonia or pulmonary edema caused by aspiration of the hydrocarbon solvent, vomiting should be induced only under professional supervision.	
<b>HOT LINE NUMBER</b> - Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact CHEMTREC: 1-800-424-9300.	

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION.** Causes moderate eye irritation. Harmful if swallowed or absorbed through skin. Avoid contact with eyes, skin or clothing.

**Personal Protective Equipment (PPE)**

Some materials that are chemical-resistant to this product are listed below.

**Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of waterproof materials, such as barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, or viton ≥ 14 mils
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**Engineering Controls:**

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [(40 CFR 170.240)(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

**USER SAFETY RECOMMENDATIONS**

**Users should:**

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent aquatic sites. Do not contaminate water when disposing of equipment washwaters or rinsate.

**Endangered Species Protection**

If endangered plant species occur in proximity to the application site, the following mitigation measures are required:

- If applied by ground, leave an untreated buffer zone of 200 feet. The product must be applied using a low boom (20 inches above the ground) and ASAE fine to medium/coarse nozzles.
- If applied by air, leave an untreated buffer zone of 170 feet. Must use straight-stream nozzles (D-6 or larger); wind can be no more than 8 mph; and release height must be 15 feet or less.

To determine whether your county has an endangered species, consult the website <http://www.epa.gov/espp/usa-map.htm>.

Endangered Species Bulletins may also be obtained from extension offices or state pesticide agencies. If the bulletin is not available for your specific area, check with the appropriate local state agency to determine if known populations of endangered species occur in the area to be treated.

#### DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. This label must be in the possession of the user at the time of pesticide application.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide protection.

Observe all cautions and restrictions in this label and the labels of products used in combination with *PAVILION 3.3 EC HERBICIDE*. The use of *PAVILION 3.3 EC HERBICIDE* not consistent with this label can result in injury to crops, animals, or persons. Keep containers closed to avoid spills and contamination.

Do not allow spray to drift from the application site and contact people, structures people occupy at any time and the associated property, parks and recreation areas, non-target crops, aquatic and wetland areas, woodlands, pastures, rangelands, or animals.

Do not enter or allow other people (or pets) to enter the treated area until sprays have dried.

Innvictis intends that this product may not be used for manufacturing products for application to turf and ornamentals.

Innvictis does not recommend or authorize the use of this product in manufacturing, processing or preparing custom blends with other products for application to field and row crops or to orchard, grove, and vineyard crops.

#### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours. Exception:** if the product is soil injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of waterproof materials, such as barrier laminate, butyl rubber  $\geq 14$  mils, nitrile rubber  $\geq 14$  mils, neoprene rubber  $\geq 14$  mils, or viton  $\geq 14$  mils
- Shoes plus socks

## Product Information

**PAVILION 3.3 EC HERBICIDE** is a selective herbicide for controlling most annual grasses and certain broadleaf weeds as they germinate. Refer to **Table 1** for a complete list of controlled weeds. **PAVILION 3.3 EC HERBICIDE** will not control established weeds.

**Table 1. Weeds Controlled:**

(see crop sections for additional weeds controlled)

Weeds controlled with <b>PAVILION 3.3 EC HERBICIDE</b> applied up to 4.8 pts/A				
GRASSES				
Annual ryegrass	Downy brome* (Cheatgrass)	Hairy chess** <sup>a</sup>	Jointed goatgrass** <sup>a</sup>	Shattercane*
Barnyardgrass	Foxtail, giant	Itchgrass*	Oat, wild*	Signalgrass*
Canarygrass** <sup>b</sup>	Foxtail, green	Italian ryegrass*	Panicum, fall	Wild proso millet*
Cheat** <sup>b</sup>	Foxtail, yellow	Japanese brome** <sup>a</sup>	Panicum, Texas	Witchgrass
Crabgrass	Goosegrass	Johnsongrass (seedling)	Sandbur, field	Woolly cupgrass*
Crowfootgrass				
BROADLEAVES				
Amaranth, Palmer	Henbit	Lambsquarters, Slimleaf	Purslane	Spurge, Annual
Bugloss, small*	Kochia	London Rocket*	Pusley, Florida	Velvetleaf*
Carpetweed	Lady's Thumb	Mustard, black <sup>o</sup>	Sheeperspurse*	Waterhemp Species
Chickweed, Common*	Lambsquarters, Common	Pigweed Species	Smartweed, Pennsylvania*	

\* Suppression, but controlled when **PAVILION 3.3 EC HERBICIDE** use rate exceeds 4.8 pts/A.

a Neither suppressed nor controlled in California.

b Not controlled in California.

Weeds controlled with <b>PAVILION 3.3 EC HERBICIDE</b> applied at 4.8 pts/A or greater				
GRASSES				
Annual Bluegrass	Grass, Guinea <sup>a</sup>	Lovegrass		Sprangletop, red
Browntop panicum	Junglerice	Sprangletop, Mexican		Swollen fingergrass
BROADLEAVES				
Dodder <sup>†</sup>	Fiddleneck	Morningglory**	Prostrate, knotweed	Puncturevine

<sup>†</sup> For optimum dodder control, use the highest labeled rate of **PAVILION 3.3 EC HERBICIDE** specified in the specific crop.

\*\* Suppression

b Not controlled in California

### Precautions

- Unusually cold, excessively wet, or hot and dry conditions that delay germination or extend germination over a long period of time can reduce weed control.
- Over application can result in crop-stand loss, crop injury, or soil residues.
- Uneven application or improper soil incorporation can decrease weed control or cause crop injury.
- Soil incorporation deeper than specified can reduce weed control.
- Seedling diseases, cold weather, excessive moisture, shallow or deep planting, low or high soil pH, high soil salt concentration, or drought can weaken seedlings and plants and increase the possibility of crop damage from **PAVILION 3.3 EC HERBICIDE**. Under these conditions, crop yields can be reduced.

### Mode of Action

**PAVILION 3.3 EC HERBICIDE** is a meristematic inhibitor that interferes with the plant's cellular division or mitosis. This and/or other products with the meristematic inhibiting mode of action may not effectively control naturally occurring biotypes of some of the weeds listed on this label. A weed biotype is a naturally occurring plant within a given species that has a slightly different, but distinct, genetic makeup from other plants. Other herbicides with the meristematic inhibiting mode of action include other dinitroaniline herbicides, such as trifluralin. If naturally occurring meristematic inhibiting resistant biotypes are present in a field, **PAVILION 3.3 EC HERBICIDE** and/or any other meristematic inhibiting mode of action herbicide should be tank mixed or applied sequentially with an appropriate registered herbicide having a different mode of action to ensure control.

### Application Rate

Use rates for *PAVILION 3.3 EC HERBICIDE* when used alone, in tank mix, or sequential applications are given in **Crop-specific Information**. Use rates of this product vary by soil texture and organic matter. See **Table 2** for soil texture groupings used in this label.

**Table 2. Soil Texture Groups**

COARSE	MEDIUM	FINE
Sands Loamy sands Sandy loams	Sandy clay loams* Sandy clays Loams Silt loams Silt	Silty clay loams* Silty clays Clay loams Clays
* Sometimes considered transitional soils and may be classified as either medium- or fine- textured soils.		
<b>Peat and Muck soils:</b> <i>PAVILION 3.3 EC HERBICIDE</i> may be used on peat and muck soils, but weed control may be inconsistent and/or reduced. Use maximum labeled use rate allowed in the specific crop.		

### Application Timings

*PAVILION 3.3 EC HERBICIDE* will provide most effective weed control when applied by ground or aerial equipment and subsequently incorporated into soil within 7 days after application by rainfall, sprinkler irrigation, or mechanical tillage prior to weed seedling emergence from soil. *PAVILION 3.3 EC HERBICIDE* can also be applied through chemigation, including flooded basin irrigation systems. *PAVILION 3.3 EC HERBICIDE* is labelled for preplant surface, preplant incorporated, surface incorporated, preemergence, early postemergence, postemergence incorporated (CULTI-SPRAY) or layby treatment. See **Crop-specific Information** for specific application directions by crop.

**Preplant Surface Applications:** For use in minimum tillage or no-tillage production systems, apply *PAVILION 3.3 EC HERBICIDE* alone or in tank mixes up to 45 days before planting. When making early preplant surface applications (15 to 45 days prior to planting), *PAVILION 3.3 EC HERBICIDE* should be tank mixed or followed by a postemergence herbicide application. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Rainfall or sprinkler irrigation within 7 days after application is required to move this product into the upper soil surface where weed seeds germinate.

**Preplant Incorporated Applications:** Apply *PAVILION 3.3 EC HERBICIDE* and incorporate into the upper (1 inch to 2 inches) soil surface up to 60 days before planting. Use an implement capable of giving uniform incorporation; two-pass incorporation usually results in a more consistent result.

**Surface Incorporated Applications:** Uniformly apply *PAVILION 3.3 EC HERBICIDE* as broadcast or banded treatment to soil surface underneath established trees and/or in ground areas between trees rows. Within 7 days after application, incorporate into upper (1 inch to 2 inches) soil surface using either rainfall, sprinkler irrigation, or shallow mechanical incorporation using an implement capable of giving uniform incorporation; two-pass mechanical incorporation usually results in a more consistent result.

**Preemergence Surface Applications:** Broadcast treatment uniformly to the soil surface at planting and up to 2 days after planting. Rainfall, sprinkler irrigation, or shallow mechanical incorporation within 7 days after application is required to move this product into the upper soil surface where weed seeds germinate. If adequate rainfall or irrigation does not occur and weed seedling emergence begins, a shallow cultivation or rotary hoeing will improve performance.

**Early Postemergence Applications:** *PAVILION 3.3 EC HERBICIDE* must be applied prior to weed seedling emergence or in a tank mix with products that control the emerged weeds. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Refer to **Crop-specific Information** for specific postemergence application specified by crop.

### Postemergence Incorporated Applications

(CULTI-SPRAY): Prior to application, crop must be cultivated in such a manner as to throw at least one inch of soil over the base of the crop plants. This will prevent direct contact of *PAVILION 3.3 EC HERBICIDE* and the zone of brace root formation. *PAVILION 3.3 EC HERBICIDE* must be applied broadcast with a ground sprayer when crop is at least 4 inches tall up to layby. Use drop nozzles if crop foliage will prevent uniform coverage of the soil surface within the rows. Thoroughly and uniformly incorporate *PAVILION 3.3 EC HERBICIDE* treatments into the soil with:

1. a sweep-type or rolling cultivator set to provide thorough incorporation in the top 1 inch of soil, or
2. adequate overhead irrigation water or rainfall. See **Crop-specific Information (Corn and Grain Sorghum)** for more details on (CULTI-SPRAY) application.

**Layby Application:** Apply *PAVILION 3.3 EC HERBICIDE* directly to the soil between rows as a directed spray following the last normal cultivation (layby). See **Crop-specific Information** for more details on layby application.

**Split Applications:** *PAVILION 3.3 EC HERBICIDE* may be applied preplant incorporated up to 60 days prior to planting and followed by a preemergence application at planting or up to 2 days after planting. The total amount of *PAVILION 3.3 EC HERBICIDE* applied per acre per season cannot exceed the highest labeled rate for any given soil type. See **Crop-specific Information** for more details on split applications.

**Fall Applications:** *PAVILION 3.3 EC HERBICIDE* may be used in fall application programs in certain crops. See **Crop-specific Information** for details on fall application timing.

## Spraying Instructions

*PAVILION 3.3 EC HERBICIDE* may be applied using either water or sprayable fluid fertilizer (such as straight 32-0-0 or 28-0-0) as the spray carrier. Additionally, *PAVILION 3.3 EC HERBICIDE* may be impregnated on dry bulk fertilizer. Sprayable fluid fertilizer as a carrier is not recommended for use after crop emergence unless the typical fertilizer burn symptoms on the crop are acceptable.

### Aerial Applications

Uniformly apply in 5 or more gallons of water per acre. Exercise caution to minimize drift. Do not apply during periods of gusty winds or when wind conditions favor drifting. Spray drift can cause injury to sensitive crops. It is recommended that a flagman or an automatic mechanical flagging unit on the aircraft be used to avoid overlapping and possible crop injury.

### Ground Applications (Broadcast)

Uniformly apply with properly calibrated ground equipment in 10 or more gallons of water per acre or 20 or more gallons of liquid fertilizer per acre. Use sprayers equipped with appropriate nozzles that provide uniform and accurate spray distribution and minimize drift. Keep the bypass line on or near the bottom of the tank to minimize foaming. Nozzle and in-line screens must be no finer than 50 mesh. Application of *PAVILION 3.3 EC HERBICIDE* during periods of gusty winds may result in uneven applications. Do not apply *PAVILION 3.3 EC HERBICIDE* postemergence in liquid fertilizers.

If liquid fertilizer/herbicide(s) mixture separates in the spray tank, clogged equipment and uneven application can result. Always predetermine the compatibility of *PAVILION 3.3 EC HERBICIDE* alone or with other herbicides based on the following compatibility "**jar test**":

1. Add 1 pint of fertilizer to a quart jar.
2. Add 1 to 4 teaspoon(s) of the Dry Flowable (DF), Wettable Powder (WP), Aqueous Solution (AS), Flowable (F) or Liquid (L) formulation (depending on mixing ratio required) to the liquid fertilizer. The number of teaspoons of the formulation to add can be determined by the following formula:
$$\frac{\text{lbs or pts of product/acre}}{\text{gallons of fertilizer/acre}} \times 11.4 = \frac{\text{number of teaspoons of herbicide to add to}}{1 \text{ pint of fertilizer}}$$
3. Close the jar and agitate until the herbicide(s) are evenly dispersed in the liquid fertilizer. If the materials do not disperse well, it may be necessary to slurry the chemicals in water before adding to the fertilizer.
4. After dispersing the materials, add appropriate number of teaspoons of *PAVILION 3.3 EC HERBICIDE* to the jar and shake well. Add water soluble concentrate herbicides to the mixture last and agitate. Let the mixture stand for 30 minutes and then observe the results. Look for signs of separation: an oily layer or globules, sludge, flakes or other precipitates.
5. Evaluate compatibility.
  - (a) If the herbicide(s) and liquid fertilizer mixture does not separate, use this mixture in your spray tank.
  - (b) If the mixture separates but mixes readily with shaking, the mixture can be used provided that good agitation is maintained in the spray tank.
  - (c) If separation of the mixture occurs and agitation does not correct this problem, a compatibility agent is needed.
6. If the need for a compatibility agent is demonstrated, the following procedure is recommended: Using a clean quart jar, repeat step 1 above and add 1/2 teaspoon of the compatibility agent to the liquid fertilizer. Mix well and repeat steps 2, 3 and 4. If separation or precipitation occurs with the compatibility agent, do not use *PAVILION 3.3 EC HERBICIDE* with that specific liquid fertilizer.

### Ground Applications (Band)

Uniformly apply the broadcast equivalent rate and volume per acre. To determine these:

$$\frac{\text{Band Width in Inches}}{\text{Row Width in Inches}} \times \text{Broadcast Rate per Acre} = \text{Band Rate per Acre}$$
$$\frac{\text{Band Width in Inches}}{\text{Row Width in Inches}} \times \text{Broadcast Volume per Acre} = \text{Band Volume per Acre}$$

### Ground Applications (Dry Bulk Fertilizer)

Apply *PAVILION 3.3 EC HERBICIDE* dry bulk fertilizer mixtures only with ground equipment. Do not impregnate *PAVILION 3.3 EC HERBICIDE* onto coated ammonium nitrate or limestone because these materials will not absorb the herbicide. Dry fertilizer blends containing mixtures of ammonium nitrate or limestone may be impregnated with *PAVILION 3.3 EC HERBICIDE*. A minimum of 200 pounds of impregnated dry bulk fertilizer, excluding the weight of ammonium nitrate or limestone, must be applied per acre.

Use the following formula to determine the amount (in pints) of *PAVILION 3.3 EC HERBICIDE* to be impregnated on a ton of dry bulk fertilizer based on the rate of fertilizer to be applied per acre:

$$\frac{2000}{\text{Pounds of Dry Fertilizer per Acre}} \times \text{Pints of } \textit{PAVILION 3.3 EC HERBICIDE} \text{ (Specified Rate per Acre)} = \text{Pints of } \textit{PAVILION 3.3 EC HERBICIDE} \text{ per Ton of Fertilizer}$$

To impregnate *PAVILION 3.3 EC HERBICIDE* on bulk fertilizer, use a closed rotary-drum mixer or other commonly used dry bulk fertilizer blender equipped with suitable spray equipment. Spray nozzles must be placed to provide uniform coverage of *PAVILION 3.3 EC HERBICIDE* onto the fertilizer during mixing.

Apply the *PAVILION 3.3 EC HERBICIDE*/dry bulk fertilizer mixture with an accurately calibrated dry fertilizer spreader. The *PAVILION 3.3 EC HERBICIDE*/dry bulk fertilizer mixture must be spread uniformly on the soil surface.

### Chemigation Applications via Sprinkler Irrigation Systems

*PAVILION 3.3 EC HERBICIDE* may be applied as a chemigation treatment through sprinkler irrigation systems. Refer to **Crop-specific Information** sections for individual crops.

Apply this product only through a sprinkler irrigation system of the following type: center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move.

Uniform distribution of *PAVILION 3.3 EC HERBICIDE*-treated irrigation water is the sole responsibility of the applicator and is required to avoid crop injury, lack of herbicide effectiveness or illegal pesticide residues in the crop. If you have any questions about calibration, you should contact state extension service specialists, equipment manufacturers, or other experts.

The system must be properly calibrated (with water only) to ensure that the amount of *PAVILION 3.3 EC HERBICIDE* applied corresponds to the specified rate. Apply *PAVILION 3.3 EC HERBICIDE* in 1/2 to 3/4 inches of water during the first sprinkler set (use at least 1 inch of water in the states of Texas, New Mexico and Oklahoma). Maintain agitation in the injection nurse tank to keep a uniform herbicide suspension during application. When application is complete, flush the system with water.

### Restrictions

- Do not apply *PAVILION 3.3 EC HERBICIDE* via chemigation to crops unless specified in **Crop-specific Information** section.
- Do not apply this product through any other type of sprinkler irrigation system.

### Chemigation Instructions (for low volume micro sprinklers)

Output of low volume sprinkler = 4 to 50 gallons per hour (gph) per emitter. Point of application must be above ground.

Irrigation system should run a sufficient amount of time prior to *PAVILION 3.3 EC HERBICIDE* injection to have all emitters functioning properly. After system is operating properly, length of injection should be such that at one period of time during the injection, the first and last emitters in the system contain *PAVILION 3.3 EC HERBICIDE*-treated water. Add *PAVILION 3.3 EC HERBICIDE* to the supply tank already filled with the volume of water required for the injection period. Maintain proper agitation in *PAVILION 3.3 EC HERBICIDE* injection tank. *PAVILION 3.3 EC HERBICIDE* should be mixed in clean water and injected down-line from filters. Following *PAVILION 3.3 EC HERBICIDE* injection, system should be flushed for a period of time sufficient to clear the line of *PAVILION 3.3 EC HERBICIDE*. (If *PAVILION 3.3 EC HERBICIDE* application is made during a normal irrigation cycle, injection should be made during the last stage.)

### Chemigation Calibration (for low volume micro sprinklers)

Calculation of use rate is based on wetted area around emitters - not on tree acres. To determine correct amount of *PAVILION 3.3 EC HERBICIDE*, use the following formula:

- Treated area per each emitter = A

$$A = 3.14 \times (\text{radius} \times \text{radius})$$

- The area in square feet wet in each acre = B

$$B = \frac{A \times \text{emitters/acre}}{144}$$

- The total area (in square feet) wet by your system = C

$$C = B \times \text{acres covered by system.}$$

- Rate per treated acre of *PAVILION 3.3 EC HERBICIDE* (based on length of control desired) = R

Amount of *PAVILION 3.3 EC HERBICIDE* to inject = S

$$S = \frac{C}{43,560} \times R = \text{Qts of PAVILION 3.3 EC HERBICIDE}$$

### Example:

If the average distance from emitter to perimeter of wetted area measured one inch below soil surface is 13 inches, then

$$A = 3.14 \times (13'' \times 13'')$$

and A = 530.7 square inches.

If there are 300 emitters per acre, then

$$B = \frac{530.7 \times 300}{144}$$

and B = 1105.6 square feet wetted per acre.

If the system covers 20 acres, then

$$C = 1105.6 \text{ square feet per acre} \times 20 \text{ acres and}$$

$$C = 22,112 \text{ square feet wetted by system.}$$

If the desired application rate per treated acre is 2.4 qts of *PAVILION 3.3 EC HERBICIDE*, then

$$S = \frac{22,112}{43,560} \times 2.4$$

and S = 1.2 qts of *PAVILION 3.3 EC HERBICIDE* should be injected into the system.

### Special Precautions for Chemigation

1. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.
2. The pesticide injection pipeline must contain a functional, automatic quick-closing check valve to prevent the flow of fluid back toward the injection pump. It must also contain a functional, normally closed solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
3. The sprinkler chemigation system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. In addition, systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
4. The sprinkler chemigation system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
5. The irrigation line or water pump must include a functional pressure switch that will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

### Restrictions for Chemigation

- Do not apply when wind speed favors drift beyond the area intended for treatment.
- Do not connect an irrigation system used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

### Chemigation Systems Connected to Public Water Systems

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
3. All chemigation systems connected to public water systems must also follow restrictions listed in the preceding section titled **Chemigation**.

### Applications via Flooded Basin Irrigation Systems

*PAVILION 3.3 EC HERBICIDE* may be applied via flooded basin irrigation systems, but only to the following crops: bearing and nonbearing fruit and nut trees, nonbearing vineyards, and alfalfa grown for forage, hay or seed production.

### Use Instructions for Flooded Basin Irrigation

- *PAVILION 3.3 EC HERBICIDE* may be applied through a flood basin irrigation system designed to uniformly distribute irrigation water along the soil surface. Solid set systems utilizing tail riser for overhead application are excluded.
- Follow all label directions for *PAVILION 3.3 EC HERBICIDE* regarding rates per acre, timing of application, and crop-specific restrictions.
- A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.
- Mix *PAVILION 3.3 EC HERBICIDE* with water at a 1:1 ratio in the injection nurse tank to assist with product flowability. Maintain agitation in the injection nurse tank to keep a uniform herbicide suspension during application. When application is complete, flush the system with water.
- Tail water (runoff water) from flood irrigation that contains *PAVILION 3.3 EC HERBICIDE* should be re-circulated and contained in the field of initial application or used only on adjacent tree or vine crops or alfalfa for which *PAVILION 3.3 EC HERBICIDE* is registered for this type of application.
- Systems using a gravity-flow pesticide dispensing system must meter the pesticide in the water at the head of the field downstream of a hydraulic discontinuity, such as a drop structure or weir box, to decrease potential for water source contamination from backflow.
- Systems utilizing a pressurized water and pesticide injection system must meet the following requirements:
  - The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located in the irrigation pipe to prevent water source contamination from backflow.
  - The pesticide injection pipeline must contain a functional automatic quick closing check valve to prevent flow of fluids back towards the injection pump.
  - The pesticide injection pipeline must also contain a functional, normally closed solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
  - The system must contain a functional interlocking control to automatically shut off the pesticide injection pump when the water pump stops.
  - The irrigation pipe or water pump must include a functional pressure switch, which will stop the pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
  - Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump), of effective design and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
  - Any alternative to the above safety devices must conform to the list of EPA-approved alternative devices.
- Be sure to regularly measure the flow in the field to ensure the correct amount of *PAVILION 3.3 EC HERBICIDE* is being metered into the irrigation water and also regularly monitor to ensure that treated water is being uniformly distributed across the field. Flow rates through metering devices and distribution of *PAVILION 3.3 EC HERBICIDE* can vary with water temperature and speed of water flow across the field.
- Uniform distribution of *PAVILION 3.3 EC HERBICIDE*-treated irrigation water is the sole responsibility of the applicator and is required to avoid crop injury, lack of herbicide effectiveness, or illegal pesticide residues in the crop.
- If you have questions about calibration, contact a state extension service specialists, equipment manufacturer or other expert.

### Restrictions for Flooded Basin Irrigation

- Do not connect an irrigation system used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.



## Managing Off-Target Movement

### SPRAY DRIFT

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment and weather-related factors determines the potential for spray drift. The applicator and grower are responsible for considering all these factors when making decisions. It is the responsibility of the applicator to avoid spray drift on nontarget areas.

The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops:

1. The distance of the outermost nozzles on the boom must not exceed 3/4 the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

Where states have more stringent regulations, they must be observed.

The applicator should be familiar with and take into account the information covered in the Spray Drift Reduction Advisory Information presented below.

### INFORMATION ON DROPLET SIZE

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential but will not prevent drift if applications are made improperly or under unfavorable environmental conditions (see **WIND, TEMPERATURE AND HUMIDITY**, and **TEMPERATURE INVERSIONS**).

### CONTROLLING DROPLET SIZE

**Volume** - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.

**Pressure** - Do not exceed the nozzle manufacturer's specified pressures. For many nozzle types, lower pressure produces larger droplets. When high flow rates are needed, use higher flow rate nozzles instead of increasing pressure.

**Number of Nozzles** - Use the minimum number of nozzles that provide uniform coverage.

**Nozzle Orientation** - Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is recommended practice. Significant deflection from the horizontal will reduce droplet size and increase drift potential.

**Nozzle Type** - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid- or straight-stream nozzles oriented straight back produce the largest droplets and the lowest drift.

### BOOM LENGTH

For some use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width.

### APPLICATION HEIGHT

Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

### SWATH ADJUSTMENT

When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the air-craft upwind. Swath adjustment distance should increase with increasing drift potential (higher wind, smaller droplets, etc.).

### WIND

Drift potential is lowest between wind speeds of 2 to 10 mph. However, many factors, including droplet size and equipment type, determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential.

**Note:** Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

### TEMPERATURE AND HUMIDITY

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

### TEMPERATURE INVERSIONS

Applications must not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light, variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

### SENSITIVE AREAS

This pesticide must only be applied when the potential for drift to adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for threatened or endangered species, or nontarget crops or plants) is minimal (e.g. when wind is blowing away from the sensitive areas).

### ADDITIVES

Spray adjuvants have little or no influence on performance of *PAVILION 3.3 EC HERBICIDE* when applications are made prior to weed emergence. However, several tank mixes with *PAVILION 3.3 EC HERBICIDE* require adjuvants to improve burndown of emerged weeds. Therefore, surfactants, liquid fertilizer (28%, 30%, or 32% UAN (urea ammonium nitrate) or ammonium sulfate), or crop oil concentrate may be used with *PAVILION 3.3 EC HERBICIDE* tank mixes applied preplant, preemergence, or early postemergence to the crop. Follow the adjuvant specifications on the tank mix partner's label.

When an adjuvant (or a specific adjuvant product, such as a drift control agent) is to be used with this product, the use of a Chemical Producers and Distributors Association (CPDA) certified adjuvant is recommended. The recommended adjuvants must contain ingredients accepted by the EPA.

## TANK MIXING INFORMATION

*PAVILION 3.3 EC HERBICIDE* may be applied in a tank mix or a sequential application with other herbicides registered for use in a given crop. Refer to the companion label for weeds controlled in addition to *PAVILION 3.3 EC HERBICIDE* alone.

When using tank mixtures or sequential applications with *PAVILION 3.3 EC HERBICIDE*, always read the companion product label(s) to determine the specific use rates by soil types, weed species, and weed or crop growth stage. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

### Uses with Other Products (Tank Mixes)

If this product is used in combination with any other product except as specifically recommended in writing by INNVICTIS, to the extent consistent with applicable law, INNVICTIS shall have no liability for any loss, damage, or injury arising out of its use in any such combination not so specifically recommended. If used in combination recommended by INNVICTIS, to the extent consistent with applicable law, the liability of INNVICTIS shall in no manner extend to any damage, loss, or injury not directly caused by the inclusion of the INNVICTIS product in such combination use, and in any event shall be limited to return of the amount of the purchase price of the product. Always perform a mixing test to check the compatibility of *PAVILION 3.3 EC HERBICIDE* with all potential tank mix partners.

### Mixing Instructions

1. Fill tank 1/2 to 3/4 full with clean water or liquid fertilizer and agitate. Prior to mixing *PAVILION 3.3 EC HERBICIDE* or *PAVILION 3.3 EC HERBICIDE* tank mixtures in liquid fertilizer, refer to appropriate label sections for specified uses in liquid fertilizer, application instructions, and compatibility determinations.

**Note:** *PAVILION 3.3 EC HERBICIDE* will not mix in high salt formulation fertilizers, such as 10-34-0. When utilizing high salt formulation fertilizers as the spray carrier, use one of the following:

- (a) Pre-slurry *PAVILION 3.3 EC HERBICIDE* in water prior to adding to tank; use 1:1 ratio of water to *PAVILION 3.3 EC HERBICIDE*.
  - (b) Add water to fertilizer solution prior to adding *PAVILION 3.3 EC HERBICIDE*. The amount of water should be equal to or greater than the amount of *PAVILION 3.3 EC HERBICIDE* to be used.
2. ***PAVILION 3.3 EC HERBICIDE* Alone**  
When using *PAVILION 3.3 EC HERBICIDE* alone, add *PAVILION 3.3 EC HERBICIDE* to the partially filled tank while agitating and then fill the remainder of the tank with water or liquid fertilizer.
  3. ***PAVILION 3.3 EC HERBICIDE* Tank Mixes**  
Add the tank mixture ingredients in the order listed below prior to adding *PAVILION 3.3 EC HERBICIDE* (for tank mixtures with 2,4-DB, paraquat or glyphosate, see mixing instructions at the end of this section):
    - (a) **Wettable Powder (WP) formulations** - Make a slurry of the WP in water (1:2 ratio). Add the slurry slowly into the partially filled tank while agitating.
    - (b) **Dry Flowable (DF)/Water Dispersible Granule (WDG) formulations** - Add the granules to the partially filled tank while agitating. Make a slurry of the granules in water before adding to liquid fertilizer.
    - (c) **Flowable (F) formulations** - Add the F formulation to the partially filled tank while agitating.
    - (d) **Water Soluble Concentrate (WSC) formulations** - Add the WSC formulation to the partially filled tank while agitating.
    - (e) **Emulsifiable Concentrate (EC) formulations** - Add the EC formulation to the partially filled tank while agitating.

After complete mixing, add *PAVILION 3.3 EC HERBICIDE* to the tank.

- (f) **Note: For tank mixes including 2,4-DB, paraquat or glyphosate:** After complete mixing of *PAVILION 3.3 EC HERBICIDE*, continue filling the sprayer with water and add 2,4-DB or paraquat or glyphosate near the end of the filling process.  
If paraquat is included in the tank mixture, add 8 ozs of non-ionic surfactant per 100 gallons of total spray mixture as the last ingredient in the tank. Fill the remainder of the tank with water or liquid fertilizer while agitating.
4. Thorough and continuous sprayer-tank agitation must be maintained during mixing and spraying of *PAVILION 3.3 EC HERBICIDE*. If the spray mixture is allowed to settle for any period of time, thorough agitation is essential to resuspend the mixture before spraying is resumed. Continue agitation while spraying.

### Cleaning Spray Equipment

Clean application equipment thoroughly by using a strong detergent or commercial spray cleaner according to the manufacturer's directions, and then triple rinsing the equipment before and after applying this product.

### Precautions:

- *PAVILION 3.3 EC HERBICIDE* is most effective in controlling weeds mechanically incorporated or when incorporated into the weed germination zone by adequate rainfall or overhead irrigation after application.
- When using tank mixtures with *PAVILION 3.3 EC HERBICIDE*, always read the companion product label(s) to determine the specific use rates by soil types, weed species, and weed or crop growth stage. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.
- In the event of a crop loss due to adverse weather conditions or other reasons, any crop registered for a preplant incorporated application of *PAVILION 3.3 EC HERBICIDE* can be replanted without adverse effects the same year (see **Crop Specific Information** for exceptions). If replanting is necessary, do not work the soil deeper than the treated zone.
- Refer to **Crop-specific Information** for crop-specific preharvest intervals and feeding and grazing restrictions.

### Restrictions

- Do not exceed the maximum labeled rate for any soil type.
- *PAVILION 3.3 EC HERBICIDE* will not control established weeds. Destroy emerged weeds prior to application.

## CROP ROTATION DIRECTIONS

- Use of *PAVILION 3.3 EC HERBICIDE* in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors, such as arid conditions, make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible. Soil characteristics and environmental conditions which may contribute to crop stress that may be accentuated by the use of *PAVILION 3.3 EC HERBICIDE* include: coarse soils, compaction, high salinity, eroded knolls/hilltops, cold and/or wet soils, drought, and heavy rainfall soon after application.
- When *PAVILION 3.3 EC HERBICIDE* is used in tank mix or sequential combinations, refer to labels of other herbicides for additional rotational crop restrictions. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.
- **Restrictions for rotational cropping after the use of *PAVILION 3.3 EC HERBICIDE* are dependent on the application use rate of *PAVILION 3.3 EC HERBICIDE* in the primary crop. Read the following restrictions thoroughly to determine the rotational crops for each specific situation, according to application use rate.**

### I. Rotational Crop Restrictions Following Applications of *PAVILION 3.3 EC HERBICIDE* to Field and Row Crops

#### 1. Application Rate less than or equal to 4.8 pts/A (2.0 lbs ai/A):

a. Crops which are labeled for preplant incorporated application may be planted the same season in which *PAVILION 3.3 EC HERBICIDE* was applied.

##### b. Sugar beets, Red beets and Spinach

To avoid crop injury, do not plant sugar beets, red beets or spinach for 12 months following a spring application of *PAVILION 3.3 EC HERBICIDE* or 14 months following a fall application of *PAVILION 3.3 EC HERBICIDE*.

These crops should not be planted for 18 months following a spring application of *PAVILION 3.3 EC HERBICIDE* or 20 months following a fall application of *PAVILION 3.3 EC HERBICIDE* if rainfall or irrigation was not sufficient to produce a crop.

To ensure thorough mixing of soil prior to planting sugar beets, red beets and spinach, land should be plowed using a moldboard plow to a depth of 12 inches.

##### c. Proso millet, Sorghum (milo), and Annual or Perennial grass crops or mixtures

Proso millet, sorghum (milo), and annual or perennial grass crops or mixtures should not be planted for 10 months after a spring application of *PAVILION 3.3 EC HERBICIDE* or 12 months after a fall application of *PAVILION 3.3 EC HERBICIDE* except in the following conditions:

In the states of **Minnesota, North Dakota** and **South Dakota**, these crops should not be planted for 18 months following a spring application of *PAVILION 3.3 EC HERBICIDE* or 21 months following a fall application of *PAVILION 3.3 EC HERBICIDE*.

To avoid the possibility of crop injury in areas that receive less than 20 inches of rainfall or irrigation to produce a crop, these crops should not be planted for 18 months following a spring application of *PAVILION 3.3 EC HERBICIDE* or 20 months following a fall application of *PAVILION 3.3 EC HERBICIDE* if rainfall or irrigation was not sufficient to produce a field or row crop.

##### d. Wheat and Barley

Wheat and barley may be planted 4 months after an application of *PAVILION 3.3 EC HERBICIDE*, except under the following conditions:

If less than 12 inches of rainfall or overhead irrigation was received between application and rotational crop planting, wheat should not be planted before 12 months after a spring application of *PAVILION 3.3 EC HERBICIDE* or 14 months after a fall application of *PAVILION 3.3 EC HERBICIDE*.

In dryland areas and/or areas where irrigation is necessary to produce the crop treated with *PAVILION 3.3 EC HERBICIDE*, do not plant winter wheat or barley as a follow crop if crop failure/destruction occurs and land is fallowed during the summer.

##### e. All Other Rotational Crops Not Specifically Addressed Above

Crops, other than those to which *PAVILION 3.3 EC HERBICIDE* may be applied as a preplant incorporated treatment, may be planted the year following application of *PAVILION 3.3 EC HERBICIDE*, except under the following condition:

If rainfall or irrigation was not sufficient to produce a crop, delay planting for 18 months following a spring application of *PAVILION 3.3 EC HERBICIDE* or 20 months following a fall application of *PAVILION 3.3 EC HERBICIDE*.

#### 2. Application Rate greater than 4.8 pts/A (2.0 lbs ai/A):


In the growing season following application of *PAVILION 3.3 EC HERBICIDE* to field and row crops at greater than 4.8 pts/A, plant only those crops for which *PAVILION 3.3 EC HERBICIDE* is labeled for preplant incorporated treatment or crop injury may occur. **DO NOT** plant other crops for 24 months.

## II. Rotational Crop Restrictions Following Applications of PAVILION 3.3 EC HERBICIDE to Orchard, Grove, and Vineyard Crops

In the growing season following application of PAVILION 3.3 EC HERBICIDE to bearing fruit and nut trees, plant only those crops for which PAVILION 3.3 EC HERBICIDE is labeled for preplant incorporated treatment or crop injury may occur.

Do not rotate to other crops (except for nut crops, fruit trees, or grapes) for 24 months following a PAVILION 3.3 EC HERBICIDE application to fruit or nut trees.

### Use Area



**Note:** Southern States includes Hawaii

#### CROP-SPECIFIC INFORMATION

**Crop Injury Disclaimer:** PAVILION 3.3 EC HERBICIDE use may result in crop injury, loss or damage to certain crops under a number of conditions, including but not limited to agronomic, cultural, mechanical, and environmental. Numerous risks of loss or damage to certain crops may be associated with the use of PAVILION 3.3 EC HERBICIDE even when directions for use are followed completely. The user or grower should take all such risks into consideration before deciding to apply the product. **INNICTIS recommends testing on a small portion of the target crop to determine if damage is likely to occur.** Each grower who is considering the product for such use should test PAVILION 3.3 EC HERBICIDE in order to determine its suitability. A grower should use PAVILION 3.3 EC HERBICIDE only to the extent that in his sole opinion the benefit of PAVILION 3.3 EC HERBICIDE use outweighs the potential injury to the grower's crop.

In addition, many factors can affect crop growth and/or yield, including but not limited to, insects, diseases, weed competition, poor seed quality, improper planting depth, mechanical cultivation, poor weather (such as freezing or excessive wind, rain, heat, or cold), lack of or excessive moisture, crusting, fertility, or hardpans. Risk of loss or damage to crops may be associated with the use of PAVILION 3.3 EC HERBICIDE and contribute to poor stands due to failure of crop to emerge, swelling of roots or other below-ground plant parts, less vigorous plant growth and development, and reduction in yield potential.

PAVILION 3.3 EC HERBICIDE may also cause injury to sensitive rotational crops.

**ALFALFA**  
**(Grown for Forage, Hay or Seed Production)**

*PAVILION 3.3 EC HERBICIDE* may be applied by ground, air, chemigation, flooded basin irrigation systems, or on dry bulk fertilizer.

**Use Directions**

**Established Alfalfa for Forage/Hay** (defined as alfalfa planted in the fall or spring that has gone through a first cutting/mowing):

Uniformly apply *PAVILION 3.3 EC HERBICIDE* at a broadcast rate of 1.2 to 4.8 quarts per acre prior to weed emergence. Applications can be made in the fall after the last mowing/cutting, during winter dormancy, in the spring, or between cuttings. Applications should be made prior to the alfalfa reaching 6 inches in regrowth.

**Established Alfalfa Grown for Seed Production** (defined as alfalfa planted in the fall or spring that has gone through a summer season of cutting/mowing):

Uniformly apply *PAVILION 3.3 EC HERBICIDE* at a broadcast rate of 1.2 to 4.8 quarts per acre prior to weed emergence in one of the following ways:

1. Apply to dormant established alfalfa.
2. Apply before alfalfa exceeds 10 inches in height after first mowing/beatng.
3. Once the alfalfa reaches 10 inches in height or if the alfalfa has been mowed/beaten two or more times, *PAVILION 3.3 EC HERBICIDE* must be applied with drop nozzles directing the spray so that there is little to no contact with the foliage.

**Seedling Alfalfa** (defined as alfalfa planted in the fall or spring which has not gone through a cutting/mowing):

Uniformly apply *PAVILION 3.3 EC HERBICIDE* at a broadcast rate of 1.2 to 2.4 pints per acre prior to weed emergence. Applications can be made once the seedling alfalfa has reached the 2nd trifoliate stage of growth. Applications should be made prior to the alfalfa reaching 6 inches in growth.

**Alfalfa Stand Establishment:**

Apply *PAVILION 3.3 EC HERBICIDE* at a broadcast rate of 1.2 to 1.8 pints per acre as a preplant incorporated or preemergence treatment in direct-seeded alfalfa. Some crop stand reduction and stunting may occur with this use of *PAVILION 3.3 EC HERBICIDE*; however, reduced weed competition will allow establishment of a quality stand. Use the lower rates on coarse-texture soil or in lower rainfall areas (receiving less than 20 inches of rainfall and irrigation a year).

**Preplant incorporated:** Uniformly incorporate *PAVILION 3.3 EC HERBICIDE* into the top 2 to 3 inches of the final seedbed prior to planting.

**Preemergence:** Apply directly after drill seeding alfalfa. Alfalfa should be planted into a seedbed that is firm and free of clods.

**Chemigation Applications**

*PAVILION 3.3 EC HERBICIDE* may be applied through sprinkler irrigation systems. Follow all directions, special instructions and precautions in the section covering **Chemigation in Spraying Instructions**.

**Flooded Basin Irrigation Systems**

*PAVILION 3.3 EC HERBICIDE* may be applied in flooded basin irrigation systems. Follow all directions, special instructions and precautions in the section covering **Flooded Basin Irrigation in Spraying Instructions**.

**Precautions**

- Follow all precautions and restrictions on the labels of all products applied in combination with *PAVILION 3.3 EC HERBICIDE*. Always follow the most restrictive label.
- Some stunting and chlorosis of the alfalfa may occur with postemergence applications.
- Applications made after the alfalfa exceeds 6 inches in height may result in poor weed control due to possible reduced spray coverage to the soil.

**Restrictions**

- Do not exceed 4.8 quarts of *PAVILION 3.3 EC HERBICIDE* per acre in any one crop season.
- Do not apply *PAVILION 3.3 EC HERBICIDE* less than 50 days prior to alfalfa harvest for forage or hay.
- Do not apply *PAVILION 3.3 EC HERBICIDE* less than 90 days prior to alfalfa harvest for seed.

**BEARING FRUIT AND NUT TREES**

*PAVILION 3.3 EC HERBICIDE* may be applied in the following individual crops within the fruit tree and tree nut crop groupings:

Citrus Fruit Crop Grouping		
Calamondin	Kumquat	Orange (sweet and sour)
Citrus citron	Lemon	Pummelo
Citrus hybrids	Lime	Satsuma mandarin
Grapefruit	Mandarin (tangerine)	Tangelo

Tree Nuts Crop Grouping		
Almond	Chestnut	Pecan
Beech nut	Chinquapin	Pistachio
Brazil Nut	Filbert (hazelnut)	Walnut
Butternut	Hickory nut	
Cashew	Macadamia nut	

**Pome Fruits Crop Grouping**

Apple	Mayhaw	Pear, oriental
Crabapple	Pear	Quince
Loquat		

**Stone Fruits Crop Grouping**

Apricot	Peach	Plum, Japanese
Aprium	Plum	Plumcot
Cherry, sweet	Plum, chicksaw	Pluot
Cherry, tart	Plum, Damson	Prune
Nectarine		

**Other Fruit Trees**

Pomegranate	Juneberry
-------------	-----------

**Use Directions**

*PAVILION 3.3 EC HERBICIDE* may only be applied by ground, chemigation or flooded basin irrigation systems.

*PAVILION 3.3 EC HERBICIDE* may be applied either in a single application or sequentially with an interval of 30 days or more. Apply *PAVILION 3.3 EC HERBICIDE* at between 2.4 to 4.8 quarts per acre (depending on desired length of control, see chart below) per application.

**Use Rate per Acre:**

Short-term control	2.4 quarts
Long-term control	4.8 quarts

**Ground Applications**

*PAVILION 3.3 EC HERBICIDE* may be applied surface incorporated of (surface) preemergence.

Apply *PAVILION 3.3 EC HERBICIDE* as a broadcast or banded treatment using ground equipment before weed emergence. Apply the spray directly to the ground beneath the trees and/or in areas between rows.

**Chemigation Applications**

*PAVILION 3.3 EC HERBICIDE* may be applied through sprinkler irrigation systems. Follow all recommendations, special instructions and precautions in the general section covering **Chemigation in Spray Instructions**.

**Flood Basin Irrigation Systems**

*PAVILION 3.3 EC HERBICIDE* may be applied in flood basin irrigation systems. Follow all recommendations, special instructions and precautions in the general section **Flood Basin Irrigation in Spraying Instructions**.

**Restrictions**

- Do not apply more than 4.8 quarts of *PAVILION 3.3 EC HERBICIDE* per acre per year in pome, stone and other fruit trees
- Do not apply more than 7.2 quarts of *PAVILION 3.3 EC HERBICIDE* per acre per year in citrus and nut trees.
- Do not apply by air.
- Do not feed forage or graze livestock in treated groves or orchards.
- Do not apply within 1 day of harvest of citrus fruit.
- Do not apply within 60 days of harvest of pome and stone fruit or other tree fruit.
- Do not apply within 60 days of harvest of nuts, except almonds.
- Do not apply within 120 days of harvest of almonds
- Do not apply to newly seeded nursery stock.
- Do not apply over the top of trees with leaves or buds or fruit. Contact by the spray mixture with leaves, shoots or buds may cause injury.
- Do not apply *PAVILION 3.3 EC HERBICIDE*-treated irrigation water over top of trees with leaves or buds or fruit. Contact with leaves, shoots or buds by spray mixture may cause injury.

## CARROTS

*PAVILION 3.3 EC HERBICIDE* may be applied by ground, air or chemigation.

### Use Directions

**Preemergence** - Make a single broadcast application by ground or by air or by chemigation at 2.4 pints per acre of *PAVILION 3.3 EC HERBICIDE* as a postplant treatment prior to emergence of the crop and before weed emergence. Apply as a preemergence treatment within 2 days after planting.

**Layby** - *PAVILION 3.3 EC HERBICIDE* may be applied only by ground equipment at layby (last mechanical cultivation) at 2.0 pints per acre as a directed spray to the soil between rows.

### Layby Precautions:

- *PAVILION 3.3 EC HERBICIDE* should be applied prior to weed emergence.
- Emerged weeds will not be controlled by this treatment.

### Chemigation Applications

*PAVILION 3.3 EC HERBICIDE* may be applied through sprinkler irrigation systems. Follow all recommendations, special instructions and precautions in the general section covering Chemigation in **Spraying Instructions**.

### Restrictions

- Do not apply more than 2.4 pints per acre per season.
- Do not apply within 60 days of harvest.
- Do not feed forage or graze livestock in treated fields.
- Do not apply as a broadcast spray over top of carrots or crop injury may result.
- Do not allow the spray to contact carrot plants or injury may occur.
- Do not allow *PAVILION 3.3 EC HERBICIDE*-treated irrigation water to contact carrot plants.
- Do not apply layby applications by chemigation or by air.
- Do not apply tank mixtures through any type of irrigation system unless the label instructions on chemigation of all products are followed.

## CARROTS GROWN FOR SEED PRODUCTION

*PAVILION 3.3 EC HERBICIDE* may be applied only by layby.

### Use Directions

**Last Cultivation (Layby)** - Apply *PAVILION 3.3 EC HERBICIDE* following the last normal mechanical cultivation (layby) at a rate of 1.2 to 4.8 pints per acre (on a broadcast basis). Uniformly apply as a directed spray to the soil between rows.

Layby applications can be applied to carrots previously treated with herbicides registered in/on carrots. Consult the labels of those herbicides for suggested treatments, rates to be used, and precautions or restrictions for use in carrots and for follow crop restrictions.

### Precautions

- Use protective shields to avoid contact with carrot foliage.
- Use properly calibrated and accurate nozzles and equipment.

### Restrictions

- Do not apply as a broadcast spray over top of carrots or crop injury may result.
- Do not apply layby applications by chemigation or by air.
- Do not apply within 60 days of carrot seed harvest.
- Do not feed, forage or graze livestock in treated fields.
- Do not harvest carrots for food or feed use.
- Do not allow the spray to contact carrot plants or injury may occur.

### SPECIAL CROP USE INSTRUCTIONS

The pesticide applicator, the producer of the crop, and the seed conditioner must be aware that use of this product according to this labeling is deemed a nonfeed/nonfood use. If the applicator of this pesticide is not the producer, the applicator should provide a copy of this labeling to the producer of the crop. Producers of this crop who use this product, or cause the product to be used on a field they operate, should provide a copy of this pesticide label to the seed conditioner.

Consequently, no portion of this carrot seed crop, including but not limited to green chop, hay, pellets, meal, whole seed, cracked seed, roots, bulbs, foliage and seed screenings, may be used or distributed for food or feed purposes.

Processed carrot seed from a field treated with this product must bear a specific tag or conspicuous container labeling, or if shipped in bulk, on the shipment invoice or bill of lading, with the following statement: "Not for human consumption or animal feed." All seed screenings from seed processing shall be disposed of in such a manner that the screenings cannot be distributed or used for human food or animal feed purposes.

The seed conditioner shall keep records of screening disposal for three years from the date of disposal and shall furnish the records immediately upon request. Conditioner disposal records shall consist of documentation of on-farm disposal, disposal at a controlled dumpsite, incinerator, compostor or other equivalent disposal site and shall include the lot numbers, amount of material disposed of, the grower(s), and the date of disposal.

**CORN**  
(Field, Pop, Seed, Sweet)

*PAVILION 3.3 EC HERBICIDE* may be applied by ground, air or chemigation.

*PAVILION 3.3 EC HERBICIDE* may be applied in conventional, minimum, or no-till as a preemergence, postemergence, or postemergence incorporated (CULTI-SPRAY) application in field corn.

*PAVILION 3.3 EC HERBICIDE* may be applied in conventional tillage as a preemergence or postemergence application in sweet corn, seed corn, or popcorn.

**Regardless of tillage system, plant corn at least 1-1/2 inches deep and completely cover with soil.**

*PAVILION 3.3 EC HERBICIDE* or *PAVILION 3.3 EC HERBICIDE* tank mix combination treatments are most effective in controlling weeds when adequate rainfall or overhead irrigation is received within 7 days after application. If cultivation is necessary because of soil crusting or weed germination, use shallow tillage and make certain corn seeds are below the tilled area.

**Precautions**

- In conventional tillage systems, plant into a seedbed that is firm and free of clods and trash. Use only where adequate tillage is practiced to provide good soil coverage of the corn seed.
- In no-till systems, utilize a no-till planter that is capable of planting through crop residue. The use of no-till planters under conditions that do not allow good soil coverage of the corn seed can result in reduced crop stand or injury if *PAVILION 3.3 EC HERBICIDE* contacts the germinating corn seed. Check equipment to ensure good seed coverage.

**Additional Weeds Controlled:** In addition to the weeds listed in **Table 1**, *PAVILION 3.3 EC HERBICIDE* will control the following weeds in corn with CULTI-SPRAY application: wild proso millet and shattercane.

**Use Directions**

**Preemergence** - Apply after planting but before weeds and crop emerge.

**Postemergence** - Apply postemergence until field corn is 30 inches tall (20 to 24 inches tall for pop, seed and sweet corn) or in the V8 growth stage, whichever is more restrictive. If the corn canopy prevents applications from reaching the soil, use drop nozzles and apply as a directed spray.

**CULTI-SPRAY** - Apply *PAVILION 3.3 EC HERBICIDE* alone or *PAVILION 3.3 EC HERBICIDE* plus atrazine when field corn is at least 4 inches tall until last cultivation (layby). *PAVILION 3.3 EC HERBICIDE* plus atrazine must be applied before the field corn reaches 12 inches in height.

Under situations of low rainfall or soil moisture when deep germinating weeds such as shattercane or field sandour are anticipated, mechanical incorporation will provide best results. If cultivation is needed after application and incorporation of *PAVILION 3.3 EC HERBICIDE*, the depth of cut should be no deeper than the depth of cut used to incorporate.

**Chemigation Applications**

*PAVILION 3.3 EC HERBICIDE* may be applied through sprinkler irrigation systems. Follow all directions, special instructions and precautions in the section covering **Chemigation in Spraying Instructions**.

**Use Rates**

**Preemergence or Postemergence Applications**

Soil Texture	Organic Matter		
	<1.5% (pts/A)	1.5 to 3.0% (pts/A)	>3.0% (pts/A)
Coarse	1.8 to 2.4	2.4 to 3.6	3.6
Medium	2.4 to 3.6	3.6	3.6 to 4.8
Fine	2.4 to 3.6	3.6 to 4.8	3.6 to 4.8

**CULTI-SPRAY Applications - Field Corn ONLY**

Soil Texture	Southern States <sup>1</sup> (pts/A)	Northern States <sup>1</sup> (pts/A)
Coarse	1.2 to 1.8	1.8 to 2.4
Medium	1.8 to 2.4	2.4 to 3.6
Fine	1.8 to 3.6	2.4 to 3.6

<sup>1</sup>See **Use Area** map for specific states.



## Restrictions

- Do not apply *PAVILION 3.3 EC HERBICIDE* in reduced, minimum or no-till sweet corn, seed corn or popcorn.
- Do not apply *PAVILION 3.3 EC HERBICIDE* in no-till in California.
- Do not apply preplant incorporated.
- Do not apply postemergence in liquid fertilizer.
- Livestock can graze or be fed forage from treated corn after 21 days following application.
- Do not exceed one application per crop season at the highest rate per acre for any given soil type and application method.
- Do not exceed 1.2 lbs ai per acre of atrazine, as specified on the atrazine label.

## COTTON

*PAVILION 3.3 EC HERBICIDE* may be applied by ground, air, or chemigation in conventional, minimum, stale seedbed, or no-till as a preplant surface, preplant incorporated, preemergence, or layby application in cotton.

Preplant surface, preemergence, and layby treatments are most effective in controlling weeds when adequate rainfall or overhead irrigation is received within 7 days after application. A shallow cultivation is recommended if soil crusting or soil compaction occurs. If weeds begin to germinate or adequate moisture is not received within 7 days after application, use shallow tillage (rotary hoe or light harrow) and make sure cotton seeds are below tilled area. The use of a postemergence herbicide treatment may be required to control weed escapes at planting or following cotton emergence.

**Additional Weeds Suppressed:** In addition to the weeds listed in **Table 1**, *PAVILION 3.3 EC HERBICIDE* will suppress Russian thistle in the state of Arizona.

### Use Directions

**Preplant Surface** - Apply *PAVILION 3.3 EC HERBICIDE* up to 15 days prior to planting. Apply *PAVILION 3.3 EC HERBICIDE* tank mixes and sequential programs as specified under the tank mix section.

**Preplant Incorporated** - Apply *PAVILION 3.3 EC HERBICIDE* up to 60 days prior to planting and incorporate within 7 days of application. Apply *PAVILION 3.3 EC HERBICIDE* tank mixes and sequential programs as specified under the tank mix section.

**Preemergence** - Apply *PAVILION 3.3 EC HERBICIDE* at planting or up to 2 days after planting. Apply to a seedbed that is firm and free of clods. Apply *PAVILION 3.3 EC HERBICIDE* tank mixes and sequential programs as specified under the tank mix section.

**Preplant Incorporated followed by Preemergence** - Apply *PAVILION 3.3 EC HERBICIDE* up to 60 days prior to planting and incorporate within 7 days of application. Apply overlay application of *PAVILION 3.3 EC HERBICIDE* at planting or up to 2 days after planting. Total amount of *PAVILION 3.3 EC HERBICIDE* applied per acre cannot exceed the highest labeled rate for a given soil type. Preplant incorporated and preemergence applications of *PAVILION 3.3 EC HERBICIDE* may be applied with the labeled tank mix herbicide(s).

**Layby Application (at last cultivation)** - Apply *PAVILION 3.3 EC HERBICIDE* directly to the soil between rows as a directed spray following the last normal cultivation (layby). Layby applications can be applied in cotton previously treated with *PAVILION 3.3 EC HERBICIDE* or any herbicide(s) registered for use in cotton. Consult the labels of those herbicides for suggested treatments, rates to be used, and precautions or restrictions for use in cotton, and for follow-crop restrictions. The total amount of *PAVILION 3.3 EC HERBICIDE* applied per acre per season cannot exceed the highest labeled rate for a given soil type.

Glyphosate-containing products may be applied with *PAVILION 3.3 EC HERBICIDE* at layby in cotton with the **Roundup Ready<sup>®</sup>** gene.

### Layby Precautions

- Avoid contact of the spray to the non-woody portion of cotton stems and to cotton foliage or serious crop injury can result.
- To reduce the potential for crop injury caused by herbicide contact with cotton foliage and stems, use protective shields when conditions favoring spray drift occur.

### Layby Restrictions

- Do not apply as a broadcast spray over the top of the cotton or serious crop injury can result.
- Do not apply glyphosate-containing products at layby on non-Roundup Ready cotton.
- Do not apply *PAVILION 3.3 EC HERBICIDE* and glyphosate tank mix as a broadcast spray over the top of cotton or CROP INJURY MAY RESULT.

### Chemigation Applications

*PAVILION 3.3 EC HERBICIDE* may be applied through sprinkler irrigation systems. Follow all directions, special instructions and precautions in the section covering **Chemigation in Spraying Instructions**.

**Fall Application** - *PAVILION 3.3 EC HERBICIDE* may be applied for weed control in cotton in the fall after October 15 (up to 140 days prior to planting cotton) in Arizona, California, Louisiana, New Mexico, Mississippi, Oklahoma and Texas. Apply *PAVILION 3.3 EC HERBICIDE* at the broadcast rate of 2.4 pints per acre on coarse or medium soils and 3.6 pints per acre on fine soils.

### Use Rates

Soil Texture	Conventional or Minimal Tillage (pts/A)	No-Till <sup>2</sup> (pts/A)
Coarse	1.2 to 2.4 <sup>1</sup>	1.8 to 2.4
Medium	1.8 to 2.4	2.4 to 3.6
Fine	2.4 to 3.6	3.6 to 4.8

<sup>1</sup>Do not exceed 1.8 pts/A on coarse-textured soils in California.

<sup>2</sup>Not recommended for soils with more than 3% organic matter.

### Restrictions

- Do not apply *PAVILION 3.3 EC HERBICIDE* in no-till in California.
- Preharvest Interval (PHI) is 60 days between the last *PAVILION 3.3 EC HERBICIDE* application and harvest.
- Do not feed forage or graze livestock in treated cotton fields.
- Do not exceed the highest seasonal rate per acre for any given soil type.

### EDIBLE BEANS

#### (Dry, Lima, Snap, Chickpeas (Garbanzo Beans), Southern Peas (Cowpeas), and Sweet Lupines)

*PAVILION 3.3 EC HERBICIDE* may only be applied (fall) preplant surface or preplant incorporated in chickpeas (garbanzo beans), dry beans, lima beans, snap beans, and Southern peas (cowpeas). *PPAVILION 3.3 EC HERBICIDE* may be applied (fall) preplant surface or preplant incorporated or preemergence in sweet lupines.

### Use Directions

**Preplant Incorporated** - Apply up to 60 days prior to planting and incorporate within 7 days of application.

**Preemergence** - Apply only to sweet lupines at planting or up to 2 days after planting. Apply to a seedbed that is firm and free of clods.

### Use Rates

Soil Texture	Southern States <sup>1</sup> (pts/A)	Northern States <sup>1</sup> <3% Organic Matter > 3% (pts/A)	
		Coarse	1.8
Medium	2.4	3.0	3.6
Fine	3.6	3.6	3.6

<sup>1</sup>See **Use Area** map for specific states.

**Fall Applications** - Fall preplant surface and preplant incorporated applications may be made in North Dakota, South Dakota, Minnesota, Oregon, Washington, Montana, Idaho, and Wyoming only. Rainfall or irrigation is required for incorporation and activation. Unpredictable weed control can be expected since factors such as length of time between application and planting as well as uncontrollable weather factors will determine herbicide activity and longevity.

Apply *PAVILION 3.3 EC HERBICIDE* and incorporate (rainfall, irrigation or mechanically) in late fall prior to planting edible beans [chickpeas (garbanzo beans)], dry beans (such as navy, great northern, red kidney, black turtle, cranberry, and small white type), lima beans, snap beans, Southern peas (cowpeas), and sweet lupines the following spring. Apply *PAVILION 3.3 EC HERBICIDE* in the late fall when soil temperatures are 45° F or below but before the ground freezes.

### Preplant Surface and Preplant Incorporated (Fall Application) Use Rates

Soil Texture	Broadcast Rate <3% Organic Matter (pts/A)	Broadcast Rate >3% Organic Matter (pts/A)
Coarse	1.2 to 2.4	2.4
Medium	1.8 to 3.0	3.0 to 3.6
Fine	2.4 to 3.6	3.6

<sup>1</sup>For use in North Dakota, South Dakota, Minnesota, Oregon, Washington, Montana, Idaho and Wyoming ONLY.

### Restrictions

- Do not feed lupine hay and forage or graze livestock in treated lupine fields.
- Do not apply *PAVILION 3.3 EC HERBICIDE* more than once per cropping season.
- Do not apply in any type of irrigation system.
- For fall applications, do not apply when the air temperature is below 45°F.

## FORAGE LEGUMES

*PAVILION 3.3 EC HERBICIDE* may be used in forage legumes used as a cover crop in federal set-aside or conservation reserve program areas.

### Use Directions

*PAVILION 3.3 EC HERBICIDE* may be applied preplant incorporated or preemergence for weed control in legume cover crops.

### Use Rates Preplant Incorporated or Preemergence

Soil Texture	Broadcast Rate (pts/A)
Coarse	1.2 to 1.8
Medium	1.8 to 2.4
Fine	2.4 to 3.0

### Precautions

- Some stand reduction of the legume cover crop may occur with this use. Consult local county extension service or the local ASC committee for recommended cover crops.
- If loss of cover crop occurs due to adverse weather conditions, any crop registered for *PAVILION 3.3 EC HERBICIDE* preplant incorporated use can be replanted the same year into *PAVILION 3.3 EC HERBICIDE*-treated soil without adverse effects.
- If replanting is necessary, do not rework the soil deeper than the *PAVILION 3.3 EC HERBICIDE*-treated zone.
- The cover crop residue should ultimately be destroyed by tillage or left on the surface to retard erosion or as directed by the local ASC committee.

### Restrictions

- Do not feed or graze legume cover crops established following *PAVILION 3.3 EC HERBICIDE* application.

## FRUITING VEGETABLES

*PAVILION 3.3 EC HERBICIDE* may be applied to the following fruiting vegetables: tomato, eggplant, groundcherry (*Physalis* spp.), pepper (includes bell pepper, chili pepper, cooking pepper, pimento, sweet pepper), pepino, tomatillo.

### Use Directions

Uniformly apply *PAVILION 3.3 EC HERBICIDE* only by ground or air as a broadcast preplant incorporated application, or as a broadcast preplant surface application prior to transplanting fruiting vegetables, or as a post-directed application to transplanted or established direct-seeded fruiting vegetables.

*PAVILION 3.3 EC HERBICIDE* can be applied as a post-directed spray on the soil at the base of the plant, beneath plants, and between rows. Following the post-directed spray and when sufficient rainfall or irrigation does not occur to activate the herbicide, mechanically incorporate at the time of blocking and thinning or at "layby." *PAVILION 3.3 EC HERBICIDE* should be applied prior to weed emergence.

### Use Rates

Soil Texture	Broadcast Rate (pts/A)
Coarse	1.2 to 1.8
Medium	1.8 to 2.4
Fine	1.8 to 3.6

### Precautions

- Avoid direct contact with foliage or stems.
- Be sure roots of transplants are established.
- Emerged weeds will not be controlled by this treatment.

### Restrictions

- Do not apply more than 3.6 pints per acre per season.
- Do not apply within 70 days of harvest.
- Do not allow *PAVILION 3.3 EC HERBICIDE*-treated soil to come in contact with transplant area.
- Do not apply if row is to later be covered with plastic.
- Do not apply prior to direct-seeded fruiting vegetables.
- Do not apply postemergence over the top of or to foliage of fruiting vegetables as severe injury may occur.

## GARLIC

*PAVILION 3.3 EC HERBICIDE* may be applied preemergence, postemergence, or split application by ground, air, or chemigation.

### Use Directions

**Preemergence** - After planting but before crop and weeds emerge.

**Postemergence** - 1st to 5th true-leaf growth stage.

**Split Application** - At both preemergence and postemergence timings.

### Chemigation Applications

*PAVILION 3.3 EC HERBICIDE* may be applied through sprinkler irrigation systems. Apply between the 2nd and 9th true-leaf stage (2nd to 6th true-leaf stage in California). Follow all directions, special instructions and precautions in the section covering **Chemigation** in **Spraying Instructions**.

### Use Rates

Soil Texture	Broadcast Rate (pts/A)
Coarse	1.8
Medium	2.4
Fine	3.6

### Restrictions

- Do not exceed 3.6 pints per acre per crop (except Idaho, Oregon, and Washington).
- Do not apply within 60 days of harvest in California and within 45 days of harvest in all other states.
- Do not feed or graze these crops.
- Do not irrigate in excess of 0.5 inch of water.

## GRAIN SORGHUM

Uniformly apply *PAVILION 3.3 EC HERBICIDE* in water by ground equipment or by aircraft.

*PAVILION 3.3 EC HERBICIDE* may be applied as a postemergence incorporated (CULTI-SPRAY) application in grain sorghum grown in all states. In addition, *PAVILION 3.3 EC HERBICIDE* may be applied early postemergence in grain sorghum grown in states east of the Mississippi River and in Arkansas, eastern Texas, Louisiana, and the Missouri "boothel."

**Additional Weeds Controlled:** In addition to the weeds listed in **Table 1**, *PAVILION 3.3 EC HERBICIDE* as a CULTI-SPRAY application will control the following weeds in grain sorghum: wild proso millet and shattercane.

### Use Directions

**CULTI-SPRAY:** *PAVILION 3.3 EC HERBICIDE* treatments can be applied from the 4-inch growth stage to as late as the last cultivation (layby) of grain sorghum. See specific directions for (CULTI-SPRAY) application under **Application Instructions**.

**Early Postemergence:** For use only in states east of the Mississippi River plus Arkansas, eastern Texas, Louisiana, and the "boothel" of Missouri.

The seedbed should be firm and free of clods and trash. Use only where adequate tillage is practiced to provide good seed coverage. Plant grain sorghum at least 1-1/2 inches deep to ensure good seed coverage.

### Use Rates

#### CULTI-SPRAY Application

Soil Texture	Southern States <sup>1</sup> (pts/A)	Northern States <sup>1</sup> (pts/A)
Coarse	1.8	2.4
Medium	2.4	3.6
Fine	3.6	3.6

<sup>1</sup>See **Use Area** map for specific states.

## Early Postemergence Application

Soil Texture	<i>PAVILION 3.3 EC HERBICIDE</i>
Coarse	<b>DO NOT USE</b>
Medium, Fine	2.4 pts/A

### Restrictions

- Do not apply *PAVILION 3.3 EC HERBICIDE* preplant incorporated or preemergence.
- Do not apply *PAVILION 3.3 EC HERBICIDE* as a CULTI-SPRAY treatment in grain sorghum planted in double row beds.
- Do not replant grain sorghum if crop loss occurs.
- Do not apply in liquid fertilizer.
- Livestock can graze or be fed forage from *PAVILION 3.3 EC HERBICIDE*- treated grain sorghum fields after 21 days following application.
- Do not apply *PAVILION 3.3 EC HERBICIDE* in grain sorghum more than once per crop season.

### GREEN ONIONS

(Leeks, Spring Onions or Scallions, Japanese Bunching Onions,  
Green Shallots or Green Eschalots)

*PAVILION 3.3 EC HERBICIDE* may be applied preemergence, postemergence, or split application by ground, air, or chemigation.

### Use Directions

Uniformly apply 2.4 pints per acre of *PAVILION 3.3 EC HERBICIDE* as a broadcast spray to the soil surface as preemergence spray or as a postemergence spray to the crop at the 2 to 3 true- leaf stage at least 30 days before harvest. If *PAVILION 3.3 EC HERBICIDE* is to be applied sequentially as both a preemergence and postemergence spray, the preemergence spray must be applied 30 days prior to the postemergence spray.

### Chemigation Applications

*PAVILION 3.3 EC HERBICIDE* may be applied through sprinkler irrigation systems. Apply at 2 to 3 true-leaf stage at least 30 days before harvest. Follow all recommendations, special instructions and precautions in the general section covering Chemigation in **Spraying Instructions**.

### Restrictions

- Do not apply more than 2.4 pints per acre per application.
- Do not apply more than 4.8 pints per acre per season.
- Do not apply within 30 days of harvest.
- Do not feed forage or graze livestock in treated fields.
- Do not irrigate in excess of 0.5 inch of water.

### LENTILS AND PEAS

(English, Dry, Garden, Dwarf, Green, Pigeon, and Edible Pod)

*PAVILION 3.3 EC HERBICIDE* may be applied (fall) preplant surface or preplant incorporated for weed control in lentils and peas.

### Use Directions

**Preplant Incorporated** - *PAVILION 3.3 EC HERBICIDE* may be applied 60 days prior to planting up to immediately before planting. After application, rotary hoeing and shallow cultivation/ tillage can be practiced without reducing weed control. Avoid tillage that will bring untreated soil to the surface.

### Use Rates

Soil Texture	Broadcast Rate (pts/A)
Coarse	1.8
Medium	2.4
Fine	3.6

**Fall Applications** - Fall preplant surface and preplant incorporated applications may be made in North Dakota, South Dakota, Minnesota, Oregon, Washington, Montana, Idaho, and Wyoming only. Rainfall or irrigation is required for incorporation and activation. Unpredictable weed control can be expected since factors such as length of time between application and planting as well as uncontrollable weather factors will determine herbicide activity and longevity.

Apply *PAVILION 3.3 EC HERBICIDE* and incorporate (via rainfall, irrigation or mechanically) in late fall prior to planting lentils or peas (English, dry, garden, dwarf, green, pigeon, and edible pod) the following spring. Apply *PAVILION 3.3 EC HERBICIDE* in the late fall when soil temperatures are 45°F or below but before the ground freezes.

**Preplant Surface and Preplant Incorporated  
(Fall Application<sup>1</sup>) Use Rates**

Soil Texture	Broadcast Rate (pts/A)
Coarse	1.2 to 1.8
Medium	1.8 to 2.4
Fine	2.4 to 3.6

<sup>1</sup>For use in Idaho, Minnesota, Montana, North Dakota, Oregon, South Dakota, Washington, and Wyoming only.

**Restrictions**

- Do not use in California.
- Do not apply *PAVILION 3.3 EC HERBICIDE* preemergence in peas.
- Do not apply *PAVILION 3.3 EC HERBICIDE* more than once per cropping season.
- Do not apply to peas, lentils, pea or lentil forage, pea silage, pea hay, or pea straw grown for livestock feed.
- Do not apply in any type of irrigation system.
- Any crop registered for a preplant incorporated application of *PAVILION 3.3 EC HERBICIDE* can be double cropped after peas.
- For fall applications, do not apply when the air temperature is below 45°F.

**MINT  
(Peppermint and Spearmint)**

*PAVILION 3.3 EC HERBICIDE* may be applied by ground or air.

**Use Directions**

Make a single broadcast preemergence application of *PAVILION 3.3 EC HERBICIDE* to mint at 1.8 pints to 4.8 pints per acre, depending on soil texture (see chart below), to dormant established mint before weed emergence.

**Use Rates**

Soil Texture	Broadcast Rate (pts/A)
Coarse	1.8 to 2.4
Medium	2.4 to 4.8
Fine	2.4 to 4.8

**Precautions**

- After a *PAVILION 3.3 EC HERBICIDE* application, some temporary crop injury may be observed early in the growing season as mint breaks dormancy and begins to grow.
- *PAVILION 3.3 EC HERBICIDE* will not cause crop injury when applied according to the label under normal growing conditions.
- Non-uniform application may result in injury to crops, poor stands, or soil residues; conversely, uneven application may reduce weed control.
- Diseases, cold weather, excessive moisture, deep planting, low or high pH, salinity, or drought may weaken seedlings and plants and make them more susceptible to herbicidal damage.

**Restrictions**

- Do not apply *PAVILION 3.3 EC HERBICIDE* to "baby" mint in the first year of growth and establishment.
- Do not apply to mint that has broken dormancy or crop injury may result. Application to mint that is near dormancy break can result in crop injury. Risk of crop injury increases the closer application is to mint dormancy break.
- Do not apply to mint stands that have been weakened by age, disease, cold weather, excessive moisture, or other factors that reduce crop vigor. Mint growing under stress is more susceptible to herbicidal damage.
- Do not apply more than 4.8 pints per acre per season.
- Do not apply within 90 days of harvest.
- Do not allow livestock to graze on treated spent hay or feed treated spent hay to livestock.
- Do not apply this product on mint through any type of irrigation system.
- Do not use in California except as directed in supplemental labeling.

**NONBEARING FRUIT AND NUT TREE CROPS and NONBEARING VINEYARDS**

*PAVILION 3.3 EC HERBICIDE* may be applied for preplant incorporated, preplant surface, surface incorporated or preemergence weed control in several nonbearing fruit and nut tree crops and nonbearing vineyards. *PAVILION 3.3 EC HERBICIDE* may be used before or after transplanting the following nonbearing crops: Almond, Apple, Apricot, Cherry, Citrus, Grape, Grapefruit, Lemon, Nectarine, Orange, Peach, Pear, Pecan, Pistachio, Plum, Prune, Tangelo, Tangerine, Walnut, English. Apply the spray directly to the ground beneath the trees or vines.

*PAVILION 3.3 EC HERBICIDE* may be applied by ground, air, chemigation or flooded basin irrigation systems.

### Use Directions

*PAVILION 3.3 EC HERBICIDE* may be applied either in a single application or sequentially with an interval of 30 days or more. Apply *PAVILION 3.3 EC HERBICIDE* at 2.4 to 4.8 quarts per acre (depending on desired length of control, see chart below) per application.

**Preplant Surface** - Prior to transplanting, uniformly apply with ground or aerial equipment.

**Preplant Incorporated** - Uniformly apply *PAVILION 3.3 EC HERBICIDE* prior to transplanting but before weeds emerge. Incorporate *PAVILION 3.3 EC HERBICIDE* to a depth of 1 to 2 inches.

**Preemergence (postplant)** - Applications may be in a band or broadcast.

### Chemigation Applications

*PAVILION 3.3 EC HERBICIDE* may be applied through sprinkler irrigation systems. Follow all directions, special instructions and precautions in the section covering **Chemigation in Spraying Instructions**.

### Flooded Basin Irrigation Systems

*PAVILION 3.3 EC HERBICIDE* may be applied in flooded basin irrigation systems. Follow all directions, special instructions and precautions in the section covering **Flooded Basin Irrigation in Spraying Instructions**.

### Use Rate per Acre

Short-term control	2.4 quarts
Long-term control	4.8 quarts

### Precautions

- Avoid root contact with treated soil when placing transplants into the hole or injury may occur.
- Application and incorporation must be made prior to transplanting to avoid mechanical injury to the crop.

### Restrictions

- Do not feed forage or graze livestock in treated fields.
- Do not apply more than 4.8 quarts of *PAVILION 3.3 EC HERBICIDE* per acre per year in pome, stone and other fruit trees.
- Do not apply more than 7.3 quarts of *PAVILION 3.3 EC HERBICIDE* per acre per year in citrus, nut trees and grapevines.
- Do not apply over the top of trees or vines with leaves or buds. Contacting leaves, shoots, or buds with the spray mixture may cause malformed plant tissue.
- Do not apply to newly seeded nursery stock.
- Do not apply *PAVILION 3.3 EC HERBICIDE*-treated irrigation water over top of trees or vines with leaves or buds.

### Restrictions for newly transplanted and one-year-old grapevines:

- Apply only to dormant grapevines.
- Do not apply if buds have started to swell. Application after buds have started to swell may result in leaf distortion.
- Do not apply to newly transplanted trees or vines until ground has settled and no cracks are present.

### ONIONS (Direct-Seeded and Transplanted Dry Bulb) and SHALLOTS (Dry Bulb)

*PAVILION 3.3 EC HERBICIDE* may be applied by ground, air or chemigation.

### Use Directions

#### Chemigation Applications

*PAVILION 3.3 EC HERBICIDE* may be applied through sprinkler irrigation systems. Apply between the 2nd and 9th true-leaf stage (2nd to 6th true-leaf stage in California) unless otherwise specified below. Follow all directions, special instructions and precautions in the section covering **Chemigation in Spraying Instructions**.

#### Restriction for Chemigation Applications

- Do not irrigate in excess of 0.5 inch of water.

### Mineral Soils

#### Use Rates

Soil Texture	Broadcast Rate (pts/A)
Coarse	1.8
Medium	2.4
Fine	3.6

**State-Specific Use Directions  
In All States Except California:**

Apply *PAVILION 3.3 EC HERBICIDE* as a broadcast treatment when onions or shallots have 2 to 9 true leaves.

**Additional Use in Colorado, Kansas, and Nebraska:**

*PAVILION 3.3 EC HERBICIDE* may be applied sequentially in seeded onions. Apply first application of *PAVILION 3.3 EC HERBICIDE* at loop stage. Apply sequential application of *PPAVILION 3.3 EC HERBICIDE* early postemergence (2nd to 9th true-leaf stage).

**Restrictions for Additional Use in Colorado, Kansas, and Nebraska**

- Do not exceed the maximum labeled rate for a given soil texture.
- Do not apply *PPAVILION 3.3 EC HERBICIDE* at loop stage through the 9th true-leaf stage if heavy rains are expected, or severe crop injury may result.

**Additional Use in Colorado and the High Plains of Texas:**

For transplanted onions only, apply and shallow incorporate (less than 2 inches deep) *PAVILION 3.3 EC HERBICIDE* into preformed beds prior to transplanting.

**Additional Use in Idaho, Oregon, and Washington:**

Apply *PAVILION 3.3 EC HERBICIDE* as a broadcast treatment when onions or shallots are between the flag leaf to 9th true-leaf stage.

*PAVILION 3.3 EC HERBICIDE* may be used at 3.6 to 4.8 pints per acre for dodder control on medium- and fine-textured soils.

*PAVILION 3.3 EC HERBICIDE* may be applied in the fall or spring to the furrow area of land bedded in the fall in preparation for planting seed of dry bulb onions the following spring. Apply *PAVILION 3.3 EC HERBICIDE* as a banded application at rates based on appropriate soil texture. Band width should be approximately 1/2 the width of the row spacing. Keep *PAVILION 3.3 EC HERBICIDE* away from the area where onion seed will be planted. Harrow-off tops of beds following *PAVILION 3.3 EC HERBICIDE* furrow applications prior to planting onions. For selective weed control in the onion row, apply *PAVILION 3.3 EC HERBICIDE* as a banded postemergence application to flag leaf onions at the labeled rates based on soil texture. Apply *PAVILION 3.3 EC HERBICIDE* only once to the furrow area and once to the onion row as a postemergence application.

**Restriction for Additional Use in Idaho, Oregon, and Washington**

- Do not apply *PAVILION 3.3 EC HERBICIDE* using chemigation at the dodder control rate.

**Additional Use in Michigan:**

For mineral soils containing >10% organic matter, follow the directions for muck soils (see following).

**In California:**

*PAVILION 3.3 EC HERBICIDE* may only be applied as a single application when onions or shallots have 2 to 6 true leaves.

**Restrictions (Mineral Soils)**

- Do not mechanically incorporate except as specified for use on dry bulb onions in Colorado and the Texas High Plains.
- Do not exceed 3.6 pints per acre per crop (except Idaho, Oregon, and Washington).
- Do not apply within 60 days of harvest in California and within 45 days of harvest in all other states.
- Do not feed or graze these crops.
- Do not apply *PAVILION 3.3 EC HERBICIDE* preemergence through the loop stage if heavy rains are expected or severe crop injury may result. If irrigating immediately after *PAVILION 3.3 EC HERBICIDE* application at the preemergence through loop stage, do not irrigate in excess of 0.5 inch of water.

**Muck Soils**

**Use Rates**

*PAVILION 3.3 EC HERBICIDE* may be applied sequentially on muck soils as follows:

Application Timing and Growth Stage	Rate (pts/A)
Preemergence through Loop Stage	4.8
Early Postemergence (2nd to 6th true-leaf stage)	4.8
Late Postemergence (6th to 9th true-leaf stage)	4.8

**Restrictions (Muck Soils)**

- Do not apply to muck soils in California.
- Do not apply within 45 days of harvest.
- Do not feed or graze these crops.
- Do not apply more than 14.4 pints per acre per growing season on muck soils. To maximize crop safety, ensure good soil coverage during planting or transplanting and delay preemergence applications to the loop stage, if possible.
- Do not apply *PAVILION 3.3 EC HERBICIDE* preemergence through the loop stage if heavy rains are expected or severe crop injury may result. If irrigating immediately after *PAVILION 3.3 EC HERBICIDE* application at the preemergence through loop stage, do not irrigate in excess of 0.5 inch of water.
- Do not plant sugar beets, red beets, spinach, winter wheat, or winter barley as rotational crops on muck soils for 12 months from the time of last application if more than 3.6 pints per acre of *PAVILION 3.3 EC HERBICIDE* is applied to the onion crop.
- **If loss** of onion crop occurs, do not replant any crop other than onions in muck soil during the same cropping year and do not work the soil deeper than 2 inches.



## PEANUTS

*PAVILION 3.3 EC HERBICIDE* may be applied by ground, air, or chemigation.

*PAVILION 3.3 EC HERBICIDE* may be applied preplant incorporated in peanuts.

*PAVILION 3.3 EC HERBICIDE* may also be applied preemergence to peanuts grown under overhead irrigation.

### Use Directions

**Preplant Incorporated** - Apply *PAVILION 3.3 EC HERBICIDE* up to 60 days prior to planting and incorporate within 7 days after applications.

**Preemergence** - Apply *PAVILION 3.3 EC HERBICIDE* at planting or up to 2 days after planting and before crop emergence. To prevent decreased crop pegging, adequate incorporation must be achieved by applying a minimum of 0.75 inch of overhead irrigation or rainfall within 48 hours of application.

### Chemigation Applications

*PAVILION 3.3 EC HERBICIDE* may be applied through sprinkler irrigation systems. Follow all directions, special instructions and precautions in the section covering **Chemigation in Spraying Instructions**.

### Use Rates

Region	Rate (pts/A)
Texas, Oklahoma and New Mexico	1.2 to 2.4
Other peanut growing states*	2.4
*For heavy weed infestations, especially of Texas panicum, up to 3.6 pts/A of <i>PAVILION 3.3 EC HERBICIDE</i> can be used in Alabama, Georgia or Florida.	

### Restrictions

- Do not use in California.

## POTATOES

*PAVILION 3.3 EC HERBICIDE* may be applied by ground, air, or chemigation.

*PAVILION 3.3 EC HERBICIDE* may be applied preemergence, preemergence incorporated, or early postemergence in potatoes.

**Additional Weeds Controlled:** In addition to the weeds listed in **Table 1**, *PAVILION 3.3 EC HERBICIDE* will control stinging nettle in potatoes.

### Use Directions

**Preemergence** - Apply *PAVILION 3.3 EC HERBICIDE* after planting, but before potatoes and weeds emerge, or after dragoff.

**Preemergence Incorporated** - Apply *PAVILION 3.3 EC HERBICIDE* and incorporate after planting but before potatoes and weeds emerge. Where dragoff is practiced, apply *PAVILION 3.3 EC HERBICIDE* and incorporate before, at, or after dragoff, but before potatoes and weeds emerge. Incorporate *PAVILION 3.3 EC HERBICIDE* within 7 days of application. *PAVILION 3.3 EC HERBICIDE* must be thoroughly and uniformly incorporated into the top 1 to 2 inches of soil. Mechanical incorporation is not required if adequate rainfall for good crop and weed emergence occurs or irrigation is received within 7 days after application. Care must be taken so that incorporation equipment does not damage seed pieces or elongating sprouts.

**Early Postemergence** - Apply *PAVILION 3.3 EC HERBICIDE* from crop emergence to the 6-inch stage of growth.

### Chemigation Applications

*PAVILION 3.3 EC HERBICIDE* may be applied through sprinkler irrigation systems. Apply *PAVILION 3.3 EC HERBICIDE* preemergence after planting, after dragoff, or early postemergence through sprinkler irrigation systems. Follow all directions, special instructions and precautions in the section covering **Chemigation in Spraying Instructions**.

### Use Rates

Soil Texture	< 3% Organic Matter > 3% (pts/A)	
Coarse	1.8	1.8
Medium	2.4	3.6
Fine	3.6	3.6

### Restrictions

- Do not apply to sweet potatoes or yams.
- Do not apply preplant.
- Do not make more than one application of *PAVILION 3.3 EC HERBICIDE* per season.
- Application of *PAVILION 3.3 EC HERBICIDE* on White Rose variety potatoes during or followed by cool and/or wet weather conditions may result in crop injury.
- Do not apply *PAVILION 3.3 EC HERBICIDE* postemergence if potatoes are under stress from cold/wet or hot/dry conditions or crop injury may occur.

## RICE

**PAVILION 3.3 EC HERBICIDE** may be applied as a pre-flood, preemergence application in dry-seeded or drilled rice or as a delayed preemergence application in drilled dry-seeded rice or as an early postemergence application in dry-seeded rice. Treatments may be applied to conventional, reduced or minimum tillage, and no-till (stale seedbed) rice. The seedbed should be firm and free of clods and must be prepared to allow for good seed coverage. The use of a planter under conditions that do not allow good soil coverage of the rice seed can result in reduced stand or stunting if **PAVILION 3.3 EC HERBICIDE** contacts germinating rice seed.

**Additional Weeds Controlled:** In addition to the weeds listed in **Table 1**, **PAVILION 3.3 EC HERBICIDE** will control the following weeds in rice: junglerice and sprangletop.

### Use Directions

**Pre-Flood, Preemergence** - **PAVILION 3.3 EC HERBICIDE** may be applied for preemergence weed control as a pre-flood, pre-rice germination herbicide in lightly incorporated dry-seeded rice or on drilled rice.

**SEEDING DIRECTIONS:** For all rice seed incorporation methods, seed must be incorporated shallowly or no more than 1 inch below soil surface. Seed left on the surface may be injured or killed by **PAVILION 3.3 EC HERBICIDE**. However, it is recommended that 15 to 20% of seed total be visible at surface in order to ensure that seed is not covered too deeply. Seeding rates should be increased by a percentage corresponding to the amount of seed left on the surface. Adjust seeding ratios to meet individual practices, incorporation depths and field conditions.

**EXAMPLE:** Target seeding rate is 150 lbs per acre. If approximately 15% of seed is left on soil surface, seeding rate should then be increased 22.5 lbs per acre to 177.5 lbs per acre.

Seeding depths can be affected by soil textures, tillage practices, irrigation, and methods of mechanical incorporation. Seed that is incorporated either mechanically and/or by irrigation flush must remain at a shallow depth of no more than 1 inch below the soil surface. Fields where rice seed is incorporated too deeply will experience reduced crop stands.

Following are examples of typical implements that can be used for rice seed incorporation: rice roller/ridger, ring roller, light harrow, or flat roller. Regardless of the implement or method of incorporation used, seed incorporation must be less than 1 inch below the soil surface.

After rice seed is incorporated, uniformly apply to soil surface as broadcast spray the tank mixture of **PAVILION 3.3 EC HERBICIDE** at 2.4 pints per acre plus FirstChoice® SafeGuard™ spray adjuvant at the labeled rate. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Use of **PAVILION 3.3 EC HERBICIDE** without tank mixing with FirstChoice® SafeGuard™ spray adjuvant can result in crop injury and loss of rice stand.

After herbicide application, flush field with irrigation water with method best employed to facilitate a thorough soaking of field and a rapid drain. Tail water (runoff water) from flood irrigation that contains **PAVILION 3.3 EC HERBICIDE** should be re-circulated and contained in the field of initial application or used only on adjacent crops for which **PAVILION 3.3 EC HERBICIDE** (or other pendimethalin-based products) is registered for use.

Rice seed covered with water for longer than 8 days may result in reduced stand and weed control.

**Delayed Preemergence** - Apply **PAVILION 3.3 EC HERBICIDE** alone or with tank mix partner for delayed preemergence weed control in grain-drilled, dry-seeded rice. Apply **PAVILION 3.3 EC HERBICIDE** alone or in tank mixture to levees after the levees are pulled and planted. Exposed seeds that come in contact with **PAVILION 3.3 EC HERBICIDE** may be injured. Apply only when growing conditions favor vigorous rice growth. The seedbed should have adequate moisture for seed germination. Not for use in California.

Uniformly apply the specified rate of **PAVILION 3.3 EC HERBICIDE** after rice planting and before rice and weed emergence (spiking). Apply after the rice seed has absorbed water and germinated and after the soil has been previously sealed over the seed by at least 1 inch of rainfall or by irrigation (flush). If the soil has not been sealed by rain or flush, apply when 80 percent of germinated seeds have a primary root (radicle) or shoot at least 1/2-inch long. If there is insufficient moisture, flushing is recommended before **PAVILION 3.3 EC HERBICIDE** application to supply moisture for root (radicle) initiation and for vigorous rice and weed growth.

If applied to soil prior to these conditions or to cracked soil, stand reduction or stunting of rice may occur. Under some conditions, use of gibberellic acid-treated seed, heavy rainfall after application, or flushing after application may result in herbicide injury to rice. Rice can overcome moderate injury with appropriate cultural practices.

Due to the residual activity of **PAVILION 3.3 EC HERBICIDE**, this treatment may be applied if rice is too small to maintain a flood on the field for weed control. However, proper water management practices must be followed for normal rice growth and activity of **PAVILION 3.3 EC HERBICIDE**.

**Early Postemergence** - Apply **PAVILION 3.3 EC HERBICIDE** as a tank mix partner. Base applications on weed and crop size guidelines of the tank mix partner. Do not apply to fields with standing water. If necessary, fields may be flushed prior to treatment to produce vigorous rice and weed growth. Since soil and weeds must be completely exposed to spray coverage, no flood water should be on the field at the time of application. Cloddy soil, standing water (puddles) at the time of application, or cracks in the soil that form after application may result in reduced weed control. Because of residual activity of **PAVILION 3.3 EC HERBICIDE**, this treatment may be applied if rice is too small to maintain a flood on the field for weed control. However, proper water management practices must be followed for normal rice growth and activity of **PAVILION 3.3 EC HERBICIDE**.

Since the residual activity of **PAVILION 3.3 EC HERBICIDE** is activated by moisture, **PAVILION 3.3 EC HERBICIDE** is most effective in controlling emerging weeds when adequate rainfall or irrigation (flush) is received within 7 days after application.

**Use Rates**  
**Delayed Preemergence Applications**

Soil Texture	Rate (pts/A)
Sands, loamy sands	<b>DO NOT USE</b>
Sandy loams	1.8
Loams, silt loams, silts, sandy clay loams	2.4
Silty clay loams, clay loams, sandy clays, silty clays, clays	2.4

**Early Postemergence Application**

Soil Texture	Rate (pts/A)
Coarse	1.8
Medium	2.4
Fine	2.4

**Restrictions**

- Do not apply *PAVILION 3.3 EC HERBICIDE* as a pre-flood, preemergence treatment in rice unless tank mixed with FirstChoice® SafeGuard™ spray adjuvant.
- Do not apply *PAVILION 3.3 EC HERBICIDE* through any type of irrigation system.
- Do not apply in liquid fertilizer.
- Do not use on water-seeded rice except as specified in other INNICTIS labeling.
- Do not apply to rice fields if fields are used for fish production, especially catfish or crayfish farming.
- Do not use water containing *PAVILION 3.3 EC HERBICIDE* residues from rice cultivation to irrigate food or feed crops that are not registered for use with *PAVILION 3.3 EC HERBICIDE*.
- In case of a crop failure due to weather conditions or disease following treatment with *PAVILION 3.3 EC HERBICIDE* alone or in a tank mixture, only drilled dry-seeded rice may be immediately replanted; however, to the extent consistent with applicable law, the grower assumes all risks and consequences associated with replanting of rice because there is the potential for stand reduction or stunting. A 10 percent increase in seeding rate is recommended. Replant seed below the herbicide layer because reduced stand or stunting may occur if *PAVILION 3.3 EC HERBICIDE* contacts germinating rice seed. Do not replant with gibberellic acid-treated seed. Do not reapply *PAVILION 3.3 EC HERBICIDE* alone or in a tank mixture.
- Do not apply *PAVILION 3.3 EC HERBICIDE* and then flush for germination.
- Do not apply to stressed rice. Stress factors include cold or hot temperature extremes, excessive moisture or drought, problem soils, poor field drainage, or deep water after application.
- Do not apply early preemergence nor preplant incorporated as severe rice injury is possible.
- Do not feed forage or graze livestock in treated fields.

**SOYBEANS**

*PAVILION 3.3 EC HERBICIDE* may be applied in conventional, minimum, or no-till as a fall surface, fall incorporated, preplant surface, preplant incorporated, or preemergence application in soybeans.

**Additional Weeds Controlled:** In addition to the weeds listed in **Table 1**, *PAVILION 3.3 EC HERBICIDE* will control or reduce competition from the following weeds in soybeans: itchgrass and red rice. For specific rates for red rice and itchgrass management, see table at end of this section.

**Use Directions**

**Fall Applied** - *PAVILION 3.3 EC HERBICIDE* may be surface applied or incorporated in the fall, after fall harvest and prior to ground freeze in states north of I-80 and the entire states of Iowa, Illinois, Indiana, Kansas, Kentucky, Missouri, Nebraska, Ohio, Oklahoma, and Texas. Fall applications of *PAVILION 3.3 EC HERBICIDE* will not provide season-long weed control.

**Preplant Surface** - Apply *PAVILION 3.3 EC HERBICIDE* up to 15 days prior to planting. *PAVILION 3.3 EC HERBICIDE* may be applied up to 45 days prior to planting when used in a tank mix or applied sequentially with glyphosate + imazethapyr, imazamox or imazethapyr herbicides. *PAVILION 3.3 EC HERBICIDE* tank mixes and sequential programs as specified under the tank mix section. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

**Preplant Incorporated** - Apply *PAVILION 3.3 EC HERBICIDE* up to 60 days prior to planting and incorporate within 7 days after application.

**Preemergence** - Apply *PAVILION 3.3 EC HERBICIDE* at planting or up to 2 days after planting. Apply to a firm seedbed free of clods.

**Preemergence Restriction**

- Do not make applications of *PAVILION 3.3 EC HERBICIDE* preemergence north of Interstate 80, except in the states of Indiana, Michigan and Ohio, or as specified in INNICTIS supplemental labeling.

**Use Rates****Fall Surface, Fall Incorporated, Preplant Surface, or Preplant Incorporated**

Soil Texture	< 3% Organic Matter > 3% (pts/A)	
Coarse	1.8	2.4
Medium	3.0 <sup>1</sup>	3.6
Fine <sup>2</sup>	3.6	3.6

<sup>1</sup>Do not exceed 2.1 pts for southern states; see **Use Area** map for specific states.

<sup>2</sup>For heavy clay soils, apply *PAVILION 3.3 EC HERBICIDE* at the broadcast rate of 3.6 pints per acre.

**Preemergence Applications**

Soil Texture	< 3% Organic Matter > 3% (pts/A)	
Coarse	1.8	1.8
Medium	2.4	2.4
Fine	2.4	3.0

**Preplant Incorporated Applications for Red Rice Control and Itchgrass Suppression**

Soil Texture	Up to 3% Organic Matter <sup>1</sup> (pts/A)
Coarse	3.6
Medium	3.6
Fine	4.8

<sup>1</sup>This use is not specified for soils with more than 3% organic matter.

**Restrictions**

- Do not **APPLY POSTEMERGENCE** or serious crop injury can result.
- Do not use *PAVILION 3.3 EC HERBICIDE* in soybeans in California.
- Livestock can graze or be fed forage from treated soybean fields.
- Do not apply within 85 days of harvest.
- Do not exceed one application per crop season at the highest rate per acre for any given soil type and application method.

**STRAWBERRY**

*PAVILION 3.3 EC HERBICIDE* may be applied by ground, air, or chemigation.

**Use Directions**

**Stunting, reduced growth, or reduction in daughter plants may occur with this use.** Uniformly apply 1.8 to 3.6 pints per acre of *PAVILION 3.3 EC HERBICIDE* as a broadcast spray to the soil surface at pre-transplant time. A second application of 1.8 to 3.6 pints per acre of *PAVILION 3.3 EC HERBICIDE* may be applied in a band to the soil between crop rows 35 days before harvest, but do not concentrate the rate per acre into the treated area and do not allow spray to contact strawberry plants. The second application rate is based on per unit of treated area.

**Chemigation Applications**

*PAVILION 3.3 EC HERBICIDE* may be applied through sprinkler irrigation systems. Follow all recommendations, special instructions and precautions in the general section covering

**Chemigation in Spraying Instructions.****Use Rates**

Soil Texture	Broadcast Rate (pts/A)
Coarse	1.8
Medium	2.4 to 3.0
Fine	3.0 to 3.6

**Restrictions**

- Do not apply more than 3.6 pints per acre per application.
- Do not apply more than 7.2 pints per acre per season.
- Do not apply within 35 days of harvest.
- Do not feed forage or graze livestock in treated fields.
- Do not apply if row is to later be covered with plastic.
- Do not allow *PAVILION 3.3 EC HERBICIDE*-treated irrigation water to contact strawberry plants.

## SUGARCANE

### Use Directions

*PAVILION 3.3 EC HERBICIDE* may be applied preemergence through layby to plant or ratoon sugarcane. Applications may be made band or broadcast. Although there may be adequate crop tolerance for postemergence applications at layby, the spray must be directed under the sugarcane canopy in order to obtain effective weed control.

*PAVILION 3.3 EC HERBICIDE* must be thoroughly and uniformly incorporated into the soil with either (a) mechanical incorporation equipment as outlined below, or (b) with rainfall or irrigation, if rainfall or irrigation is adequate for good crop and weed emergence and received within 7 days after application. If rainfall or irrigation is not obtained, *PAVILION 3.3 EC HERBICIDE* should be mechanically incorporated.

### Mechanical Incorporation

*PAVILION 3.3 EC HERBICIDE* should be applied to loosened beds and incorporated into the top 1 to 2 inches of soil within 7 days after application.

### Use Rates

Use Area	Broadcast Rate <sup>1</sup> (pts/A)
All states, except Hawaii	4.8 to 7.2
Muck soils (Florida only)	4.6 to 9.7
Hawaii	4.8 to 9.7

<sup>1</sup>Use the high rate if: clay soils; no mechanical incorporation is planned; heavy weed populations are anticipated; for grass infestation is anticipated; shaving is planned.

### Restrictions

- Do not exceed 14.4 pints of *PAVILION 3.3 EC HERBICIDE* per acre in one growing season.
- Do not use less than 11 gallons of water as a carrier when applying *PAVILION 3.3 EC HERBICIDE* for weed control.
- Ratoon sugarcane must be lightly shaved in early spring to remove the old stubble before incorporation over the line of sugarcane is possible. Carefully adjust equipment to incorporate without causing excessive damage to emerging shoots.
- Do not make aerial applications at close-in because complete and uniform coverage cannot be obtained.
- Do not apply through any type of irrigation system.
- Do not apply within 90 days of harvest.
- Do not graze treated fields or feed treated forage or fodder to livestock.

## SUNFLOWERS

*PAVILION 3.3 EC HERBICIDE* may be applied preplant incorporated in all states. Fall preplant incorporated applications may be made in North Dakota, South Dakota and Minnesota only. *PAVILION 3.3 EC HERBICIDE* may be applied preemergence in conventional tillage sunflowers, except in the state of California.

**Plant sunflowers 1.5 inches to 2 inches deep and completely cover with soil.**

### Use Directions

**Preplant Incorporated (Spring)** - Apply up to 60 days prior to planting and incorporate within 7 days after application.

**Preplant Incorporated (Fall applications in North Dakota, South Dakota and Minnesota)** – Apply *PAVILION 3.3 EC HERBICIDE* and immediately incorporate in late fall prior to planting sunflowers the following spring. Apply *PAVILION 3.3 EC HERBICIDE* in the late fall when soil temperatures are 45°F or below but before the ground freezes.

Prior to sunflower planting in the spring, fields treated with *PAVILION 3.3 EC HERBICIDE* should receive at least one shallow additional incorporation. Spring incorporation should be at an angle to the last tillage operation.

**Preemergence** - Apply *PAVILION 3.3 EC HERBICIDE* at planting or up to 2 days after planting. If dry conditions with limited precipitation exist or unseasonably cool temperatures following planting are forecast, apply *PAVILION 3.3 EC HERBICIDE* prior to planting and mechanically incorporate with tillage.

### Preemergence Precautions

- Preemergence applications of *PAVILION 3.3 EC HERBICIDE* to sunflowers may increase the likelihood of crop injury, especially when sunflowers are grown in stress situations, such as compacted soils.
- Decreased herbicide performance compared to preplant incorporated applications may also result from a preemergence application.

**Use Rates****Preplant Incorporated (Spring) or Preemergence (Conventional Tillage)**

Soil Texture	Southern States <sup>1</sup> (pts/A)	Northern States <sup>1</sup> (pts/A)	
		<3% Organic Matter	>3%
Coarse	1.8	2.4	2.4
Medium	2.4	3.0	3.6
Fine	3.6	3.6	3.6

<sup>1</sup>See **Use Area** map for specific states.

**Preplant Incorporated (Fall) Application<sup>1</sup>**

Soil Texture	< 3% Organic Matter > 3% (pts/A)	
Coarse	3.0	3.0
Medium	3.0	4.2
Fine	4.2	4.2

<sup>1</sup>For use in North Dakota, South Dakota and Minnesota Only.

**Restrictions**

- For preplant incorporated, do not apply when the air temperature is below 45°F.

**NO-TILL SUNFLOWERS**

*PAVILION 3.3 EC HERBICIDE* may be applied at 3.6 pints per acre up to 30 days before planting (preplant) to immediately after planting (preemergence).

*PAVILION 3.3 EC HERBICIDE* is most effective in controlling weeds when adequate rainfall or overhead irrigation is received within 7 days after application.

**Restrictions (all tillage types)**

- Do not apply *PAVILION 3.3 EC HERBICIDE* postemergence.
- Do not feed forage or graze livestock in treated sunflower fields.
- Do not use in California.

**TOBACCO**

*PAVILION 3.3 EC HERBICIDE* may be applied preplant incorporated or as a layby application in transplanted tobacco.

**Use Directions**

**Preplant Incorporated** - Apply *PAVILION 3.3 EC HERBICIDE* with ground sprayer up to 60 days prior to transplanting tobacco and incorporate within 7 days after application.

**Preplant Incorporated Precautions**

- Applied according to directions and under normal growing conditions, *PAVILION 3.3 EC HERBICIDE* will not harm transplanted tobacco.
- Under stress conditions for plant growth such as cold/wet or hot/dry weather, *PAVILION 3.3 EC HERBICIDE* can produce a temporary retardation of tobacco development.

**Layby** - *PAVILION 3.3 EC HERBICIDE* may be applied as a directed spray following the last normal cultivation (layby), usually 4 to 6 weeks after transplanting tobacco. Apply *PAVILION 3.3 EC HERBICIDE* in a 16- to 24-inch band between the crop rows.

**Layby Precaution**

- The spray should not contact tobacco plants.

**Use Rates**  
**Preplant Incorporated Application**

Use Area	Soil Texture	Rate (pts/A)
Florida Georgia Maryland North Carolina South Carolina Virginia	Coarse	2.4
	Medium sandy clay loams, loams	2.4
	silt loams, silts	3.0
	Fine	3.0
Other states	Coarse	2.4
	Medium	3.6
	Fine	3.6

**Layby Application**

Soil Texture	Broadcast Rate (pts/A)
Coarse	1.8
Medium	2.4
Fine	2.4

**Restrictions**

- Do not apply as a broadcast spray as contact may cause malformed tobacco leaves.

**WHEAT**

*PAVILION 3.3 EC HERBICIDE* may be applied by ground or air.

*PAVILION 3.3 EC HERBICIDE* may be applied preemergence, delayed preemergence, or postemergence to wheat for weed control in fall, winter or spring seeded wheat.

**Use Directions**

Apply to a seedbed that is firm and free of clods and trash. The seedbed **MUST** be prepared to ensure good seed coverage by the soil and seed-to-soil contact. Use high quality seed. When applications of *PAVILION 3.3 EC HERBICIDE* are intended to be made preemergence or delayed preemergence, plant seed at least 1 inch deep to avoid possible crop injury, but not too deep for proper germination. When applications of *PAVILION 3.3 EC HERBICIDE* are intended to be made postemergence, plant seed at least 0.5 inch to 1.0 inch to avoid crop injury.

Uniformly apply *PAVILION 3.3 EC HERBICIDE* as a preemergence, or delayed preemergence (after wheat seed has germinated), or postemergence treatment from the 1st-leaf stage of wheat until before the flag leaf is visible/emerged for weed control. *PAVILION 3.3 EC HERBICIDE* should be applied prior to weed emergence. Adequate rainfall or irrigation within 7 days after application will provide the most consistent weed control.

For control of established weeds, *PAVILION 3.3 EC HERBICIDE* may be tank mixed with any postemergence herbicide registered for use in wheat. *PAVILION 3.3 EC HERBICIDE* will provide residual control of the weeds listed in this label. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Always perform a mixing test to check the compatibility of *PAVILION 3.3 EC HERBICIDE* with all potential tank mix partners.

**Use Rates**

Soil Texture	Southern States <sup>1</sup> (pts/A)	Northern States <sup>1</sup> (pts/A)
Coarse	1.8 to 2.4	1.8
Medium	1.8 to 3.6	3.0
Fine	2.4 to 3.6	2.4 to 3.6

<sup>1</sup>See Use Area map for specific states.

## Precautions

- Emerged weeds will not be controlled by this treatment.

## Restrictions

- Do not apply more than 3.6 pints per season.
- **Note:** If loss of grain crop occurs, any crop registered for *PAVILION 3.3 EC HERBICIDE* preplant incorporated use may be replanted the same year without adverse effects.
- Do not replant wheat.
- Do not apply *PAVILION 3.3 EC HERBICIDE* within 60 days of harvest of wheat grain or straw.
- Do not apply *PAVILION 3.3 EC HERBICIDE* within 28 days of harvest of wheat hay.
- Do not apply *PAVILION 3.3 EC HERBICIDE* within 11 days of harvest of wheat forage

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

**PESTICIDE STORAGE: DO NOT STORE BELOW 40° F.** - Extended storage at temperatures below 40° F can result in the formation of crystals on the bottom of the container. If crystallization does occur, store the container on its side at room temperature (70° F) and rock occasionally until crystals redissolve.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of federal law. If these wastes cannot be disposed of by use according to label directions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

### CONTAINER HANDLING

**Nonrefillable Container. Do not reuse or refill this container.** Triple rinse or pressure rinse container (or equivalent) promptly after emptying; then offer for recycling, if available, or reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

**Triple rinse containers small enough to shake (capacity ≤ 5 gallons) as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

**Triple rinse containers too large to shake (capacity > 5 gallons) as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Repeat this procedure two more times.

**Pressure rinse as follows:** Empty the remaining contents into application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank, or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

**Refillable Container.** Refill this container with pesticide only. Do not reuse this container for any other purpose. Triple rinsing the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller.

**Triple rinse as follows:** To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

When this container is empty, replace the cap and seal all openings that have been opened during use; return the container to the point of purchase or to a designated location. This container must only be refilled with a pesticide product. Do not reuse the container for any other purpose. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, worn-out threads and closure devices. Check for leaks after refilling and before transport. Do not transport if this container is damaged or leaking. If the container is damaged, or leaking, or obsolete and not returned to the point of purchase or to a designated location, triple rinse emptied container and offer for recycling, if available, or dispose of container in compliance with state and local regulations.

## CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of INVICTIS CROP CARE, LLC or Seller. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold INVICTIS CROP CARE, LLC and Seller harmless for any claims relating to such factors.

INVICTIS CROP CARE, LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or INVICTIS CROP CARE, LLC, and TO THE EXTENT CONSISTENT WITH APPLICABLE LAW Buyer and User assume the risk of any such use. INVICTIS CROP CARE, LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law, neither INVICTIS CROP CARE, LLC nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF INVICTIS CROP CARE, LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF INVICTIS CROP CARE, LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.

INVICTIS CROP CARE, LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of Sale and Limitation of Warranty and Liability which may not be modified except by written agreement signed by a duly authorized representative of INVICTIS CROP CARE, LLC.

All trademarks are the property of their respective owners.