

SPECIMEN



PONCHO[®] **600**

Net
Contents:

2.5 Gallons



Bayer SeedGrowth™

For: seed treatment use on canola, cereal grains (except corn, rice, sorghum and wild rice), rapeseed, corn, rice, sorghum, and soybean for the control of listed insect pests.

All seed treated with this product must be conspicuously coloured at the time of treatment.

ACTIVE INGREDIENT:

Clothianidin: **48.00%**

OTHER INGREDIENTS: **52.00%**

TOTAL: 100.00%

Contains 5.00 pounds clothianidin per U.S. gallon

EPA Reg. No. 264-789

KEEP OUT OF REACH OF CHILDREN CAUTION

For **MEDICAL** And **TRANSPORTATION** Emergencies **ONLY**

Call 24 Hours A Day 1-800-334-7577

For **PRODUCT USE** Information Call 1-866-99BAYER (1-866-992-2937)

Please refer to booklet for additional precautionary statements and directions for use.

Produced for:
Bayer CropScience LP
P.O. Box 12014, 2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
PONCHO is a registered trademark of Bayer.
©2018 Bayer CropScience

171108C 05/18

US61379957C

FIRST AID

If Swallowed:	<ul style="list-style-type: none">• Immediately call a poison control center or doctor for treatment advice.• Do not induce vomiting unless told to by a poison control center or doctor.• Have person sip a glass of water if able to swallow.• Do not give anything by mouth to an unconscious person.
If Inhaled:	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or ambulance, and then give artificial respiration, preferably mouth-to-mouth if possible.• Call a poison control center or doctor for further treatment advice.
If on Skin or Clothing:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
<p>In case of emergency call toll free the Bayer CropScience Emergency Response Telephone No. 1-800-334-7577.</p> <p>Have a product container or label with you when calling a poison control center or doctor, or going for treatment.</p>	
Note to Physician: No specific antidote is available. Treat the patient symptomatically.	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

- Harmful if swallowed. Harmful if absorbed through the skin.
- Harmful if inhaled.
- Avoid contact with skin, eyes, or clothing. Avoid breathing spray mist.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt
- Long pants
- Chemical-resistant gloves made out of: barrier laminate, nitrile rubber ≥ 14 mils, butyl rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, polyvinyl chloride rubber ≥ 14 mils or Viton ≥ 14 mils
- Shoes plus socks.

USER SAFETY RECOMMENDATIONS

- Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

USER SAFETY REQUIREMENTS

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENVIRONMENTAL HAZARDS

This product is toxic to birds and aquatic invertebrates. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high-water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwater or rinsate. Do not apply where runoff is likely to occur. Drift and runoff from treated areas may be hazardous to aquatic organisms in water adjacent to treated areas. Apply this product only as specified on the label.

This chemical has properties and characteristics associated with chemicals detected in ground water. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow. Therefore contain any product spills or equipment leaks and dispose of wastes according to the disposal instructions on this label.

Treated seed exposed on soil surface maybe hazardous to birds and mammals. Cover or collect spilled seeds.

This compound is highly toxic to bees exposed directly (contact).

CONDITIONS OF SALE AND LIMITATIONS OF WARRANTY AND LIABILITY

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

Treatment of highly mechanically damaged seed, or seed of known low vigor and poor quality, may result in reduced germination and/or reduction of seed and seedling vigor. Treat and conduct germination tests on a small portion of seed before committing the total seed lot to a selected chemical treatment. Due to seed quality conditions beyond the control of Bayer CropScience, no claims are made to guarantee germination of carry-over seed.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience. To the extent consistent with applicable law, all such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE'S ELECTION, THE REPLACEMENT OF PRODUCT.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read the entire label before using this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas (that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water), is:

- Long-sleeved shirt
- Long pants
- Chemical-resistant gloves made out of: barrier laminate, nitrile rubber ≥ 14 mils, butyl rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, polyvinyl chloride rubber ≥ 14 mils or Viton ≥ 14 mils
- Shoes plus socks.

PRODUCT INFORMATION

Poncho[®] 600:

- is a systemic insecticide seed treatment for use on canola, cereal grains (except corn, rice, sorghum and wild rice), corn, rapeseed, rice, sorghum, and soybean for the control of listed insects.
- provides early season protection against the listed insect pests.
- provides protection under low to moderate corn rootworm pressure.

APPLICATION INSTRUCTIONS

Poncho 600:

- is to be used in liquid or slurry treaters. Mix thoroughly before use or use entire container at one time. Pre-test all tank mixes to determine physical compatibility between formulations. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions and precautionary language of the products in the mixture.
- Treatment of highly mechanically damaged seed, or seed of known low vigor or poor quality, may result in reduced germination and/or reduction of seed and seedling vigor. Treat and conduct germination tests on a small portion of seed before committing the total seed lot to a selected chemical treatment. Due to seed quality conditions beyond the control of Bayer CropScience, no claims are made to guarantee germination of carry-over seed.

USE RESTRICTIONS

- All seed treated with this product must be conspicuously coloured at the time of treatment. This product contains no colorant. An EPA approved dye or colorant must be added when this product is applied to seed to distinguish and prevent subsequent inadvertent use as a food for man or feed for animals. Regulations pertaining to coloration of treated seed enforced by 40 CFR 153.155 must be strictly adhered to when using this product.
- For use only in commercial seed treatment facilities. Not for use in hopper box, planter box, slurry box, or other on-farm seed treatment applications except for cereal grains (except corn, rice, sorghum and wild rice).
- Regardless of method of application, (seed treatment, soil, or foliar) DO NOT apply more than 0.2 lbs of active ingredient clothianidin per acre per year.
- Not for use on sorghum seed in the state of California.

Refer to the specific use directions and restrictions in each crop table below.

SPECIFIC CROP DIRECTIONS

COMMERCIAL SEED TREATMENT FACILITIES

CANOLA; RAPESEED			
Insect Pests	Product Rate (fl oz per 100 lbs seed)	Product Rate (mg ai/seed)¹	Remarks
Flea beetle Wireworm	3.84	0.00664	Use this rate under low to moderate flea beetle pressure.
	5.12	0.008863	Use this rate under moderate flea beetle pressure.
	10.23	0.01770	Use this rate under high to extreme flea beetle pressure where extended control is required.
• ¹ Rate is calculated based on number of seeds/lb averages of canola (102500).			

CEREAL GRAINS (Except Corn, Rice, Sorghum and Wild Rice) Barley, Buckwheat, Millet (pearl and proso), Oats, Rye, Teosinte, Triticale, Wheat

Insect Pests	Product Rate (fl oz per 100 lbs seed)	Product Rate (mg ai/seed)¹
Wireworm	0.25 - 1.79	0.00324 - 0.02318
Aphid (including Bird cherry-oat, English grain, Greenbug, and Russian wheat aphid)	0.75 - 1.79	0.00971 - 0.02318
Grasshopper ² Hessian Fly	1.79	0.02318

Crop Specific Directions:

- For wireworm seed protection, use a 0.25 fl oz/100 lbs seed rate or higher, depending on field history and site populations.
- For Hessian fly protection to wheat seedlings in the fall season, use 1.79 fl oz/100 lbs seed.
- For maximum effectiveness, treat seed uniformly. Use the higher rate to provide increased length of protection and from heavy insect pressure and to reduce potential spread of Barley yellow dwarf virus due to aphid vectors.
- Grasshopper feeding damage to seedling plants under low to moderate insect pressure will be reduced when Poncho 600 is used. Treated seed may be planted as a 50 - 60 foot border around the edges of the crop field as a barrier to grasshoppers advancing into the interior of the field.
- ¹ Rate is calculated based on number of seeds/lb averages of wheat (13700).
- ² Not registered for use in California.

CORN (Field and Pop)		
Insect Pests	Product Rate (mg ai per seed)	Product Rate (fl oz per 80,000 Seeds)
Corn rootworm (including Northern, Western, Southern and Mexican) Southern corn billbug	1.25	5.64
Chinch bug Corn flea beetle Corn leaf aphid Cutworm, black Grape colaspis Seedcorn maggot Southern corn leaf beetle Southern corn rootworm Southern green stinkbug Storage pests ¹ (including Indian meal moth, Lesser grain borer, Red flour beetle, Rice weevil, Rusty grain beetle) Sugarcane beetle White grub (including European, Chafer larvae, May/June beetle larvae, Japanese beetle larvae) Thrips Wireworm	0.25 - 0.50	1.13 - 2.26
Crop Specific Directions:		
<ul style="list-style-type: none"> • ¹ It is recommended that seed with existing populations of storage pests be fumigated prior to treating and bagging seed. 		

CORN (Sweet)		
Insect Pests	Product Rate (mg ai per seed)	Product Rate (fl oz per 80,000 Seeds)
Chinch bug Corn flea beetle Corn leaf aphid Cutworm, black Grape colaspis Seedcorn maggot Southern corn leaf beetle Southern corn rootworm Southern green stinkbug Storage pests ¹ (including Indian meal moth, Lesser grain borer, Red flour beetle, Rice weevil, Rusty grain beetle) Sugarcane beetle White grub (including European, Chafer larvae, May/June beetle larvae, Japanese beetle larvae) Thrips Wireworm	0.25 - 0.50	1.13 - 2.26
Crop Specific Directions:		
<ul style="list-style-type: none"> • ¹ It is recommended that seed with existing populations of storage pests be fumigated prior to treating and bagging seed. 		

RICE		
Insect Pests	Product Rate (fl oz per 100 lbs seed)	Remarks
Aphid Black Bug Chinch Bug Grape Colaspis Rice Water Weevil Thrips	1.93	Provides early season protection against listed pests.
Crop Specific Restrictions: <ul style="list-style-type: none"> • Do not apply to Wild Rice. • Not registered for use in California. • Poncho 600 is to be used only as a dry-seeded rice seed treatment. • Do not use Poncho 600 in areas where rice/crawfish aquaculture practices are in place or in or near fish farms, shrimp, prawn or crab pond or nursery operations. • Poncho 600 is not to be followed by a spray treatment of another neonicotinoid insecticide. Follow approved guidelines established for insect resistance management by consulting your local extension pest management expert. 		

SORGHUM		
Insect Pests	Product Rate (fl oz per 100 lbs seed)	Product Rate (mg ai/seed)¹
Black granulate cutworm Aphid (including Greenbug, Yellow sugarcane aphid, Corn leaf aphid and English grain aphid) Chinch bugs White grubs Wireworms	5.1 - 6.4	0.05 - 0.06
Crop Specific Restrictions: <ul style="list-style-type: none"> • Not registered for use in California. • ¹Rate is calculated based on number of seeds/lb averages of sorghum (18000). 		

SOYBEAN		
Insect Pests	Product Rate (fl oz per 140,000 seeds)	Product Rate (mg ai/seed)
Aphids Bean leaf Beetle Grape Colaspis Leafhoppers Seed Corn Maggot Three Cornered Alfalfa Hopper Thrips White Grubs Wireworm	0.87	0.11
<ul style="list-style-type: none"> • Not registered for use in California. 		

ON FARM SEED TREATMENT

CEREAL GRAINS (Except Corn, Rice, Sorghum and Wild Rice) Barley, Buckwheat, Millet (pearl and proso), Oats, Rye, Teosinte, Triticale, Wheat

Insect Pests	Product Rate (fl oz per 100 lbs seed)	Product Rate (mg ai/seed) ¹
Wireworm	0.25 - 1.79	0.00324 - 0.02318
Aphid (including Bird cherry-oat, English grain, Greenbug, and Russian wheat aphid)	0.75 - 1.79	0.00971 - 0.02318
Hessian Fly	1.79	0.02318

Crop Specific Directions:

- For wireworm seed protection, use a 0.25 fl oz/100 lbs seed rate or higher, depending on field history and site populations.
- For Hessian Fly protection to wheat seedlings in the Fall season, use 1.79 fl oz/100 lbs seed.
- For maximum effectiveness, treat seed uniformly. Use the higher rate to provide increased length of protection and from heavy insect pressure and to reduce potential spread of Barley yellow dwarf virus due to aphid vectors.
- Grasshopper feeding damage to seedling plants under low to moderate insect pressure will be reduced when Poncho 600 is used. Treated seed may be planted as a 50 to 60 foot border around the edges of the crop field as a barrier to grasshoppers advancing into the interior of the field.

Crop Specific Restrictions:

- Regardless of application method (seed treatment, soil, or foliar), do not apply more than 0.2 lb active ingredient clothianidin/A/yr.
- ¹ Rate is calculated based on number of seeds/lb averages of wheat (13700).

ROTATIONAL CROP PLANT-BACK INTERVALS

Crop Planted	Immediate Plant-back	30-Day Plant-back	4-Month Plant-back	8-Month Plant-back	12-Month Plant-back
Field corn, popcorn, soybean, cereal grains, broccoli, carrot, leeks, onions	Cereal Grains, Canola, Soybeans, Cotton, Root And Tuber Vegetables, Leafy Green Vegetables, Brassica (Cole) Vegetables, Fruiting Vegetables, Cucurbit Vegetables, and Bulb Vegetables	Grasses, non-grass animal feeds, dry beans		Sugar cane	Any crops without an earlier plant-back interval
Sweet corn, canola, rapeseed, sorghum	Cereal Grains, Canola, Soybeans, Cotton, Root And Tuber Vegetables, Leafy Green Vegetables, Brassica (Cole) Vegetables, Fruiting Vegetables, Cucurbit Vegetables, and Bulb Vegetables	Grasses, non-grass animal feeds, dry beans	Any crop without an earlier plant-back interval		

SEED TAG LABELING

The Federal Seed Act requires that the container of seed treated with **PONCHO 600** must be labeled with the following statements:

- This seed has been treated with:
 - **PONCHO 600**, which contains **48.0% clothianidin**.
- Do not use treated seed for food, feed, or oil production.

(continued)

In addition, the US Environmental Protection Agency requires the following statements on the container of seed treated with **PONCHO 600**:

- **Store treated seeds away from food and feedstuffs.**
- **Do not allow children, pets, or livestock to have access to treated seed.**
- **Wear long-sleeved shirt, long pants and chemical-resistant gloves when handling treated seed.**
- Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by-products that are used in agronomic practice.
- Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading.
- Dispose of all excess treated seed. Leftover treated seed may be double sown around the headland or buried away from water sources in accordance with local requirements.
- Do not contaminate water bodies when disposing of planting equipment wash waters.
- Dispose of seed packaging or containers in accordance with local requirements. Do not use empty seed bags for any other purpose.
- This compound is highly toxic to bees exposed directly (contact). Ensure that planting equipment is functioning properly in accordance with manufacturing recommendations to minimize seed coat abrasion during planting to reduce dust, which can drift to blooming crops or weeds.
- Rapeseed greens and seed grown or harvested from Poncho® 600-treated seed must not be used for feed or human consumption. Rapeseed grown and harvested from Poncho® 600-treated seed is only for industrial uses and cannot be used for edible oil or any other human/feed consumption.
- This compound is toxic to birds and mammals. Treated seeds exposed on soil surface may be hazardous to birds and mammals. Cover or collect treated seeds spilled during loading.
- In soybean, foliar sprays of clothianidin can be applied 45 days after planting Poncho 600 treated seeds for mid-to-late season foliar insects.
- For all other crops do not apply a neonicotinoid insecticide foliar spray after planting treated seed. Follow approved guidelines established for insect resistance management by consulting your local extension pest management expert.
- **For Field Corn, Popcorn:** Immediate Plant-back: Cereal grains, Field corn, popcorn, sweet corn, rapeseed, canola, soybeans, root and tuber vegetable, 30-Day Plant-back: Grasses, non-grass animal feeds, dry beans, 8-Month Plant-back: Sugar cane, 12-Month Plant-back: Any crops without an earlier plant-back interval.
- **For Sweet corn, Canola, Rapeseed, and Sorghum:** Immediate Plant-back: Cereal grains, Field corn, popcorn, sweet corn, rapeseed, soybeans, canola, sorghum, root and tuber vegetables, 30-Day Plant-back: Grasses, non-grass animal feeds, dried beans, 4-Month Plant-back: Any crop without an earlier plant-back interval.
- **For Soybeans:** Immediate Plant-back: Cereal grains, Field corn, popcorn, sweet corn, cotton, rapeseed, canola, soybeans, root and tuber vegetable, 30-Day Plant-back: Grasses, non-grass animal feeds, dry beans, 8-Month Plant-back: Sugar cane, 12-Month Plant-back: Any crops without an earlier plant-back interval.

- **For wheat, barley, oats, rye, and triticale:** Treated areas may be replanted with any crop specified on clothianidin label, or any crop for which a tolerance exists for clothianidin, as soon as practical following the last application. Areas planted with treated seed may be replanted immediately with cotton, corn, rapeseed, sorghum; sugar beet, potato, cereal grains (Crop Group 15 except rice), canola, and the following vegetable crops: root vegetables (Crop Group 1), bulb vegetables (Crop Group 3), leafy green vegetables (Crop Group 4A), Brassica (cole) leafy vegetables (Crop Group 5), fruiting vegetables (Crop Group 8), cucurbit vegetables (Crop Group 9). These areas may also be replanted after 30 days with soybeans, dried beans and dried peas. Do not plant any other crop in the treated area for at least one year after treated seeds are planted.
- Follow-up foliar sprays of registered insecticide may be needed under heavy pest pressure. Consult your local Extension office for recommendations.
- Do not graze or feed soybean forage and hay to livestock
- Regardless of method of application (seed treatment, soil or foliar) DO NOT apply more than 0.2 lbs of active ingredient clothianidin per acre per year.
- Not for use on sorghum or rice seed in the state of California.
- Poncho 600 is to be used only as a dry-seeded rice seed treatment.
- Do not use Poncho 600 in areas where rice/crawfish aquaculture practices are in place or in or near fish farms, shrimp, prawn or crab pond or nursery operations.
- For canola and rapeseed: treated at the rate of XX fluid ounces per 100 lbs of seed will result in XX grams of clothianidin per 100 lbs of seed.
- For corn (field and pop): this seed has been treated at the rate of XX mg ai/seed.
- For corn, for the pest insect Southern corn billbug: Follow up foliar sprays of a locally registered insecticide may be needed under heavy pest pressure. Consult your local Extension Office for recommendations.
- For corn (sweet): this seed has been treated at the rate of XX mg ai/seed.
- For rice: treated at the rate of XX fluid ounces per 100 lbs of seed will result in XX lbs of clothianidin per 100 lbs of seed.
- For sorghum: treated at the rate of XX fluid ounces per 100 lbs of seed will result in XX grams of clothianidin per 100 lbs of seed.
- For soybean: this seed has been treated with 0.11 mg clothianidin/seed.
- For wheat, barley, buckwheat, millet (pearl and proso), oats, rye, teosinte, or triticale: treated at the rate of XX fluid ounces per 100 lbs of seed will result in XX grams of clothianidin per 100 lbs of seed.

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)
For MEDICAL, TRANSPORTATION or OTHER Emergencies ONLY Call 24
Hours A Day 1-800-334-7577

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.

Pesticide storage

Do not contaminate water, food or feed by storage or disposal. Store in a cool, dry secured storage area.

Pesticide disposal

To avoid waste, use all material in this container by application according to label directions. If waste cannot be avoided, offer remaining product to a waste facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

(continued)

Container handling

Non-Refillable Plastic Containers

Rigid, Non-refillable containers (equal to or less than 5 gallons)

Non-refillable container. Do not reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration.

Bayer

Poncho® 600

For: seed treatment use on canola, cereal grains (except corn, rice, sorghum and wild rice), rapeseed, corn, rice, sorghum, and soybean for the control of listed insect pests.

All seed treated with this product must be conspicuously coloured at the time of treatment.

ACTIVE INGREDIENT:

Clothianidin: 48.00%

OTHER INGREDIENTS: 52.00%

Contains 5.00 pounds clothianidin per U.S. gallon

TOTAL: 100.00%

EPA Reg. No. 264-789

KEEP OUT OF REACH OF CHILDREN CAUTION

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

Please refer to booklet for additional precautionary statements and directions for use.

FIRST AID

If Swallowed:	<ul style="list-style-type: none">• Immediately call a poison control center or doctor for treatment advice.• Do not induce vomiting unless told to by a poison control center or doctor.• Have person sip a glass of water if able to swallow.• Do not give anything by mouth to an unconscious person.
If Inhaled:	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or ambulance, and then give artificial respiration, preferably mouth-to-mouth if possible.• Call a poison control center or doctor for further treatment advice.
If on Skin or Clothing:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.

In case of emergency call toll free the Bayer CropScience Emergency Response Telephone No. 1-800-334-7577.

Have a product container or label with you when calling a poison control center or doctor, or going for treatment.

Note to Physician: No specific antidote is available. Treat the patient symptomatically.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

- Harmful if swallowed. Harmful if absorbed through the skin.
- Harmful if inhaled.
- Avoid contact with skin, eyes, or clothing. Avoid breathing spray mist.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read the entire label before using this product.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.

Pesticide storage

Do not contaminate water, food or feed by storage or disposal. Store in a cool, dry secured storage area.

Pesticide disposal

To avoid waste, use all material in this container by application according to label directions. If waste cannot be avoided, offer remaining product to a waste facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

Container handling

Non-Refillable Plastic Containers

Rigid, Non-refillable containers (equal to or less than 5 gallons)

Non-refillable container. Do not reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration.

Bayer CropScience LP
P.O. Box 12014, 2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
©2018 Bayer CropScience

US61379957C

171108C 05/18