

PROMOTE™



Contains halosulfuron, the active ingredient used in Permit®.

PROMOTE is a herbicide for selective pre-emergent and post-emergent control of listed weeds including both broadleaf weeds and nutsedge in labeled crop and non-crop sites.

ACTIVE INGREDIENT:	% BY WT.
Halosulfuron-methyl	75%
OTHER INGREDIENTS:	25%
TOTAL:	100%

EPA Reg. No. 91234-31

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail)

SEE BELOW FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

FIRST AID	
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call poison control center or physician for treatment advice.
IF SWALLOWED:	<ul style="list-style-type: none"> • Call poison control center or physician immediately for treatment advice. • Remove visible particles from mouth. • Have person rinse mouth thoroughly with water, spit out rinse water. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency medical assistance, call SafetyCall: 1-844-685-9173.	

For Chemical Emergency: Spill, Leak, Fire, Exposure, or Accident, Call CHEMTREC Day or Night Within USA and Canada:
1-800-424-9300 or +1 703-527-3887 (collect calls accepted)

Promote™ is not manufactured, or distributed by Gowan Company LLC, seller of Permit®.

Product of China



Manufactured for:
Atticus, LLC
940 NW Cary Parkway, Suite 200
Cary, NC 27513

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Causes moderate eye irritation. Harmful if swallowed. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- long-sleeved shirt and long pants, and
- shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROL STATEMENTS:

When handlers use closed systems, or enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS:

Users should:

- Remove PPE immediately after handling this product.
- Wash the outside of the gloves before removing.
- As soon as possible, wash thoroughly and change into clean clothing.
- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to non-target vascular plants. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Halosulfuron-methyl is known to leach through soil into ground water under certain conditions as a result of label use. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow.

PHYSICAL OR CHEMICAL HAZARDS

Do not mix with or allow to come into contact with oxidizing agents. A hazardous chemical reaction may occur.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. This product can only be used in accordance with the Directions for Use on this label or in separately published Atticus, LLC Supplemental Labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Read the entire label before using this product. Use only according to label instructions.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forest, nurseries and green houses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on the label about personal protective equipment (PPE), restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during this restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- Coveralls
- Shoes plus socks
- Chemical-resistant gloves, such as nitrile rubber, neoprene rubber or polyethylene.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep people and pets off treated areas until spray solution has dried.



Product Information

PROMOTE is a sulfonylurea herbicide that works by inhibition of acetolactate synthase (ALS). Many factors such as application rate, weed species, weed pressure, conditions of weeds including size and climatic factors impact the degree of weed control. Applications made to actively growing weeds at the early stages of development as described below will optimize performance. In post-emergent weed applications, early treatment is best to control the weeds vying (competing) with the crop. For residual control from early post-emergent treatments (in corn) a second application may be needed to control later germination of weeds.

PROMOTE is quick to act on targeted weeds by stunting growth allowing the crop to overtake the development of the targeted weeds. Once the development of the targeted weeds is stunted, the leaves and growing point begin to discolor and die.

Complete control typically occurs within 7 to 14 days depending on the weed size, species and growing conditions. Depending on the stage and development of the targeted weeds, control generally takes place in 7 to 14 days.

Resistant Management Guidance

PROMOTE is a Group 2 herbicide. Any weed population may contain or develop plants naturally resistant to **PROMOTE** and other Group 2 herbicides. Weed species with acquired resistance to Group 2 may eventually dominate the weed population if Group 2 herbicides are used repeatedly in the same field or in successive years as primary method of control for target species. This may result in partial or total loss of control of those species by **PROMOTE** or other Group 2 herbicides.

To delay resistance consider:

- Avoiding the consecutive use of **PROMOTE** or other target site action Group 2 herbicides that have a similar target site of action, on the same weed species.
- Using tank-mixtures or premixes with herbicides from different target site of action Groups as long as the involved products are all registered for the same use, have different sites of action, and are both effective at the tank mix or prepack rate on the weed(s) of concern.
- Basing herbicide use on a comprehensive IPM program.
- Monitoring treated weed populations for loss of field efficacy.
- Contacting your local extension specialist, certified crop advisor, and/or manufacture and/ or integrated weed management specialist for specific crops and resistant weed biotypes.

Mixing Instructions

PROMOTE is a water dispersible granule designed to be diluted with water at the rates listed in the specific crop use directions. Fill the spray tank with approximately ½ of the desired volume with water or carrier. With the agitation operating, add the specified amount of the formulation as listed in the targeted crop use directions. Complete the filling process while maintaining agitation. Remove the hose from the mixing tank immediately after filling to avoid siphoning back into the carrier source. Add nonionic surfactant and other spray additives as the last ingredients in the tank. Allow time to fully disperse.

Since this product forms a suspension in water, it is important to maintain good agitation during mixing and spraying. If the spray suspension is allowed to settle for a short period of time, be sure to agitate the spray suspension for a minimum 10 minutes. Apply spray solutions within 24 hours after mixing.

Spray Additives

Spray additives such as nonionic surfactant (NIS), or Crop Oil Concentrate (COC) and liquid nitrogen fertilizer (e.g. 28-0-0) are used with **PROMOTE** to improve performance. The typical nonionic surfactant contains a minimum of 80% NIS and is accepted by the EPA for use on food crops. The use rate is 0.25 to 0.5% NIS concentrate (1 to 2 quarts per 100 gallons of spray mixture). An alternative for the nonionic surfactant is a Crop Oil Concentrate. The typical Crop Oil Concentrate is a phytoblend oil (petroleum) or crop origin (vegetable) based product that containing a minimum 14% surfactant to allow it to be miscible with water. The use rate for the Crop Oil Concentrate is 1% vol/vol (1 gallon per 100 gallons of spray mixture). NIS or COC is the only spray additives required to improve efficacy. Do not use both NIS and COC in the spray mixture. Use liquid nitrogen for those tank mix partners which required a liquid nitrogen additive to improve performance. Consult the tank mixture partner's labels for specific additive requirements and interactions. In place of the liquid nitrogen fertilizer, a high quality, spray grade ammonium sulfate (e.g. 21-0-0) is used at a use rate of 2 to 4 pounds per acre. Use either NIS or COC in the spray mixture.

For specific details, consult the use directions in crop section of the label.

Use Rate Equivalency

Since **PROMOTE** contains 75% active ingredient per lb. of product, the following table expresses the use rate equivalency of oz. of this product in term of lb. active ingredient on a per acre basis.

oz. of Product	lb. Active Ingredient
½	0.0235
⅔	0.031
1	0.047
1 ⅓	0.062
2	0.094
2 ⅔	0.125
5 ⅓	0.250

Application Methods

Apply this product by ground or with aerial equipment to produce uniform coverage on growing weeds or soil to achieve consistent weed control.

Uniform, thorough spray coverage is important to achieve consistent weed control. Calibrate application equipment according to manufacturer's specifications. Use nozzle type arrangements that provide optimum spray distribution and maximum coverage while avoid contact to sensitive crop foliage.

Thoroughly clean application equipment immediately after use and prior to spraying a crop other than corn or grain sorghum. See Spray Equipment Cleanout section of this label for complete details.

Ground Applications

When **PROMOTE** is applied by ground equipment, use in a minimum of 10 gallons of water per acre for a broadcast application. In dense weed populations and thick canopy cover, higher spray volumes are necessary, e.g. 15 – 20 gallons of water per acre. Use the proper spray volume and nozzles that will ensure thorough and uniform coverage of the targeted weeds. Use directed applications to avoid contacting sensitive crop foliage. Select nozzles that will provide optimum spray volume, distribution and coverage at a pressure (psi) that minimizes spray drift. Inspect nozzle distribution during application to avoid streaking and overspray.

Aerial Applications

When **PROMOTE** is applied by air, use in a minimum 3 - 15 gallons of water per acre. Properly calibrate the spray equipment. Follow the Spray Drift Management guidelines presented below. Inspect nozzle distribution during application to avoid streaking, overspray and spray drift.

Spray Drift Management

Do not allow this product to drift onto neighboring crops or non-crop area or use in a manner or at a time other than in accordance with label directions because animal, plant or crop injury, illegal residues or other undesirable results may occur.

AVOIDING SPRAY DRIFT AT THE APPLICATION SITE IS THE RESPONSIBILITY OF THE APPLICATOR. The interaction of many equipment – and weather – related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions. Where states have more stringent regulations, they must be observed.

The following drift management requirements must be followed to avoid off-target drift movement from aerial application to agricultural field crops. These requirements do not apply to forestry applications, public health uses or to applications using dry formulations.

1. The distance of the outer most nozzles on the boom must not exceed ¾ the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees. Where states have more stringent regulations, they must be observed.



The importance of spray droplet size:

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential but may not prevent drift if applications are made improperly or under unfavorable environmental conditions (see the following "Wind", "Temperature and Humidity" and "Temperature Inversion" sections of this advisory).

Controlling initial droplet size:

- Volume – Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher flow rates produce larger droplets.
- Pressure – Use the lower spray pressures listed for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- Number of nozzles – Use the minimum number of nozzles that provide uniform coverage.
- Nozzle orientation – Orienting nozzles so the spray stream is released backwards, parallel to air stream will produce larger droplets than other orientations. Significant deflection from the horizontal will reduce droplet size and increase drift potential.
- Nozzle type- Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce larger droplets than other nozzle types.

Controlling placement of spray droplets:

- Boom Length – For some use patterns, to further reduce drift without shorting the swath width, reduce the effective boom length to less than ¾ of the wingspan or rotor length.
- Application height – Do not apply greater than 10 feet above the top of the tallest plants unless a great height is required for aircraft safety. Greater application heights result in greater droplet size reduction through evaporation and greater movement in air currents. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.
- Application speed – Slower aircraft speeds within a safe range will produce less air turbulence and fewer small droplets.

Swath adjustment – When applications are made with a cross-wind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicators must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distances should increase drift potential (wind speed, droplet size, etc.)

Key environment factors:

- Wind – Drift potential is the lowest between wind speeds of 2 to 10 mph. However, many factors including droplet size and equipment type determine drift potential at any given point. Application must be avoided when wind speeds are below 2 mph due to variable wind direction and high inversion potential. NOTE: Local terrain can influence wind patterns. Applicators must be familiar with local wind patterns and how they affect drift.
- Temperature and Humidity – when making applications in low relative humidity set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.
- Temperature Inversions – Do not apply during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable air currents that are common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke detector. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Sensitive areas:

Use pesticide products adjacent to sensitive areas only when there is minimal potential for drift or off-target movement, e.g. wind is blowing away from non-target crops, residential areas, known habitats for threatened or endangered species, etc.

In California (only), particularly sensitive crops are identified as cotton and prunes. In applications near these sensitive crops utilize the following buffer zones:

- Do not apply aerial applications within 4 miles of sensitive crops.
- Do not apply ground applications within 1 mile of sensitive crops except when wind direction during the application is away from sensitive crops. When wind direction during the ground application is away from sensitive crops, do not apply within 0.5 miles of sensitive crops.
- Do not apply Direct Dry Applications on rice by air within 360 feet of sensitive crops.

Spray Equipment Cleanout

The mix tank and spray equipment cleanout is an important stewardship activity to avoid injury to desirable crops. It is important to clean all mixing and spraying equipment immediately after use and before using pesticide products including **PROMOTE**. This is especially important prior to spraying a crop other than grain sorghum and corn.

To clean the spraying equipment, follow the procedure outlined below:

- Completely drain the mix tank and/or sprayer, and then wash thoroughly the tank, sprayer, boom and nozzles with clean water. Drain the system again.
- Fill the mixing or spray tank half full with clean water and add domestic ammonium, normally a 3% solution, at a dilution rate of 1% vol/vol ammonium or 1 gallon per 100 gallons of rinsate.
- Completely fill the tank(s) with additional clean water. Agitate and recirculate and flush out the boom and hoses. Let the system run for 10 – 15 minutes. Drain the system completely.
- Remove nozzles and screens and dislodge any visible solid material. Then soak them in a 1% vol/vol ammonium solution. Inspect the nozzles and screen and remove any visual residues.
- Repeat the above procedure for a second time.
- Flush the mix tank and/or sprayer, boom and hoses with clean water. Drain the system again and inspect for any visible residues. If present, repeat the cleaning cycle again.
- If the rinsate cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Tank Mixtures

To improve this product's effectiveness, apply in combination with other pesticide products that are registered for the same crop and application techniques.

A list of potential herbicide tank mixture partners is provided in the use direction section under each crop. This list is an example of products used but is not an all inclusive list. For current information on the best tank mixture partner in your area, consult with the local dealer, distributor or State Agricultural Extension service.

It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

If **PROMOTE** is to be tank mixed with other herbicides, conduct a compatibility test prior to mixing. Use a small container and mix all components in a small amount, usually 0.5 to 1 qt. of spray. Combine all products in the same ratio and order of addition as in the proposed spray mixture. Observe the mixture for indication of incompatibility which usual occurs in 10 to 30 minutes after mixing. If incompatibility is observed, try changing the order of addition of the components. The guideline on tank mixture partners is driven by formulation type. Start with wettable powders (WP's) including water soluble bags (WSB's), water dispersible granules (WDG's), suspension concentrated (SC's) or flowable (F's), all with very good agitation. Next follow with water miscible concentrates and emulsifiable concentrates (EC's) before adding drift control additives, nonionic surfactants (NIS's) or crop oil concentrates (COC's). After vigorous agitation, there must be a homogeneous suspension. Let the final tank mixture stand and observe for any rapid settling or floating of components. If any indications of physical incompatibility develop, do not use this mixture for spraying.



Application Restrictions

- Do not use air assisted (air blast) sprayers to apply this product.
- Do not apply this product through any type of irrigation system.
- Do not apply when wind speed exceeds 15 mph.
- Do not apply more than 2 ounces of this product per acre per 12-month period (includes applications to the crop and to row middles/furrows) on crops except on fallow ground, field corn, and sugarcane.
- Do not apply more than 2 2/3 oz. of this product (0.125 lb. active ingredient) per acre per use season on fallow ground, field corn, and sugarcane.
- Do not allow this product to drift outside of targeted area.
- Do not apply tank mixtures if the crop is under heavy stress due to drought, water-saturated soils, poor fertility (especially low nitrogen levels), hail, frost, insects or when the maximum daytime temperature is above 92° F.
- Do not use this product if the target weeds or crop are under stress due to drought, water saturated soils, low fertility (especially low nitrogen levels) or other poor growing conditions.

Application Precautions

- Avoid spraying when conditions favor rainfall or using overhead sprinkler irrigation within 4 hours of this application.
- Significant crop injury may occur when spray residue from broadcast application of this product over plastic mulch is concentrated in the plant hole by irrigation or rainfall. To minimize this potential injury, ensure that planting beds are crowned properly.
- Under cool and wet growing conditions that delay early seedling emergence, vigor or growth, this product may cause injury or crop failure. These conditions are likely to occur during the first planting of the season.
- Loss in effectiveness or crop injury may result if weeds are under drought, stress, disease or insect damage.
- The maturity of the treated crops may be delayed by use of this product.
- Soil or foliar-applied organophosphate insecticides applied on crops treated with this product, may increase the potential for crop injury and/or the severity of the crop injury.
- Increase in crop injury may result if the seeding depth is too shallow and excessive amounts of water (greater than 1 inch) from rainfall or sprinkler irrigation occurs.
- Use nozzles and pressures that minimize the production of fine particles that drift outside of the targeted area
- Apply this product to labeled crops (including cultivars and/or hybrids of these). However, not all hybrids/varieties have been tested for sensitivity to this product. For untested varieties, treat a small amount of the field and determine potential sensitivity to its use. To the extent consistent with applicable law, the user assumes responsibility for such use and any plant injury that may occur.
- Applications of this product may cause temporary yellowing or stunting of the crop.
- Observe resistant management guidelines, especially on tolerant weeds.
- In California and Arizona due to environmental conditions that delay degradation of this product, extend the crop rotation intervals on drip irrigated crops.
- When this product is applied over-the-top of a blooming crop, bloom loss may occur under certain environmental conditions.
- If rainfall or irrigation occurs within 4 hours after application, reduce effectiveness may occur.
- Avoid disturbing (e.g. cultivation) treated areas for at least 7 days following application.

For Best Performance

Many factors such as application rate, weed species, weed pressure, conditions of weeds including size and climatic conditions impact the degree of weed control. Applications made to actively growing weeds at the early stages of development as described below will optimize performance. In post-emergent weed applications, early treatment is best to control the weeds vying (competing) with the crop. For residual control from early post-emergent treatments (in corn) a second application may be needed to control later germination of weeds.

PROMOTE is quick to act on targeted weeds by stunting growth allowing the crop to overtake the development of the targeted weeds. Once the development of the targeted weeds is stunted, the leaves and growing point begin to discolor and die.

Complete control typically occurs within 7 to 14 days depending on the weed size, species and growing conditions. Depending on the stage and development of the targeted weeds, control generally takes place in 7 to 14 days.

When using spray additives, carefully follow the listed use instructions.

- In pre-emergence applications:
 - If the targeted weeds are present prior to crop emergence, use a nonionic surfactant identified in the "Spray Additives" section of the label.
 - For optimum pre-emergent weed control, activate the soil moisture.
 - Pre-emergent weed control is improved by incorporating this product with irrigations (1/4 - 1/2 inch maximum).
- In post-emergence applications:
 - Better control is obtained when applied early to actively growing, small (1-3 inches in height) broadleaf weeds. Large broadleaf weeds may not be adequately controlled.
 - Nutsedge plants are best controlled at the actively growing, 3 - 5 leaf stage.
 - After a post-emergence application, delay overhead sprinkler irrigation for 2 to 3 days.
 - If weeds are under drought, stress, disease, or insect damage, do not use.
- Under heavy weed infestation, use early before the weeds become too competitive with the crop.
- To control suppressed weeds, large weeds that exceed the size limitations, weeds that emerge after an application, or weed species not listed, cultivate the treated soil 7 - 10 days after a post-emergence application unless specified otherwise.
- Avoid disturbing (e.g. cultivation) treated areas for at least 7 days following application.
- Annual weeds may have multiple flushes of seedlings, or treated perennials may sometimes re-grow from underground stems or roots, depending upon rainfall and other environmental conditions. To maximize control of such weeds, apply a sequential application of this product.

Pre-emergent Weed Activity Table - PROMOTE - by Weed Species

Common Name	Scientific Name	Control	Suppression	Directions
Amaranth, Spiny	<i>Amaranth spinosus</i>	YES		Certain biotypes of this weed are known to be resistant to ALS herbicides. ¹
Cocklebur, common	<i>Xanthium strumarium</i>	YES		
Corn Spurry	<i>Spergula arvensis</i>	YES		
Dayflower	<i>Commelina erecta</i>	YES		
Eclipta	<i>Eclipta prostrata</i>	YES		
Flatsedge, Rice	<i>Cyperus iria</i>		YES	
Galinsoga	<i>Galinsoga</i>	YES		
Goosefoot		YES		
Groundsel, common	<i>Senecio vulgaris</i>	YES		
Horseweed/ Maretail	<i>Erigeron canadensis</i>	YES		
Jimsonweed	<i>Datura stramonium</i>	YES		
Kochia	<i>Kochia scoparia</i>	YES		Certain biotypes of this weed are known to be resistant to ALS herbicides. ¹
Ladysthumb	<i>Polygonum persicaria</i>	YES		
Lambsquarter, common	<i>Chenopodium album</i>	YES		
Mustard, wild	<i>Sinapis arvensis</i>	YES		
Nutsedge, Yellow	<i>Cyperus esculentus</i>		YES	Use higher specified rates for suppression
Nutsedge, Purple	<i>Cyperus rotundus</i>		YES	Use higher specified rates for suppression
Pigweed, redroot	<i>Amaranthus retroflexus</i>	YES		Certain biotypes of this weed are known to be resistant to ALS herbicides. ¹
Pigweed, smooth	<i>Amaranthus hybridus</i>	YES		Certain biotypes of this weed are known to be resistant to ALS herbicides. ¹
Purslane	<i>Portulaca oleracea</i>		YES	
Radish, wild	<i>Raphanus raphanistrum</i>	YES		
Ragweed, common	<i>Ambrosia artemisiifolia</i>	YES		Certain biotypes of this weed are known to be resistant to ALS herbicides. ¹
Shepardspurse	<i>capsella bursapastoris (L.) medicus</i>	YES		
Smartweed, Pennsylvania	<i>Polygonum pennsylvanicum</i>	YES		
Sunflower	<i>Helianthus annuus</i>	YES		
Velvetleaf	<i>Abutilon theophrasti</i>	YES		

¹ If ALS resistant weeds are present, use another mode of action herbicide registered on the crop against the target weeds alone or as a tank mixture partner.

It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Post-emergent Weed Activity Table - PROMOTE - by Weed Species

Common Name	Scientific Name	Control	Suppression	Directions
Amaranth, Spiny	<i>Amaranth spinosus</i>	YES		Certain biotypes of this weed are known to be resistant to ALS herbicides. ¹
Barnyardgrass	<i>Echinochloa crusgalli</i>	YES		
Bindweed	<i>Calyptegia sepium</i>	YES		
Burcucumber	<i>Sicyas angulatus</i>	YES	YES	
California Arrowhead	<i>Sagittaria ontariensis</i>	YES		1-1 ½ ounce rate required.
Cocklebur, common	<i>Xanthium strumarium</i>	YES		
Corn Spurry	<i>Spergula arvensis</i>	YES		
Cupgrass, Woolly	<i>Eriochloa villosa</i>	YES		
Dayflower	<i>Commelina erecta</i>		YES	
Dogbane Hemp	<i>Apocynum annabinum</i>		YES	
Eclipta	<i>Eclipta prostrata</i>		YES	
Flatsedge, Rice	<i>Cyperus iria</i>	YES		
Fleabane, Philadelphia	<i>Erigeron philadelphicus</i>	YES		
Foxtail, giant, yellow, green bristly		YES		
Galinsoga	<i>Galinsoga</i>	YES		
Golden Crownbeard	<i>Verbesina encliodes</i>	YES		
Goosefoot		YES		
Horsenettle	<i>Solanum carolinense</i>	YES		
Horsetail	<i>Equisetum</i>		YES	
Itchgrass	<i>Rottboellia cochinchinensis</i>	YES		
Jointvetch	<i>Aeschynomene</i>	YES		
Johnsongrass rhizome, seedling	<i>Sorghum halepense</i>	YES		
Kochia	<i>Kochia scoparia</i>		YES	Certain biotypes of this weed are known to be resistant to ALS herbicides. ¹
Ladysthumb	<i>Polygonum persicaria</i>	YES		
Mallow, Venice	<i>Hibiscus trionum</i>	YES		
Milkweed, Common	<i>Asclepias syriaca</i>		YES	
Milkweed, honeyvine	<i>Ampelamus albidus</i>		YES	
Millet, Wild Proso	<i>Panicum miliaceum</i>	YES		
Morningglory, ivyleaf	<i>Ipomoea hederacea</i>		YES	Use higher labeled rates for suppression.
Morningglory, Tall	<i>Ipomoea purpurea</i>		YES	Use higher labeled rates for suppression.
Mustard, wild	<i>Sinapis arvensis</i>	YES		
Nightshade, Black	<i>Solanum americanum</i>	YES		
Nutsedge, Yellow	<i>Cyperus esculentus</i>	YES		Heavy infestation requires sequential applications.
Nutsedge, Purple	<i>Cyperus rotundus</i>	YES		Heavy infestation requires sequential applications.
Oats		YES		
Panicum, Fall	<i>Panicum dichotomiflorum</i>	YES		
Panicum, Texas	<i>Panicum texanum</i>	YES		

(continued)

Common Name	Scientific Name	Control	Suppression	Directions
Passionflower, Maypop	<i>Passiflora incarnata</i>	YES		
Pigweed, redroot	<i>Amaranthus retroflexus</i>	YES		Certain biotypes of this weed are known to be resistant to ALS herbicides. ¹
Pigweed, smooth	<i>Amaranthus hybridus</i>	YES		Certain biotypes of this weed are known to be resistant to ALS herbicides. ¹
Pokeweed, common	<i>Phytolacca</i>	YES		
Quackgrass	<i>Elytrigia repense</i>	YES		
Radish, wild	<i>Raphanus raphanistrum</i>	YES		
Ragweed, common	<i>Ambrosia artemisiifolia</i>	YES		Certain biotypes of this weed are known to be resistant to ALS herbicides. ¹
Ragweed, giant	<i>Ambrosia trifida</i>	YES		Certain biotypes of this weed are known to be resistant to ALS herbicides. ¹
Redstem	<i>Ammania auriculata</i>	YES		1-1 ½ ounce rate required.
Ricefield Bulrush	<i>Scirpus mucronatus</i>	YES		Certain biotypes of this weed are known to be resistant to ALS herbicides. ¹
Ryegrass, Italian	<i>Lolium Multiflorum</i>	YES		
Sandbur		YES		
Sesbania, Hemp	<i>Sesbania exaltata</i>	YES		
Shattercane	<i>Sorghum bicolor</i>	YES		
Signalgrass, broadleaf		YES		
Shepardspurse	<i>capsella bursapastoris(L.) medicus</i>		YES	
Sida, prickly		YES		
Smallflower	<i>Umbrellaplant</i>	YES		1-1 ½ ounce rate required.
Smartweed, Pennsylvania	<i>Polygonum Pensylvanicum</i>	YES		
Sorghum Alnum		YES		
Thistle, Canada	<i>Cirsium arvense</i>	YES		
Sunflower	<i>Helianthus annuus</i>	YES		
Velvetleaf	<i>Abutilon theophrasti</i>	YES		

¹ If ALS resistant weeds are present, use another mode of action herbicide registered on the crop against the target weeds alone or as a tank mixture partner.

It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

PROMOTE Crop/Use Site Index

	Page
Corn:	
Field corn and Field corn grown for seed,	8-10
Sweet corn and popcorn.....	13-14
Cotton.....	7
Fallow ground.....	7
Grain sorghum (milo).....	10
Rice.....	11-12
Soybeans, seed.....	12
Sugarcane	13
Dry beans	7

The use rate for **PROMOTE** is expressed in terms of the oz. of this product by weight per acre as Rate Oz./Acre.

The pre-harvest interval (PHI) is the required days between the last application of **PROMOTE** and the harvesting of the crop.

For the minimum acceptable intervals between the last application of **PROMOTE** and the planting of a rotational crop, see the Crop Rotation Guideline section of this label.

If **PROMOTE** is utilized with a tank mixture partner(s), refer to the specific partner label(s) and observe all the precautionary statements and use directions including pre-harvest intervals, crop rotation restrictions, mixing and application instructions. Observe the most restrictive of the labeling limitations, precautions, directions and restrictions of all products used in mixtures.

Crop	Rate Oz./Acre	PHI	RESTRICTIONS
COTTON (except CA)	⅔ - 1 ½	28	Do not apply more than 1 ½ oz. of this product per acre per crop cycle, not to exceed 1 ½ oz. (0.062 lb. active ingredient) per acre per 12-month period.
For post-emergent weed control in emerged cotton, apply this product as a directed spray in hooded equipment. Make application any time after cotton emergence until row closure prohibits the use of hooded spray equipment. Use this product any time after cotton emergence until row closure inhibits use of hooded spray equipment. The applicator is responsible for maintaining proper spray speed and equipment position so spray mist does not contact cotton plants.			

Crop	Rate Oz./Acre	PHI	RESTRICTIONS
DRY BEANS	½ - 1		Do not apply more than 1 oz. of this product per acre per crop cycle, not to exceed 2 oz. (0.094 lb. active ingredient) per acre per 12- month period (includes applications to the crop and to Row Middles/Furrows).

For spray applications, cover the treatment area with sufficient water to provide uniform coverage and distribution of the spray mixture to the weeds or soil. For ground equipment, use a minimum of 15 gallons of water per acre.

Direct-Seeded Pre-emergence:

Use Rate: ½ - ⅔ oz.

Use this product after planting, but before cracking. For lighter textured soils with low organic matter, use the lower rate.

Row Middle/Furrow Applications:

Use Rate: ½- 1 oz.

Apply this product between rows of crop for the control of nutsedge and labeled broadleaf weeds. Avoid contact of this product with the planted crop. If plastic is used on the planted row, adjust equipment to keep the application off the plastic. Adjust the rate and spray volume proportionally to the actual treated area.

Tank Mixture Partner

A tank mixture of **PROMOTE** partnered with Eptam® 7-E will provide a broader spectrum of weed control than either product used separately.

It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Use **PROMOTE** at a rate of ½ - ⅔ oz. with labeled rate of Eptam® 7-E and incorporate into the soil at a depth of approximately 2 inches before planting. If any crust appears on the soil, break it up by lightly rotary hoeing during or shortly after the emergence of the beans.

For lighter textured soils with low organic matter, use the lower rate.

Crop	Rate Oz./Acre	PHI	RESTRICTIONS
FALLOW GROUND	⅔ - 1 ½		Do not make more than 2 applications per use season. Do not apply more than 2 ⅔ oz. of this product (0.125 lb. active ingredient) per acre per use season.

Apply this product as a broadcast spray to fallow ground. For spray applications, cover the treatment area with sufficient water to provide uniform coverage and distribution of the spray mixture to the weeds or soil.



Crop	Rate Oz./Acre	PHI	RESTRICTIONS
FIELD CORN AND FIELD CORN GROWN FOR SEED	2/3 - 1 1/2	30	Do not make more than 2 applications per crop use season. Do not apply more than 2 2/3 oz. of this product (0.125 lb. active ingredient) per acre per use season. After application to foliage, allow 30 days before grazing domestic livestock, harvesting forage, or harvesting silage.

If used alone, apply a broadcast spray over-the-top or with drop nozzles from the spike through lay-by stage of field corn. For large corn or dense competing canopy, use drop nozzles. Avoid spraying an excessive amount directly over the rows and into the whorl of the corn stalk.

Corn Weed Height Activity Table

Weed Activity	Control		Suppression	
	2/3 oz.	1 - 1 1/2 oz.	2/3 oz.	1 - 1 1/2 oz.
Weed Height	Inches	Inches	Inches	Inches
Burcucumber			1 - 3	4 - 12
Cocklebur, common	1 - 9	9 - 14		
Fleabane, Philadelphia	1 - 3			
Kochia ¹	1 - 3			3 - 6
Lambsquarter, common			1 - 2	
Mallow, Venice	1 - 3	4 - 12		
Milkweed, common			3 - 5	6 - 12
Milkweed, honeyvine		1 - 6	1 - 3	
Morningglory				1 - 3
Mustard, wild		4 - 6		
Nutsedge: yellow ²	3 - 6	3 - 12		
purple	3 - 6	3 - 12		
Passionflower, maypop	1 - 3			
Pigweed, redroot ^{1,3}	1 - 3	4 - 6		
Pokeweed, common	1 - 6			
Radish, wild		4 - 6		
Ragweed: common ¹	1 - 9	9 - 12		
Giant ¹	1 - 3	4 - 6		
Smartweed, Pennsylvania	1 - 2			
Sunflower, common	1 - 12	12 - 15		
Velvetleaf ³	1 - 9	9 - 12		

¹ See Pre-emergent and Post-emergent Weed Activity Tables

² Heavy infestations of nutsedge require sequential applications. To prevent nutsedge from competing with the crop an earlier application is required.

³ For large velvetleaf and pigweed control apply with liquid nitrogen fertilizer (2 to 4 quarts per acre) plus crop oil concentrate or nonionic surfactant is suggested.

Tank Mixture Partners

It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

2,4-D	See tank mixture partner label for its rates. NIS Apply broadcast spray on corn up to 8 inches tall. If corn exceeds 8 inches, use directed spray with drop nozzles. Broadleaf weeds. Avoid sprays onto corn leaves just after unfolding, as injury may occur. Apply during the period from corn emergence through the 5 leaf stage or 8 inches tall, whichever comes first.
Accent® Herbicide	See tank mixture partner label for its rates. COC or NIS or ammonium nitrogen fertilizer (e.g. 28%). Apply broadcast spray or with drop nozzles on emerged corn up to 24 inches tall. (free standing) For corn 24 to 36 inches tall, use directed spray with drop nozzles. Annual broadleaf weeds and annual grasses. Avoid sprays directly into the whorls of large cornstalks. Refer to Accent® label for use restrictions on corn varieties.
Accent Gold® Herbicide	See tank mixture partner label for its rates. COC or ammonium nitrogen fertilizer (e.g. 28%). Apply broadcast spray on corn up to 12 inches tall. Annual broadleaf weeds and annual grasses. Do not apply to seed corn.
Atrazine 4L Herbicide	See tank mixture partner label for its rates. COC Apply broadcast spray on corn up to 12 inches tall. Apply when broadleaf weeds are small (3 inches or less). Post-emergence control of labeled broadleaf weeds. Aids in the burndown and control of many grass weeds (1.5 inches or less) which have escaped pre-emergence herbicide treatments. Atrazine mixtures may result in reduced control (antagonism) of larger broadleaf weeds. Smaller weeds are easier to control.
Atrazine 90DF Herbicide	See tank mixture partner label for its rates. COC Apply broadcast spray on corn up to 12 inches tall. Apply when broadleaf weeds are small (3 inches or less). Post-emergence control of labeled broadleaf weeds. Aids in the burndown and control of many grass weeds (1.5 inches or less) which have escaped pre-emergence herbicide treatments. Atrazine mixtures may result in reduced control (antagonism) of larger broadleaf weeds. Small weeds are easier to control.
Banvel® Herbicide or Clarity® Herbicide (dicamba)	See tank mixture partner label for its rates. NIS Apply broadcast spray on corn from emergence up to 36 inches tall. Use lower Banvel rates or directed sprays on corn taller than 8 inches. Broadleaf weeds. Avoid direct sprays into the whorls of large cornstalks. Do not make applications after corn exceeds 36 inches or 15 days before tassel emergence, whichever comes first. COC may cause crop injury, especially with higher Banvel® or Clarity® rates.

(continued)

Basis® Gold Herbicide	See tank mixture partner label for its rates. COC or NIS or ammonium nitrogen fertilizer (e.g. 28 %). Apply broadcast spray on corn up to 12 inches tall. Broader spectrum. Do not apply to seed corn.
Beacon® Herbicide	See tank mixture partner label for its rates. COC or NIS or ammonium nitrogen fertilizer (e.g. 28 %). Apply broadcast spray or with drop nozzles on corn from 4 - 20 inches tall. For corn 20 - 36 inches tall to pre-tassel, use drop nozzles. Broader spectrum. Avoid spraying directly into whorls of larger corn. See your dealer or seed supplier representative for a list of susceptible hybrids.
Buctril® Herbicide	See tank mixture partner label for its rates. NIS Apply broadcast spray on corn up to tassel emergence. For post-emergence control of annual broadleaf weeds. Leaf burn may occur. Use of COC or ammonium nitrogen fertilizer (e.g. 28%) may cause additional leaf burn.
Buctril® Herbicide plus Atrazine	See tank mixture partner label for its rates. NIS Apply broadcast spray on corn up to 12 inches tall. For post-emergence control of annual broadleaf weeds. Leaf burn may occur. Use of COC or ammonium nitrogen fertilizer (e.g. 28%) may cause additional leaf burn.
Callisto® 4L Herbicide	See tank mixture partner label for its rates. COC or ammonium nitrogen fertilizer (e.g. 28 %). Apply broadcast spray or with drop nozzles on seed or field corn up to 30 inches tall or 8 leaf collars, whichever is more restrictive. Broader spectrum.
Distinct® Herbicide	See tank mixture partner label for its rates. NIS Apply broadcast spray or with drop nozzles on corn 4 - 36 inches tall, e.g. V2 to V10 stage or 15 days prior to tassel emergence, whichever comes first. For corn taller than 20 inches, use drop nozzles. Broader spectrum. Avoid sprays directly into the whorls of large cornstalks. Do not use COC.
Glyphosate (various formulations)	See tank mixture partner label for its rates. NIS or spray grade ammonium sulfate at 17 lb. /100 gal. Apply broadcast spray or with drop nozzles on Glyphosate Tolerant (GT) field corn up to 30 inches tall or 8 leaf collars, whichever is more restrictive. For GT field corn between 24 - 36 inches, use drop nozzles. For corn taller than 20 inches, use drop nozzles. For burndown of emerged annual grasses, broadleaf weeds and nutsedge. Check product formulation label for specific restrictions. For use ONLY on corn hybrids tolerant to glyphosate herbicide.
Glyphosate (various formulations)	Use PROMOTE at 2/3 oz. For glyphosate, see product formulation label. NIS Apply broadcast spray. For pre-plant burndown of emerged annual grasses, broadleaf weeds and nutsedge. To improve burndown of broadleaf weed control use dicamba or 2,4-D. Use only on Pioneer IR corn hybrids.
Impact® 2.8L Herbicide	See tank mixture partner label for its rates. NIS (preferred) or COC or ammonium nitrogen fertilizer (e.g. 28%). Apply broadcast spray or with drop nozzles on seed or field corn up to 36 inches tall. For a density canopy, drop nozzles are preferred. Broader spectrum.
Liberty® 1.67L Herbicide	See tank mixture partner label for its rates. Spray grade ammonium sulfate (17 lbs./100 gallons of spray mix). Apply broadcast spray or with drop nozzles on field corn up to 24 inches tall or 7 leaf collars whichever is more restrictive. For field corn taller than 24 inches up to 36 inches tall, use drop nozzles. Broadleaf weeds and annual grasses. Do not add NIS or COC. For use ONLY on corn hybrids tolerant to Liberty® Herbicide.
Marksman® Herbicide	See tank mixture partner label for its rates. NIS Apply broadcast spray on corn up to 8 inches tall. Broader spectrum. COC may cause crop injury.
Option® 35WDG Corn Herbicide	See tank mixture partner label for its rates. COC or ammonium nitrogen fertilizer (e.g. 28%) or spray grade ammonium sulfate (17 lb. /100 gal.). Apply broadcast spray or with drop nozzles on field corn 4 - 16 inches tall e.g. V2 to V6. For field corn taller than 16 up to 36 inches e.g. V6 to V10, use drop nozzles. Broader spectrum. Do not apply Option® to seed corn. Avoid spraying directly into the whorls of large cornstalks.
Status® Herbicide	See tank mixture partner label for its rates. NIS Apply broadcast spray or with drop nozzles on corn up to 20 inches tall. For corn taller than 20 inches use drop nozzles. Broader spectrum. Do not use COC.
Steadfast® 75DF Herbicide	See tank mixture partner label for its rates. COC (preferred) or NIS or ammonium nitrogen fertilizer (e.g. 28%) or spray grade ammonium sulfate (17 lbs. /100 gal.). Apply broadcast spray or with drop nozzles on field corn up to 20 inches tall or 6 leaf collars whichever is more restrictive. Broader spectrum. Avoid spraying directly into the whorls of large cornstalks. Do not apply to seed corn.
Soil Residual Tank Mix Partners	Use PROMOTE at 2/3 oz. See tank mixture partner label for its rates. Micro-Tech® or Bullet® or Harness® Xtra or Harness® Xtra 5.6L or Degree® or Degree Xtra®. NIS (1 qt./100 gallons of spray) and 28% nitrogen fertilizer (4 gal/100 gallons of spray). Apply as broadcast spray in 15 - 30 gallons of spray/acre to emerged grasses at the 2 leaf stage or less and on corn less than 11 inches tall (5 inches tall for Micro-Tech® and Bullet®). For early post-emergence control of additional small broadleaf, nutsedge and emerged grasses and pre-emergence control or reduced competition of annual broadleaf weeds and grasses as listed on the partner product label. To control emerged lambsquarter less than 4 inches tall, use Banvel® or Clarity® [dicamba].

(continued)

Soil Residual Tank Mix Partners (continued)	<p>Use PROMOTE at $\frac{2}{3}$ oz. and Accent[®]. See tank mixture partner label for its rates. Micro-Tech[®] or Bullet[®] or Harness[®] Xtra or Harness[®] Xtra 5.6L or Degree[®] or Degree Xtra[®]. NIS (1 qt./100 gallons of spray) and 28% nitrogen fertilizer (4 gal/100 gallons of spray). Apply as broadcast spray in 15 – 30 gallons of spray/acre to emerged grasses at the 2 leaf stage or less, foxtail less than 2 inches tall and on corn less than 11 inches tall (5 inches tall for Micro-Tech[®] and Bullet[®]). For early post-emergence control of additional small broadleaf and emerged grasses, including foxtail and pre-emergence control or reduced competition of annual broadleaf weeds and grasses as listed on the partner product label. To control emerged lambsquarter less than 4 inches tall, use Banvel[®] or Clarity[®] [dicamba].</p> <p>Use PROMOTE plus Accent[®], Beacon[®], Option[®] or Steadfast[®]. See tank mixture partner label for its rates. Alachlor, acetochlor, metolachlor and dimethenamid. NIS (1 qt./100 gallons of spray) and 28% nitrogen fertilizer (4 gal/100 gallons of spray). Apply as broadcast spray in 15 – 30 gallons of spray/acre to emerged foxtails and other grasses. For early post-emergence and residual control of emerged foxtails and other grass weeds in seed and field corn. Provides residual control or reduced competition of annual grasses and certain broadleaf weeds as listed on the specific herbicide labels. Follow all directions and restrictions on maximum corn height for post applications on this label and the tank mix partner's label. Use the more restrictive guidelines.</p>
Pioneer IR Field Corn Hybrids	<p>Use PROMOTE at 1½ - 2 oz. Apply broadcast spray to soil. For residual control of velvetleaf, common cocklebur, common lambsquarter, common ragweed, pigweed, smartweed, sunflower and other difficult to control weeds. Use only on Pioneer IR corn hybrids.</p>
Pre-plant, Pre-emergent.	<p>Use PROMOTE plus Accent[®], Beacon[®], Option[®] or Steadfast[®]. See tank mixture partner label for its rates. Such as Harness[®], Harness[®] Xtra, Harness[®] Xtra 5.6L, Degree[®], Degree Xtra[®], Micro-Tech[®], Bullet[®], Lasso[®], alachlor, acetochlor, metolachlor and dimethenamid. NIS (1 qt./100 gallons of spray) and 28% nitrogen fertilizer (4 gal/100 gallons of spray). Apply as broadcast spray in 15 – 30 gallons of spray/acre. Apply as an early pre-plant surface-applied, pre-plant incorporated or pre-emergence treatment. For effective broadleaf control in tank partner combinations with pre-emergence grass herbicides across all tillage systems. Follow all directions and restrictions on this label and the tank mix partner's label. Use the more restrictive guidelines.</p>

Crop	Rate Oz./Acre	PHI	RESTRICTIONS
Grain Sorghum (Milo)	$\frac{2}{3}$ - 1	30	Do not make more than 1 application per use season. Do not apply more than 1 oz. of this product (0.047 lb. active ingredient) per acre per use season. Following application to foliage, allow 30 days before grazing domestic livestock, harvesting forage, or harvesting silage.

If used alone, apply at the 2-leaf through lay-by stage of grain sorghum (before the grain head emerges).

If grain sorghum is under stress, temporary stature reduction occurs to the crop following application of this product. After application this effect will be evident in 7 – 10 days but under normally growing conditions will quickly recover.

Sorghum Weed Height Activity Table

Weed Activity	Control		Suppression
	$\frac{2}{3}$ oz.	1 oz.	$\frac{2}{3}$ oz.
Rate of Product	$\frac{2}{3}$ oz.	1 oz.	$\frac{2}{3}$ oz.
Weed Height	Inches	Inches	Inches
Burcucumber			1 - 3
Cocklebur, common	1 - 9		
Fleabane, Philadelphia	1 - 3		
Kochia ¹	1 - 3		
Lambsquarter, common			1 - 2
Mallow, Venice	1 - 3		
Milkweed, common			3 - 5
Milkweed, honeyvine			1 - 3
Nutsedge: yellow ²	3 - 6	3 - 12	
purple	3 - 6	3 - 12	
Passionflower, maypop	1 - 3		
Pigweed, redroot	1 - 3		
Pokeweed, common	1 - 6		
Ragweed: common	1 - 9		
Giant	1 - 3		
Smartweed, Pennsylvania	1 - 2		
Sunflower, common	1 - 12		
Velvetleaf	1 - 9		

¹ See Pre-emergent and Post-emergent Weed Activity Tables.

² Heavy infestations of nutsedge require sequential applications. To prevent nutsedge from competing with the crop an earlier application is required.

Tank Mixture Partners

It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

2,4-D	<p>See tank mixture partner label for its rates. NIS Apply broadcast spray on sorghum 6 to 15 inches tall. If sorghum exceeds 8 inches, use directed spray with drop nozzles and avoid spray on foliage. Broadleaf weeds. Do not treat during the boot, flower or dough stage. Do not make applications when sorghum exceeds 15 inches.</p>
Atrazine 4L Herbicide	<p>See tank mixture partner label for its rates. COC Apply broadcast spray on sorghum up to 12 inches tall. Apply when broadleaf weeds are small (3 inches or less). Post-emergence control of labeled broadleaf weeds. Aids in the burndown and control of many grass weeds (1.5 inches or less) which have escaped pre-emergence herbicide treatments. Atrazine mixtures may result in reduced control (antagonism) of larger broadleaf weeds. Smaller weeds are easier to control.</p>
Buctril [®] Herbicide	<p>See tank mixture partner label for its rates. NIS Apply broadcast spray on sorghum. For post-emergence control of annual broadleaf weeds.</p>
Buctril [®] Herbicide plus Atrazine	<p>See tank mixture partner label for its rates. NIS Apply broadcast spray on sorghum. For post-emergence control of annual broadleaf weeds.</p>



Crop	Rate Oz./Acre	PHI	RESTRICTIONS
RICE	$\frac{2}{3}$ - 1 $\frac{1}{2}$	48*	<p>Do not make more than 3 applications (including pre-plant and at-planting applications) per year.</p> <p>Do not apply more than 1 $\frac{1}{2}$ oz. of this product (0.062 lb. active ingredient) per acre per use year.</p> <p>After application to foliage, allow 30 days before grazing domestic livestock, harvesting forage, or harvesting silage.</p> <p>*Do not apply within 69 days of harvest in California. For Direct Dry Applications by air:</p> <p>Do not apply to dry rice fields.</p> <p>Apply aerial applications at a maximum of no greater than $\frac{1}{2}$ the wing span.</p> <p>Do not use a swath width greater than 120 feet.</p> <p>Do not mix this product with any other additives except as directed by this label.</p> <p>Do not apply within 360 feet of sensitive crops.</p> <p>Do not apply when wind speed is less than 3 mph or exceeds 15 mph.</p>

Rice Weed Height Activity Table

Weed Activity	Control		Suppression	
	Rate of Product	Weed Height	Rate of Product	Weed Height
	$\frac{2}{3}$ oz.	1 - 1 $\frac{1}{2}$ oz.	$\frac{2}{3}$ oz.	1 - 1 $\frac{1}{2}$ oz.
	Inches	Inches	Inches	Inches
Burcucumber			1 - 3	4 - 12
California Arrowhead		Yes		
Cocklebur, common	1 - 9	9 - 14		
Dayflower	1 - 2	3 - 4		
Eclipta	1 - 4	4 - 8		
Flatsedge rice	1 - 9	9 - 12		
Fleabane, Philadelphia	1 - 3			
Jointvetch	1 - 2	3 - 4		
Kochia ¹	1 - 3			3 - 6
Lambsquarter, common			1 - 2	
Mallow, Venice	1 - 3	4 - 12		
Milkweed, common			3 - 5	6 - 12
Milkweed, honeyvine		1 - 6	1 - 3	
Morningglory				1 - 3
Mustard, wild		4 - 6		
Nutsedge: yellow ²	1 - 6	6 - 12		
purple	1 - 6	6 - 12		
Passionflower, maypop	1 - 3			
Pigweed, redroot	1 - 3	4 - 6		
Pokeweed, common	1 - 6			
Radish, wild		4 - 6		
Ragweed: common	1 - 9	9 - 12		
Giant	1 - 3	4 - 6		
Redstem	1 - 3	Yes		
Ricefield Bulrush		Yes		
Sesbania Hemp	1 - 3	3 - 6		
Sida, Prickly	1 - 2	3 - 4		
Smallflower Umbrellaplant		Yes		
Smartweed, Pennsylvania	1 - 2			
Sunflower, common	1 - 12	12 - 15		
Velvetleaf ³	1 - 9	9 - 12		

¹ See the Post-emergent Weed Activity Table.

² Heavy infestations of nutsedge require sequential applications. To prevent nutsedge from competing with the crop an earlier application is required.

³ For large velvetleaf and pigweed control, apply with liquid nitrogen fertilizer (2 - 4 qts./acre).

Pre-plant, at-plant, post-emergent and prior to emergence of rice through permanent flood:

Use $\frac{2}{3}$ - 1 $\frac{1}{2}$ oz. of this product per acre per use season.

Apply foliar applications of this product at the 3 - 5 leaf stage of rice when weeds have 2 - 4 leaves. For foliar applications, use nonionic surfactant at rate of 0.25 - 0.5% in the spray mixture.

For aerial foliar applications, use a minimum of 3 - 15 gallons of water per acre. For ground foliar applications, use a minimum of 10 gallon of water per acre.

After mixing, apply spray suspensions the same day for best results.

Precautions:

Best control of emerged weeds with foliar applications occurs when 70% - 80% of the weed foliage is exposed. For best control of submerged weeds, apply when weeds have 2 leaves or less.

Check spray drift management section of this label.

Following the foliar applications of this product, do not reintroduce water into rice fields or checks for at least 24 hours. To improve the spectrum of weed control, tank mix this product with Shark[®].

It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Sequential Applications:

To improve the spectrum of weed control, apply this product sequentially with Bolero[®], Clincher[®], Regiment[®] or Shark[®].

Direct Dry Application:

Apply this product post flood as a dry broadcast application at a rate of 1 - 1 $\frac{1}{2}$ oz. per acre per use season. When weeds have 2 leaves or less, apply the dry broadcast treatment of this product at 1 - 2 leaf stage of rice.

Water levels in rice fields and checks should remain static (3 - 6 inch depth) after dry broadcast applications of this product. Do not reintroduce water into rice fields or checks for at least 5 days after dry broadcast treatments.

Rice fields and checks may be irrigated to maintain water level, but this may reduce weed control. Co-application with Shark is allowed.



Tank Mixture Partners	
It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.	
Glyphosate (various formulations)	Use PROMOTE at $\frac{2}{3}$ oz. See Glyphosate label for its rates. NIS Broadcast spray. For pre-plant or at-planting burndown of emerged annual grasses, broadleaf weeds and nutsedge. If applied as a pre-plant burn down treatment, consult the Crop Rotational Guidelines of this product and the Glyphosate label.
Stam® M4 and Propanil 4E (propanil)	Use PROMOTE at $\frac{2}{3}$ - 1 $\frac{1}{2}$ oz. See propanil labels for its rates. Broader spectrum weed control. If applied as a pre-plant burn down treatment, consult the Crop Rotational Guidelines.

Crop	Rate Oz./Acre	PHI	RESTRICTIONS
SOYBEANS, Soybean Seed (Except CA)	$\frac{2}{3}$ - 1 $\frac{1}{2}$	88	Do not make more than 1 application per year. Do not apply more than 1 $\frac{1}{2}$ oz. of this product (0.062 lb. active ingredient) per acre per year. Do not apply to frozen ground. After application to foliage, allow 30 days before grazing domestic livestock or harvest forage, silage and hay. Do not apply this product if plans include planting Adzuki beans since unacceptable crop injury may occur. Do not use post-emergence applications of this product to straight Roundup Ready or conventional soybean varieties as severe crop injury will occur. Do not use more than $\frac{2}{3}$ oz. of this product in post-emergence application to sulfonyl-urea tolerant soybeans (STS).

Pre-plant Burndown – Fall Application:
Use Rate: $\frac{2}{3}$ - 1 $\frac{1}{2}$ oz.
For control or suppression of listed broadleaf winter annual weeds prior to planting soybeans in the following Spring, apply this product as a fall burndown herbicide and/or preventative application.
If listed broadleaf weed are visible, use a high quality crop oil concentrates (1 - 2% vol/vol) and granular AMS (2 - 4 lbs./A) or UAN (1 -2% vol/vol) to the spray suspension to improve performance.
Apply this product from anytime from after harvest up until the ground freezes. Apply this product by ground or air.
In research trials, no instances of crop injury from Fall applied applications have occurred but not all soybean varieties have been tested for crop tolerance to halosulfuron-methyl. For the latest halosulfuron-methyl tolerance information, consult the local seed agronomists and seed supplier.
For broadleaf winter annual weed activity, this product must contact the emerged weeds and reach the soil surface.
In reduce tillage systems to maintain the active ingredient in the top layer of soil where weed seeds germinate, apply this product after fall chisel, disking etc.

Tank Mixture Partners	
It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.	
2,4-D Amine or LV ester (various formulations)	See tank mixture partner label for its rates. For broader spectrum control in pre-plant burndown of emerged annual broadleaf weeds or under heavy weed infestation.
Glyphosate (various formulations)	See tank mixture partner label for product rates. For control in pre-plant burn down of emerged grass weeds.

Pre-emergence or Pre-plant Spring Application to Soybean Varieties Tolerant to Sulfonyl-urea Herbicides (STS) Only:
Use Rate: $\frac{2}{3}$ oz.
For contact and residual control or suppression of listed broadleaf winter and early germinating summer annual weeds, apply this product from 21 days before planting until prior to emergence (i.e. cracking).
For best performance, apply this product to actively growing weeds free from environmental stress.
If listed broadleaf weed are visible, use a high quality crop oil concentrates (1 % vol/vol) and granular AMS (2 - 4 lbs./A) or UAN (1 -2% vol/vol) to the spray suspension to improve performance.
For use on any soybean varieties tolerant to sulfonyl-urea herbicides (STS) unless prohibited by the seed supplier.
In research trials, no instances of crop injury from Spring pre-plant or pre-emergence applied applications have occurred but not all sulfonyl-urea tolerant soybean (STS) varieties have been tested for crop tolerant to halosulfuron-methyl. For the latest halosulfuron-methyl tolerance information, consult the local seed agronomists and seed supplier.
In reduce tillage systems to maintain the active ingredient in the top layer of soil where weed seeds germinate, apply this product after all tillage operations.

Tank Mixture Partners	
It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.	
2,4-D LV ester (various formulations)	See tank mixture partner label for its rates. For control in early germinating summer weeds.
Glyphosate (various formulations)	See tank mixture partner label for product rates. For broader spectrum control in pre-plant burndown of emerged annual broadleaf weeds or grasses.

Post-emergence Application to Soybean Varieties Tolerant to Sulfonyl-urea Herbicides (STS) Only:
Use Rate: $\frac{2}{3}$ oz.
For contact and residual control of listed broadleaf weeds and nutsedge, apply this product post-emergent from V1 through R2 stages of sulfonyl-urea tolerant soybean (STS) varieties only.
If the tolerant soybean variety is also stacked with glyphosate or glufosinate tolerant trait, then glyphosate or glufosinate respectively may be used as a tank mixture partner.
For best performance, apply this product to actively growing weeds free from environmental stress.
For use on any soybean varieties tolerant to sulfonyl-urea herbicides (STS) unless prohibited by the seed supplier.
Always use a NIS(1 to 2 qts./100 gallons of spray) or high quality crop oil concentrates (1 % vol/vol) and granular AMS (2 - 4 lbs./A) or UAN (1 -2% vol/vol) to the spray suspension to improve performance.
Some phytotoxicity from post-emergent applications may occur on susceptible sulfonyl-urea tolerant soybeans (STS) varieties. These symptoms may include stunting (seen as reduction in leaf size or internodal length), yellowing of leaves and or red veins and necrosis of leaves and petioles. Sulfonyl-urea tolerant soybeans (STS) that have exhibit these symptoms tend to recover after the product is metabolized by the plant. Soybean injury is most noticeable when the plants are under environmental stress conditions such as hot, humid conditions, or wide fluctuations in climatic conditions, drought, etc.
For the latest halosulfuron-methyl tolerance information, consult the local seed agronomists and seed supplier.

Tank Mixture Partners	
It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.	
Glyphosate (various formulations)	See tank mixture partner label for product rates. For broader spectrum control of emerged annual broadleaf and grass weeds or under heavy weed infestation.



Crop	Rate Oz./Acre	PHI	RESTRICTIONS
SUGARCANE	¾ - 1 ½	30	Do not make more than 3 applications (including pre-plant applications) per year. Do not apply more than 2 ¾ oz. of this product (0.125 lb. active ingredient) per acre per year. After application to foliage, allow 30 days before grazing domestic livestock, harvesting forage, or harvesting silage.

If used alone, apply this product prior to planting, prior to emergence or after the emergence of the sugarcane and until row closure. Use mechanical cultivation to control weed species not on this label. If mechanical cultivation is used, apply a sequential treatment to control weeds in areas of disturbed soil.

Sugarcane Weed Height Activity Table

Weed Activity	Control		Suppression	
	¾ oz.	1 - 1 ½ oz.	¾ oz.	1 - 1 ½ oz.
Rate of Product	¾ oz.	1 - 1 ½ oz.	¾ oz.	1 - 1 ½ oz.
Weed Height	Inches	Inches	Inches	Inches
Burcucumber			1 - 3	4 - 12
Cocklebur, common	1 - 9	9 - 14		
Fleabane, Philadelphia	1 - 3			
Kochia ¹	1 - 3			3 - 6
Lambsquarter, common			1 - 2	
Mallow, Venice	1 - 3	4 - 12		
Milkweed, common			3 - 5	6 - 12
Milkweed, honeyvine		1 - 6	1 - 3	
Morningglory				1 - 3
Mustard, wild		4 - 6		
Nutsedge: yellow ²	3 - 6	3 - 12		
purple	3 - 6	3 - 12		
Passionflower, maypop	1 - 3			
Pigweed, redroot ³	1 - 3	4 - 6		
Pokeweed, common	1 - 6			
Radish, wild		4 - 6		
Ragweed: common	1 - 9	9 - 12		
Giant	1 - 3	4 - 6		
Smartweed, Pennsylvania	1 - 2			
Sunflower, common	1 - 12	12 - 15		
Velvetleaf ³	1 - 9	9 - 12		

¹ See Pre-emergent and Post-emergent Weed Activity Tables.

² Heavy infestations of nutsedge require sequential applications. To prevent nutsedge from competing with the crop an earlier application is required.

³ For large velvetleaf and pigweed control, apply with liquid nitrogen fertilizer (2 - 4 qts. /acre) plus NIS or COC.

Tank Mixture Partners

It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

2,4-D amine	See tank mixture partner label for its rates. NIS Apply to sugarcane before crop emergence or post-emergence until 6 weeks before harvest. Broadleaf weeds. Do not make more than 3 applications per year.
Asulam Herbicide	See tank mixture partner label for its rates. NIS or COC Apply to sugarcane before crop emergence or post-emergence until 90 days before harvest. Broader spectrum. Do not make more than two applications per year.
Atrazine 4L Herbicide	See tank mixture partner label for its rates. NIS or COC Apply broadcast spray on sugarcane. Apply when broadleaf weeds are small (1.5 inches or less). Post-emergence control of labeled broadleaf weeds. Aids in the burndown and control of many grass weeds which have escaped pre-emergence herbicide treatments. Atrazine mixtures may result in reduced control (antagonism) of larger broadleaf weeds. Smaller weeds are easier to control.
Ametryn Herbicide	See tank mixture partner label for its rates. NIS Apply broadcast spray on sugarcane before crop emergence or post-emergence until row closure. Broadleaf weeds and grasses. Reduced efficacy occurs if temperatures exceed 85°F during application.
Glyphosate (various formulations)	See tank mixture partner label for its rates. NIS Apply as broadcast spray. For pre-plant burndown of emerged annual grasses, broadleaf weeds and nutsedge.

Crop	Rate Oz./Acre	PHI	RESTRICTIONS
Sweet Corn and Popcorn	¾	30	The maximum application rate is ¾ oz. of this product. Two applications of this product are allowed per year. Do not exceed with a total application of 1 ½ oz. of product (0.062 lb. active ingredient) per acre per use season. After application to foliage, allow 30 days before grazing domestic livestock, harvesting forage, or harvesting silage. Do not use on "Jubilee" sweet corn. Do not apply this product to sweet corn or popcorn previously treated with soil applied organophosphate insecticides. Do not apply an organophosphate insecticide within 7 days before or 3 days after any application of this product. Do not apply this product to sweet corn or popcorn unless the seed company, processor or State Agricultural Extension service has tested this product on the particular hybrid or variety and specifically approves and supports the use. Do not apply this product if the sweet corn and popcorn is under severe stress due to drought, water-saturated soils, low fertility (especially low nitrogen levels) or other poor growing conditions.

(continued)



Spray Applications

If used alone, apply a broadcast spray over-the-top or with drop nozzles from the spike through lay-by stage of corn. Use mechanical cultivation to control weeds species not on this label. Apply a sequential treatment, if necessary, only with drop nozzles semi-directed or directed to avoid application into the corn plant whorl.

Precautions:

Not all sweet corn and popcorn hybrids or varieties have been tested for sensitivity to this product. To the extent consistent with applicable law, the user assumes all responsibility for such use. After application, avoid cultivation for at least 7 days.

Sweet Corn and Popcorn Weed Height Activity Table

Weed Activity	Control	Suppression
Rate of Product	$\frac{2}{3}$ oz.	$\frac{2}{3}$ oz.
Weed Height	Inches	Inches
Burcucumber		1 - 3
Cocklebur, common	1 - 9	
Fleabane, Philadelphia	1 - 3	
Kochia ¹	1 - 3	
Lambsquarter, common		1 - 2
Mallow, Venice	1 - 3	
Milkweed, common		3 - 5
Milkweed, honeyvine		1 - 3
Nutsedge: yellow ²	3 - 6	
purple	3 - 6	
Passionflower, maypop	1 - 3	
Pigweed, redroot	1 - 3	
Pokeweed, common	1 - 6	
Ragweed: common	1 - 9	
Giant	1 - 3	
Smartweed, Pennsylvania	1 - 2	
Sunflower, common	1 - 12	
Velvetleaf	1 - 9	

¹ See Pre-emergent and Post-emergent Weed Activity Tables.

² Heavy infestations of nutsedge require sequential applications. To prevent nutsedge from competing with the crop an earlier application is required.

CROP ROTATIONAL GUIDELINES

Following applications of this product, the crop rotational intervals listed below provide for adequate safety to newly planted crops. If the crop is planted in a shorter interval, crop injury may result. If the degradation of halosulfuron-methyl is slowed down by the conditions such as drought, cool conditions or drip irrigation in Arizona and California, the time lines need to be extended. Since all possible environmental and application scenarios, have not been tested, Atticus, LLC suggests that the end user test this product in order to determine its suitability for such intended use. In areas where local experience has demonstrated crop safety, use the shorter intervals. In the event of crop failure, labeled crops may be planted back into the treated area at the user's risk for potential phytotoxicity to the subsequent crop.

TIME INTERVAL (MONTHS) BEFORE PLANTING AFTER USE OF PROMOTE

CROP	MONTHS	EXCEPTIONS
CROP NOT SPECIFICALLY LISTED	36	
Alfalfa	9	
Barley (winter)	2	
Beans, Dry	9	In the northeast, southeast, TX and CO: 2 months.
Beans, Snap	9	In the northeast and southeast: 2 months; In TX: 3 months.
Broccoli	18	In muck soils areas of FL: 3 months.
Cabbage	15	In muck soils areas of FL: 3 months.
Canola	15	
Carrot	15	
Cauliflower	18	In muck soils areas of FL: 3 months.
Cereal crops, Spring	2	
Clovers	9	
Collards	18	
Corn, IR/IMR Field	0	
Corn, IT Field	1	
Corn, Normal Field	1	
Corn, Seed	2	
Corn, Sweet and Popcorn	3	For sweet corn and popcorn, the application rates of this product are specific to those crops. For re-planting sweet corn and popcorn crops in those treated areas, that are lost, terminated or harvested, the crop rotational interval must be adhered to.
Cotton	4	
Cucumbers	9	In the northeast and southeast: 2 months; In TX: 3 months.
Eggplant	12	For FL transplants: 4 months.
Forage Grasses	2	
Lettuce Crops	18	In muck soils areas of FL: 3 months.
Melons	9	In southeast and TX: 2 months.
Mint	15	
Oats	2	
Onions and Leeks	18	
Peanuts	6	
Peas	9	
Peas, Fields	9	
Peppers	10	For FL transplants: 4 months and for TX transplants: 3 months.

(continued)



CROP	MONTHS	EXCEPTIONS
Potatoes	9	
Pumpkins	9	In southeast: 2 months.
Proso Millet	2	
Radish	12	In muck soils areas of FL: 3 months.
Red Beet	24	If irrigation is required or rainfall is sparse, the time interval is 36 months.
Rice	2	
Rye (winter)	2	
Sorghums	2	
Soybeans	9	
Spinach	24	In muck soils areas of FL: 3 months.
Squash	9	In southeast: 2 months.
Strawberries	36	For annual FL transplants: 6 months.
Sugar beet	24	If irrigation is required or rainfall is sparse, the time interval is 36 months; In MI: 21 months; In MN, ND, Red River Valley: 36 months.
Sugarcane	0	
Sunflowers	18	
Tomato (transplant)	8	In the northeast and southeast: 2 months; In TX: 3 months.
Wheat (winter)	2	

When used with tank mixture partners, consult the partner product labels to determine rotational crop restrictions. Follow the most restrictive label when planning and applying the tank mixture combinations.

Southeast: AL, FL, GA, LA, MS, NC, Puerto Rico, SC, TN.

Northeast: CT, DE, IA, IL, IN, KY, MA, MD, ME, MI, MN, MO, ND, NE, NH, NJ, NY, OH, PA, RI, SD, VA, VT, WI, WV.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a dry and secure location.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

PLASTIC BOTTLES

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Once triple rinsed, recycle if available. Some agricultural pesticide containers can be taken to a container collection site or pick up for recycling. To find the nearest site, contact your chemical dealer or manufacturer. If recycling is not available, dispose of in a sanitary landfill or by incineration if allowed by state and local ordinances.

Conditions of Sale and Limitation of Warranty and Liability

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials, resistant strains or other influencing factors in the use of the product, which are beyond the control of Atticus, LLC or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Atticus, LLC and Seller harmless for any claims relating to such factors.

To the extent allowed by applicable laws, Atticus, LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of the product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or Atticus, LLC and Buyer and User assume the risk of any such use. TO THE EXTENT ALLOWABLE BY APPLICABLE LAW, ATTICUS, LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent allowed by applicable laws, in no event shall Atticus, LLC or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT ALLOWABLE BY APPLICABLE LAW THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF ATTICUS, LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF ATTICUS, LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.

Atticus, LLC and Seller offer this product, and Buyer and User accept it, subject to foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by a duly authorized representative of Atticus, LLC.

Promote™ is a trademark of Atticus, LLC.

Permit® is a registered trademarks of Gowen Company, LLC.

All other trademarks are the property of their respective owners.

F20180514a

