



**GUARANTEED ANALYSIS**

Boron (B) ..... 10.0%

Derived From: Boric Acid

**WARNING:** This product contains boron (B) and may be harmful to certain crops.

Use only according to manufacturer's directions.

**ALSO CONTAINS NON-PLANT FOOD INGREDIENTS**

01.00% ..... Organic Matter (derived from leonardite)

00.15% ..... Carbohydrates (derived from gluconic acid)

98.85% ..... Inert Ingredients

**FOR AGRICULTURAL AND PROFESSIONAL USE ONLY**

**KEEP OUT OF REACH OF CHILDREN**



**WARNING**

Avoid breathing mists. Harmful if swallowed. Avoid contact with skin, eyes or clothing.

**FIRST AID**

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. **If on skin:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice. **If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person. **If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

**FOR A MEDICAL EMERGENCY INVOLVING THIS PRODUCT, CALL 1-866-944-8565.**

**DIRECTIONS FOR USE**

Uses include but are not limited to the research verified crop specific directions in the table below.

CROP	APPLICATION RATE PER ACRE	APPLICATION DIRECTIONS AND TIMING
(01) Root and Tuber Vegetables including Carrot, Ginseng, Horseradish, Parsley (turnip-rooted), Potato, Radish, Sugar Beet, Sweet Potato, Turnip	<b>Soil:</b> 0.5 to 4.0 quarts <b>Foliar:</b> 1.0 to 2.0 pints	
(03) Bulb Vegetables including Garlic, Leek, Onion	<b>Soil:</b> 0.5 to 4.0 quarts <b>Foliar:</b> 1.0 to 2.0 pints	
(04) Leafy Vegetables including Celery, Head and Leaf Lettuce and Spinach	<b>Soil:</b> 0.5 to 4.0 quarts <b>Foliar:</b> 1.0 to 2.0 pints	<b>Caution:</b> Foliar applications close to harvest may leave a brown residue on the harvestable portion of the crop. Rainfall, sprinkler irrigation or high humidity should alleviate this condition.
(05) Brassica Vegetables including Broccoli, Cabbage, Cauliflower, Kale, and Mustard greens	<b>Soil:</b> 0.5 to 4.0 quarts <b>Foliar:</b> 1.0 to 2.0 pints	<b>Caution:</b> Foliar applications close to harvest may leave a brown residue on the harvestable portion of the crop. Rainfall, sprinkler irrigation or high humidity should alleviate this condition.
(06) Legume Vegetables (Succulent or Dried) including Bean (all types), Peas and Soybeans**	<b>Soil:</b> 0.5 to 4.0 quarts <b>Foliar:</b> 1.0 to 2.0 pints	
(08) Fruiting Vegetables including Eggplant, Pepper and Tomato	<b>Soil:</b> 0.5 to 4.0 quarts <b>Foliar:</b> 1.0 to 2.0 pints	<b>Caution:</b> Foliar applications close to harvest may leave a brown residue on the harvestable portion of the crop. Rainfall, sprinkler irrigation or high humidity should alleviate this condition.
(09) Cucurbit Vegetables including Cantaloupe, Cucumber, Honeydew, Muskmelon, Squash (summer and winter) and Watermelon	<b>Soil:</b> 0.5 to 4.0 quarts <b>Foliar:</b> 1.0 to 2.0 pints	<b>Caution:</b> Foliar applications close to harvest may leave a brown residue on the harvestable portion of the crop. Rainfall, sprinkler irrigation or high humidity should alleviate this condition.
(10) Citrus Fruit including Grapefruit, Lemon, Lime, Sweet Orange and Tangerine	<b>Soil:</b> 0.5 to 4.0 quarts <b>Foliar:</b> 1.0 to 2.0 pints	
(11) Pome Fruits including Apple and Pear	<b>Soil:</b> 0.5 to 4.0 quarts <b>Foliar:</b> 1.0 to 2.0 pints	
(12) Stone Fruits including Apricot, Cherry, Peach and Plumcot	<b>Soil:</b> 0.5 to 4.0 quarts <b>Foliar:</b> 1.0 to 2.0 pints	
(13) Berry and Small Fruit including Blackberry, Blueberry, Caneberry, Kiwi, Raspberry and Grape	<b>Soil:</b> 0.5 to 4.0 quarts <b>Foliar:</b> 1.0 to 2.0 pints	
(14) Tree Nuts including Almonds, Cashews and Pecans	<b>Soil:</b> 0.5 to 4.0 quarts <b>Foliar:</b> 1.0 to 2.0 pints	
(15) Grain, Cereal	<b>Soil:</b> 0.5 to 4.0 quarts <b>Foliar:</b> 1.0 to 2.0 pints	
(16) Forage, Fodder and Straw of Cereal Grains	<b>Soil:</b> 0.5 to 4.0 quarts <b>Foliar:</b> 1.0 to 2.0 pints	
(17) Grass Forage, Fodder and Hay	<b>Soil:</b> 0.5 to 4.0 quarts <b>Foliar:</b> 1.0 to 2.0 pints	
(18) Nongrass Animal Feeds including Alfalfa, Clover, Hay and Vetch	<b>Soil:</b> 0.5 to 4.0 quarts <b>Foliar:</b> 1.0 to 2.0 pints	
(19) Herbs and Spices including Basil, Dill, Mustard and Sage	<b>Soil:</b> 0.5 to 4.0 quarts <b>Foliar:</b> 1.0 to 2.0 pints	
(20) Oil Seed Crops including Canola**, Flax and Sunflower	<b>Soil:</b> 0.5 to 4.0 quarts <b>Foliar:</b> 1.0 to 2.0 pints	
(22) Stalk, Stem and leaf petiole vegetables, Asparagus, bamboo shoots and celery	<b>Soil:</b> 0.5 to 4.0 quarts <b>Foliar:</b> 1.0 to 2.0 pints	
(99) Cotton, Peanut, Sugarcane, Tobacco**	<b>Soil:</b> 0.5 to 4.0 quarts <b>Foliar:</b> 1.0 to 2.0 pints	<b>Foliar:</b> Repeat every 2 weeks as needed.

CROP	APPLICATION RATE PER ACRE	APPLICATION DIRECTIONS AND TIMING
Ornamental and Landscape	<b>Soil:</b> 0.5 to 4.0 quarts <b>Foliar:</b> 1.0 to 2.0 pints	<b>Foliar:</b> Repeat every 2 weeks as needed.
Turf	<b>Soil:</b> 0.3 to 3 oz per 1000 square feet <b>Foliar:</b> 0.3 to 0.75 oz per 1000 square feet	

\*The underlined applications are considered the single most important application.

**Fertigation:**

Can be applied through Center Pivot, Drip and Micro Sprinkler Irrigation systems. Continue irrigation for a minimum of 2 hours after application to flush the lines and to maximize nutrient distribution with drip and micro sprinkler use. Inject after the irrigation water filter.

Mixing with water for field storage is generally not recommended. Products blended with water should be used immediately and not stored for prolonged periods.

**Aerial Application:**

In aerial application use adequate water to obtain thorough coverage.

**Caution:**

Foliar applications close to harvest may leave a brown residue on the harvestable portion of the crop. Rainfall, sprinkler irrigation or high humidity should alleviate this condition.

**Caution:**

Always check compatibility with other products prior to use.

Do NOT apply with conventional calcium fertilizers.

Fertilizer solutions with a pH of 4 or less may not be compatible.

Combining with liquid fertilizer solutions is more effective if done at a blending plant with vigorous agitation in blending process.

**STORAGE AND DISPOSAL**

**STORAGE:** Store in original containers only. Keep containers tightly closed when not in use. Store in a cool, dry well-ventilated area, preferably in a locked storage area away from children, feed and food products, and seed. Store at ambient temperatures. Do not contaminate water, food, or feed by storage or disposal. **DISPOSAL:** Do not reuse container. Completely empty container into application equipment. Thoroughly rinse container, adding rinse water to the spray tank.

For help with any spill, leak, fire or exposure involving this material, call day or night CHEMTREC 1-800-424-9300.

**CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY**


BEFORE BUYING OR USING THIS PRODUCT, read the Directions for Use and the following Conditions of Sale and Limitation of Warranty and Liability. By buying or using this product, the buyer or user accepts the following Conditions of Sale and Limitation of Warranty and Liability, which no employee or agent of LOVELAND PRODUCTS, INC. or the seller is authorized to vary. LOVELAND PRODUCTS, INC. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, when the product is used in accordance with such Directions for Use under normal conditions of use. LOVELAND PRODUCTS, INC. MAKES NO OTHER WARRANTY, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. THE BUYER'S OR USER'S EXCLUSIVE REMEDY FOR ANY INJURY, LOSS, OR DAMAGE RESULTING FROM THE HANDLING OR USE OF THIS PRODUCT SHALL BE LIMITED TO ONE OF THE FOLLOWING, AT THE ELECTION OF LOVELAND PRODUCTS, INC. OR THE SELLER: DIRECT DAMAGES NOT EXCEEDING THE PURCHASE PRICE OF THE PRODUCT OR REPLACEMENT OF THE PRODUCT. LOVELAND PRODUCTS, INC. AND THE SELLER SHALL NOT BE LIABLE TO THE BUYER OR USER OF THIS PRODUCT FOR ANY CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES, OR DAMAGES IN THE NATURE OF A PENALTY.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.regulatory-info-lpi.com/>

Tests have not been carried out under all possible use conditions and Seller cannot and does not represent or warrant that the goods are compatible with all other chemicals or under all use conditions.

Reax Boron is a registered trademark of Loveland Products, Inc.

F1548



Density: 11.4 lbs/gallon @ 68 °F

**Net Contents: 265 gallons (1,003.13 liters)**

**Net Weight: 3,021 lbs (1,370.30 kg)**

**Product Code: 1000947977**