

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY INSTRUCTIONS BEFORE OPENING OR USING

CLETHODIM

240EC SELECTIVE HERBICIDE

ACTIVE CONSTITUENT:

240 g/L CLETHODIM

SOLVENT:

594 g/L LIQUID HYDROCARBON

GROUP

A

HERBICIDE

For the control of certain grass weeds in beetroot, cabbage, canola, celery, chickpeas, cotton, faba beans, field peas, forestry, lentils, lettuce, lupins, mung beans, non-bearing fruit trees, onions, ornamentals, peanuts, potatoes and soybeans as per the Directions for Use Table

IMPORTANT: READ THIS LEAFLET BEFORE USE

RELYON

APVMA Approval No:
89898/126312

DIRECTIONS FOR USE:

Restrains: DO NOT apply without the addition of an adjuvant.
DO NOT apply to plants that are stressed by moisture or temperature extremes.
DO NOT apply if rain is expected within one hour of application
DO NOT apply Relyon Clethodim 240EC Selective Herbicide more than once to any one crop.

CROP	CROP GROWTH STAGE	WEEDS CONTROLLED	RATE (mL/ha)
Canola, Chickpeas, Faba beans, Field peas, Lentils ⁽¹⁾ , Lupins*	Canola: DO NOT APPLY after flower buds become visible (green buds) Chickpeas: DO NOT APPLY beyond full flowering Faba beans: DO NOT APPLY beyond full flowering Field Peas: DO NOT APPLY beyond full flowering Lentils: Apply up to the 7 node/early branching stage of crop growth. Lupins: DO NOT APPLY after 80% of flowers have opened	Annual ryegrass (<i>Lolium rigidum</i>) Annual phalaris (<i>Phalaris minor</i>)	150 to 500
		Barley grass (<i>Hordeum leporinum</i>), Brome grass (<i>Bromus diandrus</i>), Wild Oats (<i>Avena</i> spp.)	175 to 500
		Volunteer wheat (<i>Triticum aestivum</i>), Volunteer oats (<i>Avena sativa</i>)	200* to 500 *(175 to 500 in WA only)
		Volunteer barley (<i>Hordeum vulgare</i>)	500
		Silver grass (<i>Vulpia bromoides</i>) – suppression only	250 to 500
		Paradox grass (<i>Phalaris paradoxa</i>)	250 to 375 375 - 500
Cotton (Qld, NSW and NT only) Peanuts, Mung beans, Soybeans	Cotton: DO NOT APPLY after full flowering (mid bloom) Peanuts: DO NOT APPLY after the pod fill stage of crop development Mung beans: DO NOT APPLY after the first flower buds are visible. Soybeans: DO NOT APPLY after the first flower buds are visible.	Barnyard grass (<i>Echinochloa</i> spp.) Blown grass (<i>Agrostis aveacea</i>), Crow'sfoot grass (<i>Eleusine indica</i>) Feathertop Rhodes grass (<i>Chloris virgata</i>), Liverseed grass (<i>Urochloa panicoides</i>), Red sprangletop grass (<i>Leptochloa filiformis</i>), Seedling Johnson grass ¹ (<i>Sorghum halepense</i>), Summer grass (<i>Digitaria</i> spp), Volunteer sorghum (<i>Sorghum</i> spp).	250 to 375 375 - 500

STAGE OF WEED GROWTH	STATE	CRITICAL COMMENTS
2-leaf to fully tillered	Qld, NSW, ACT, Vic, Tas, SA, WA only	Always apply with D-C-Trate at 2 L/100 L or Hasten at 1 L/100 L or Kwickin at 1 L/100 L or Uptake at 500 mL/100 L spray volume. The lower doses will provide effective control if applied under ideal conditions to weeds that are smaller, actively growing and free from temperature or water stress. See COMPATIBILITY AND CAUTIONS in GENERAL INSTRUCTIONS for mixture recommendations with insecticides, fungicides and other herbicides. ⁽¹⁾ Application up to 7-node/early –branching crop growth stage only *Not Queensland
2 to 5 leaf stage 5 leaf to fully tillered	NSW, ACT, Vic, Tas only	
2 to 5 leaf stage 5 leaf to fully tillered	Qld, NSW, ACT only	Always apply with D-C-Trate at 2 L/100 L or Hasten at 1 L/100 L or Kwickin at 1 L/100 L or Uptake at 500 mL/100 L spray volume. The lower doses will provide effective control if applied under ideal conditions to weeds that are smaller, actively growing and free from temperature or water stress. See COMPATIBILITY AND CAUTIONS in GENERAL INSTRUCTIONS for mixture recommendations with insecticides, fungicides and other herbicides. ¹ Only Johnson Grass seedlings germinating from seed will be controlled. Seedlings that appear later as shoots from underground stems will not be controlled, and more than one application may be required to control emerging shoots.

CROP	WEEDS CONTROLLED	RATE (mL/ha)
Beetroot, Cabbage, Celery, Lettuce, Potatoes, Onions	Barnyard grass (<i>Echinochloa</i> spp.) Blown grass (<i>Agrostis aveacea</i>), Crowsfoot grass (<i>Eleusine indica</i>) Feathertop Rhodes grass (<i>Chloris virgata</i>), Liverseed grass (<i>Urochloa panicoides</i>), Paradoxa grass (<i>Phalaris paradoxa</i>), Red sprangletop grass (<i>Leptochloa filiformis</i>), Seedling Johnson grass ¹ (<i>Sorghum halepense</i>), Summer grass (<i>Digitaria</i> spp), Volunteer sorghum (<i>Sorghum</i> spp).	250 to 375 375 - 500
	Annual ryegrass (<i>Lolium rigidum</i>) Annual phalaris (<i>Phalaris minor</i>)	150 to 500
	Barley grass (<i>Hordeum leporinum</i>), Brome grass (<i>Bromus diandrus</i>), Wild Oats (<i>Avena</i> spp.)	175 to 500
	Volunteer wheat (<i>Triticum aestivum</i>), Volunteer oats (<i>Avena sativa</i>)	200* to 500 *(175 to 500 in WA only)
	Volunteer barley (<i>Hordeum vulgare</i>)	500
	Silver grass (<i>Vulpia bromoides</i>) – suppression only (not Qld, WA)	250 to 500
	Winter grass (<i>Poa annua</i>)	500

STAGE OF WEED GROWTH	STATE	CRITICAL COMMENTS
2 to 5 leaf stage 5 leaf to fully tillered	All States	<p>Always apply with D-C-Trate at 2 L/100 L or Hasten at 1 L/100 L or Kwicken at 1 L/100 L or Uptake at 500 mL/100 L spray volume.</p> <p>The lower doses will provide effective control if applied under ideal conditions to weeds that are smaller, actively growing and free from temperature or water stress.</p> <p>Use a spray volume of 150 L/ha when spraying dense grass populations.</p> <p>¹ Only Johnson Grass seedlings germinating from seed will be controlled. Seedlings that appear later as shoots from underground stems will not be controlled, and more than one application may be required to control emerging shoots.</p>
2-leaf to fully tillered	All States	<p>Always apply with D-C-Trate at 2 L/100 L or Hasten at 1 L/100 L or Kwicken at 1 L/100 L or Uptake at 500 mL/100 L spray volume.</p> <p>The lower doses will provide effective control if applied under ideal conditions to weeds that are smaller, actively growing and free from temperature or water stress.</p> <p>Use a spray volume of 150 L/ha when spraying dense grass populations.</p>

CROP	WEEDS CONTROLLED	RATE (mL/ha)
Forestry, non-bearing fruit trees and ornamentals	Annual ryegrass (<i>Lolium rigidum</i>) Annual phalaris (<i>Phalaris minor</i>) Barley grass (<i>Hordeum leporinum</i>), Barnyard grass (<i>Echinochloa</i> spp.), Blown grass (<i>Agrostis aveacea</i>), Brome grass (<i>Bromus diandrus</i>), Crowsfoot grass (<i>Eleusine indica</i>) Feathertop Rhodes grass (<i>Chloris virgata</i>), Liverseed grass (<i>Urochloa panicoides</i>), Paradoxa grass (<i>Phalaris paradoxa</i>), Red sprangletop grass (<i>Leptochloa filiformis</i>), Seedling Johnson grass ¹ (<i>Sorghum halepense</i>) Silver grass (<i>Vulpia bromoides</i>) – suppression only (not Qld, WA), Summer grass (<i>Digitaria</i> spp), Volunteer sorghum (<i>Sorghum</i> spp). Volunteer wheat (<i>Triticum aestivum</i>), Volunteer oats (<i>Avena sativa</i>) Volunteer barley (<i>Hordeum vulgare</i>) Winter grass (<i>Poa annua</i>)	500

STAGE OF WEED GROWTH	STATE	CRITICAL COMMENTS
2 leaf to fully tillered	All States	Apply after plants have recovered from transplant shock and are showing signs of active growth. Always apply with D-C-Trate at 2 L/100 L or Hasten at 1 L/100 L or Kwickin at 1 L/100 L or Uptake at 500 mL/100 L spray volume. Use a spray volume of 150 L/ha when spraying dense grass populations. Do NOT use on Gymnosperms (pines, conifers, etc) unless a prior test has been conducted to check safety on the relevant species. See COMPATIBILITY AND CAUTIONS in GENERAL INSTRUCTIONS for particular species. ¹ Only Johnson Grass seedlings germinating from seed will be controlled. Seedlings that appear later as shoots from underground stems will not be controlled, and more than one application may be required to control emerging shoots.

**NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL
UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

WITHHOLDING PERIODS:

HARVEST:

CANOLA, CHICKPEAS, COTTON, FABA BEANS,
FIELD PEAS, LENTILS, LUPINS, MUNG BEANS,
PEANUTS, SOYBEANS:

NOT REQUIRED WHEN USED AS DIRECTED

FORESTRY:

NOT REQUIRED WHEN USED AS DIRECTED

BEETROOT, CABBAGE:

DO NOT APPLY LATER THAN 7 DAYS BEFORE HARVEST.

ONIONS:

DO NOT APPLY LATER THAN 14 DAYS BEFORE HARVEST.

LETTUCE, POTATOES:

DO NOT APPLY LATER THAN 4 WEEKS BEFORE HARVEST.

CELERY:

DO NOT APPLY LATER THAN 9 WEEKS BEFORE HARVEST.

GRAZING:

CANOLA, CHICKPEAS, FABA BEANS, FIELD PEAS, LENTILS, LUPINS, MUNG BEANS, PEANUTS, SOYBEANS: DO
NOT GRAZE OR CUT FOR STOCK FEED FOR 21 DAYS AFTER APPLICATION.

COTTON: DO NOT GRAZE OR CUT COTTON FORAGE OR STUBBLE FOR STOCK FEED.

GENERAL INSTRUCTIONS

RESISTANT WEEDS WARNING

GROUP	A	HERBICIDE
--------------	----------	------------------

Relyon Clethodim 240EC Selective Herbicide is a member of the cyclohexanedione group of herbicides. Relyon Clethodim 240EC Selective Herbicide has the inhibitor of acetyl coA carboxylase mode of action. For weed resistance management Relyon Clethodim 240EC Selective Herbicide is a Group A Herbicide. Some naturally occurring weed biotypes resistant to Relyon Clethodim 240EC Selective herbicide and other Group A Herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Relyon Clethodim 240EC Selective Herbicide or other Group A Herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Nutrien Ag Solutions Limited accepts no liability for any losses that may result from the failure of Relyon Clethodim 240EC Selective Herbicide to control resistant weeds.

CLEANING SPRAY EQUIPMENT

Before using Relyon Clethodim 240EC Herbicide

Ensure that the recommended clean-out procedure for the previous product (particularly sulfonyleurea herbicides) sprayed with the equipment was done properly.

After using Relyon Clethodim 240EC Herbicide

Empty the tank and drain the whole system.

Thoroughly wash inside of the tank using a pressure hose, drain the tank and clean all filters in the tank, pump line and nozzles.

Use of a household detergent will aid in cleaning the equipment. Add detergent to the part- filled spray tank and thoroughly circulate through pumps, hoses and nozzles. Drain the system and thoroughly rinse twice with clean water.

Rinse water should be discharged onto a designated disposal area, or if this is unavailable, onto unused land away from desirable plants and water sources.

MIXING

To ensure even mixing, half-fill the spray tank with clean water and add the required amount of product. Add spray additive and agitate thoroughly, then add the remainder of the water. Agitate again before spraying commences.

APPLICATION

Ground Application

Relyon Clethodim 240EC Selective Herbicide should be applied with calibrated spray equipment producing a median droplet range of 200 to 300 microns VMD. Apply in a minimum of 50 litres of water per hectare. Use 150 L/ha when spraying dense populations.

Aerial Application

Relyon Clethodim 240EC Selective Herbicide can be applied through aircraft fitted with boom or Micronair equipment. A spray volume of 20 to 30 L/ha is recommended and equipment should be adjusted to deliver droplets in the range of 200 to 250 microns VMD.

Best results will be obtained when aerial applications are made in a light crosswind. Applications should not be

made during temperature inversions or in conditions of very low relative humidity. Care should be taken to avoid drift damage to adjoining grass crops.

COMPATIBILITY

Relyon Clethodim 240EC Selective Herbicide is compatible with the following grass herbicides: Quizalofop-p-ethyl (99.5 g/L), Fluazifop-p (212 g/L), and Haloxyfop (520 g/L) NOTE: in all situations where Relyon Clethodim 240EC Selective Herbicide is mixed with another grass herbicide, Relyon Clethodim 240EC Selective Herbicide should be used at the full label rate specified for target weed species.

Relyon Clethodim 240EC Selective Herbicide may also be applied in tank mixtures with one of the following products:

Alphacypermethrin 100g/L Insecticide and Omethoate 290 g/L .

Relyon Clethodim 240EC Selective Herbicide should not be applied in a tank mix with Diflufenican (500 g/L), Basagran* or Blazer.

Apply as separate applications.

For information on compatibility of products not listed, please refer to Nutrien Ag Solutions Limited.

CAUTION

Ornamentals and Trees – While Relyon Clethodim 240EC Selective Herbicide is generally selective to broadleaved plants (i.e. it is active against grasses), tests should always be made on a smaller number of plants not previously tested.

DO NOT use on Gymnosperms (pines, conifers, etc) unless a prior test has been conducted to check safety on the relevant species.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS:

DO NOT apply Relyon Clethodim 240EC Selective Herbicide if wind is likely to cause drift onto susceptible crops/plants, cropping lands or pastures. Relyon Clethodim 240EC Selective Herbicide should not be applied through misting equipment or any other method likely to cause excessive drift. Care should be taken to avoid damage to adjoining native grasses or grass crops.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate wetlands or watercourses with this product or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Protect from frost. This container can be recycled if it is clean, dry, free of visible residues and has the drumMUSTER logo visible.

Triple-rinse container for disposal. Dispose of rinsate by adding it to the spray tank. Do not dispose of undiluted chemical on site. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any drumMUSTER collection or similar container management program site. The cap should not be replaced, but may be taken separately.

If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

Refillable Containers: Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. When preparing spray and using the prepared spray wear cotton overalls buttoned to the neck and wrist, washable hat, elbow-length PVC gloves, face shield or goggles. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766. If swallowed, do NOT induce vomiting.

SAFETY DATA SHEET

For further information refer to the Safety Data Sheet (SDS), which can be obtained from the supplier.

CONDITIONS OF SALE

The use of this product is beyond the control of Nutrien Ag Solutions Limited. Any provisions or rights under the Australian Consumer Law which cannot be excluded by law are not intended to be excluded by these conditions of sale. Subject to the foregoing, all warranties, conditions, rights and remedies expressed or implied under common law, statute or otherwise, in relation to the sale, supply, storage, use or application of this product are excluded. Nutrien Ag Solutions Limited does not accept any liability (including consequential loss and/or negligence) for any loss or damage connected with the sale, supply, storage, use or application of this product except for liability which cannot be excluded by statute.

RELYON®

Nutrien Ag Solutions Limited

Level 5, Building A, 26 Talavera Road, Macquarie Park, NSW 2113.

Tel: (02) 9889 5400

Product Support Tel: 1800 44 88 92

relyonaustralia.com.au

®RELYON is a Registered Trademark of Ruralco Holdings Limited