

GROUP 4 HERBICIDE

RhonoX®

MCPA Low Volatile Ester Herbicide

Controls broadleaf weeds in small grains, flax, rangeland and pasture, turf and other non-crop areas.

ACTIVE INGREDIENT:

2-ethylhexyl ester of 2-methyl-4-chlorophenoxyacetic acid* 67.9%

OTHER INGREDIENTS:..... 32.1%

TOTAL:..... 100.0%

Contains Petroleum Distillates

*2-methyl-4-chlorophenoxyacetic acid equivalent 44.1% by weight or 3.7 pounds per gallon.

**KEEP OUT OF REACH OF CHILDREN
CAUTION / PRECAUCION**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300
For Medical Emergencies Only, Call (877) 325-1840

EPA Reg. No. 11685-21-71368

Manufactured for
Nufarm Inc.
11901 S. Austin Avenue
Alsip, IL 60803

 **Nufarm**
Grow a better tomorrow



Net Contents
2.5 Gal.
(9.46 L)
Nonrefillable Container

12938000

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION / PRECAUCION**

Harmful if swallowed, inhaled or absorbed through the skin. Avoid contact with eyes, skin or clothing. Avoid breathing vapors or spray mist. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are made of barrier laminate, butyl rubber, nitrile rubber or viton.

Applicators and other handlers must wear: long-sleeved shirt and long pants, chemical-resistant gloves such as barrier laminate, butyl rubber, nitrile rubber or viton and shoes plus socks. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Additional PPE requirements for mixers and loaders supporting aerial application to rangelands, pasture lands, or noncropland. These mixers/loaders also must wear: - a chemical-resistant apron, and - a NIOSH-approved respirator with a dust/ mist filter with MSHA/NIOSH approval number prefix TC-21C or any N2, R, P, or HE filter. See engineering controls for additional requirements.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d) (4-6)], the handler PPE (personal protective equipment) may be reduced or modified as specified in the WPS.

Engineering Controls: Pilots must use an enclosed cockpit that meets the requirements listed in the WPS for agricultural pesticides [40 CFR 170.240(d)(6)].

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

FIRST AID

IF ON SKIN OR CLOTHING

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15 to 20 minutes.
- Call a poison control center or doctor for treatment advice.

IF INHALED

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.
- Call a poison control center or doctor for further treatment advice.

IF SWALLOWED

- Immediately call a poison control center or doctor for treatment advice.
- Do not induce vomiting unless told to do so by the poison control center or doctor.
- Do not give any liquid to the person.
- Do not give anything by mouth to an unconscious person.

IF IN EYES

- Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.

NOTE TO PHYSICIAN

No specific antidote is available. All treatments should be based on observed signs and symptoms of distress in the patient. Overexposure to materials other than this product may have occurred. This product contains petroleum distillates. If large amounts (greater than 1 ml/kg body weight) of the product have been ingested, the stomach should be evacuated by gastric intubation with aid of cuffed endotracheal tube to prevent aspiration of petroleum distillates. After removal of stomach contents, wash stomach by instilling 30 to 50 grams of activated charcoal in 3 to 4 ounces of water through the stomach tube and again remove stomach contents. Avoid oily laxatives.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish, aquatic invertebrates, and aquatic plants. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Ground Water Contamination: Most cases of ground water contamination involving phenoxy herbicides such as MCPA have been associated with mixing/loading and disposal sites. Caution should be exercised when handling MCPA pesticides at such sites to prevent contamination of ground water supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent ground water contamination. Cleaning of Equipment: Do not use the same spray equipment for other purposes unless thoroughly cleaned. When cleaning equipment, do not pour wash water on the ground; spray or drain over a large area away from wells and other water sources.

DIRECTIONS FOR USE

It Is A Violation Of Federal Law To Use This Product In A Manner Inconsistent With Its Labeling. Read entire label before using this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard (WPS), 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the WPS.

Do not enter or allow entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the WPS and that involves contact with anything that has been treated such as plants, soil or water is: coveralls, chemical-resistant gloves such as barrier laminate, butyl rubber, nitrile rubber or viton and shoes plus socks.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter or allow people (or pets) to enter the treated area until sprays have dried.

SPRAY DRIFT MANAGEMENT

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment-and-weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

Apply only as a medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

Apply only when the wind speed is 2 to 10 mph at the application site.

Additional requirements for aerial applications:

The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter. Release spray at the lowest height consistent with efficacy and flight safety. Do not release spray at a height greater than 10 feet above crop canopy. When applications are made with a crosswind, the swath will be displaced downwind. The applicator must compensate for this displacement at the downwind edge of the application area by adjusting the path of the aircraft upwind. Do not make applications into temperature inversions.

Additional requirements for ground boom applications:

Do not apply with a nozzle height greater than 4 feet above the crop canopy.

RESTRICTIONS

Do not apply this product through any type of irrigation system.

Wheat, barley, oat, rye and flax treated with MCPA may be replanted with any crop specified on an MCPA label or any crop for which a residue tolerance exists for MCPA. For crops not listed on an MCPA label, or on crops for which no residue tolerances for MCPA have been established, a 60 day plantback interval must be observed.

COMPATIBILITY

If this product is to be tank mixed with other pesticides, compatibility should be tested prior to mixing. To test for compatibility, use a small container and mix a small amount (0.5 to 1 quart) of spray, combining all ingredients in the same ratio as the anticipated use. If any indications of physical incompatibility develop, do not use this mixture for spraying. Indications of incompatibility usually will appear within 5 to 15 minutes after mixing. Read and follow all directions and precautions on this label and on the labels of any product(s) for which a tank mixture is being considered.

USE IN LIQUID NITROGEN FERTILIZER

This product may be combined with liquid nitrogen fertilizer suitable for foliage application on grass, pastures, or small grains in one operation. Use product according to directions on this label for those crops. Use liquid nitrogen fertilizer at rates recommended by supplier or Extension Service Specialist. Mix the product and fertilizer according to the following instructions: Fill the spray tank approximately half full with the liquid nitrogen fertilizer. Add the product while agitating the tank. Add the remainder of the fertilizer while continuing to agitate. Apply immediately, maintaining agitation during application until tank is empty. **Do not apply during cold (near freezing) weather.** Spray mixture must be used immediately and may not be stored. Do not allow mixture to stand overnight.

NOTE: If good, continuous agitation is not maintained, separation of the spray mixture and/or clogging of the nozzles is likely to occur. Fertilizers can increase foliage contact burn of herbicides. Reducing the fertilizer rate and concentration will reduce the hazard of leaf burn.

PRODUCT INFORMATION

This product is effective on a wide range of broadleaf weeds and is useful for controlling these weeds in certain crops. Several crops, such as flax, oats, and small grains are more tolerant of MCPA than they are of 2,4-D.

This product is an isooctyl ester formulation of MCPA that mixes easily in water, even in the hard waters found in some areas. Mix thoroughly while diluting this material and mix again if allowed to stand long before spraying.

WEEDS CONTROLLED BY PROPERLY APPLIED TREATMENTS OF THIS HERBICIDE INCLUDE THE FOLLOWING:

<i>Amaranthus</i> species (pigweeds)	dandelion field pennycress	pepperweeds (except perennial)	ragweed (common and giant)
common burdock	mustard (except blue mustard)	plantains	shepherdspurse
common cocklebur		purslanes	thistle (bull and Russian)
common lambsquarters			yellow rocket

WEED RESISTANCE

Any weed population may contain plants that are naturally resistant to MCPA, the active ingredient in this product, and to other herbicides with the same mode of action. **ATTENTION:** These resistant weed biotypes will not be controlled by this product. Consult advisors such as your local agricultural extension service for agronomic management practices to minimize the occurrence of MCPA resistance and considerations for supplemental control measures.

Weed Management

To minimize the occurrence of MCPA resistant biotypes, observe the following general weed management practices:

- Scout application site before and after herbicide applications.
- Start with a clean application site, using either a burndown herbicide application or tillage.
- Control weeds early when they are relatively small.
- Add other herbicides (e.g. a selective and/or a residual herbicide) and cultural practices (e.g. tillage or crop rotation) where appropriate.
- Utilize the specified label rate for the most difficult to control weed in your field. Avoid tank mixtures with other herbicides that reduce this product's efficacy (through antagonism), or tank mixture directions that encourage application rates of this product below the label directions.
- Control weed escapes and prevent weeds from setting seeds.
- Clean equipment before moving from field to field to minimize the spread of weed seed or plant parts.
- Report any incidence of repeated non-performance of this product on a particular weed to your Nufarm representative, local retailer, or county extension agent.

Management of MCPA Resistant Biotypes

Since the occurrence of new MCPA weeds cannot be determined until after product use and scientific confirmation, manufacturer is not responsible for any losses that may result from the failure of this product to control MCPA resistant weed biotypes.

The following good agronomic practices are recommended to reduce the spread of confirmed MCPA resistant biotypes:

- If a naturally occurring resistant biotype is present in your application site, this product should be tank-mixed or applied sequentially with an appropriately labeled herbicide with a different mode of action to achieve control.
- Cultural and mechanical control practices (e.g. crop rotation or tillage) may also be used as appropriate.
- Scout treated application site after herbicide applications and control escaping weeds including resistant biotypes before they set seed.
- Thoroughly clean equipment before leaving fields known to contain resistant biotypes.

DIRECTIONS FOR APPLYING RHONOX® HERBICIDE CEREAL GRAINS

CROP	AMOUNT OF RHONOX PER ACRE	DIRECTIONS
WHEAT, BARLEY, OATS, and RYE (not underseeded with legumes):		
Annual weeds	1/2 pint	Apply after grain is in the 3 to 4 leaf stage, but not forming joints in the stem. Do not spray in the boot to dough stage.
Annual and Biennial weeds	1 to 1-1/2 pints*	Apply after grain is fully tillered (usually about 4 to 8 inches high, but not forming joints in the stem). Do not apply in the boot to dough stage.
Perennial broadleaf weeds	1-1/2 pints*	

*Use the lower rate if small annual and biennial weeds are the major problem. Use the higher rate if perennial weeds or annual and biennial weeds are present which are in the hard-to-kill categories as determined by local experience. The higher rates increase the risk of grain injury and should be used only where the weed control problem justifies the risk of grain damage.

RESTRICTIONS AND LIMITATIONS FOR USE ON SMALL GRAINS

Use a minimum of 10 gallons of water per acre for ground application and 2 gallons of water per acre for aerial application.

Do not allow livestock animals to forage or graze treated areas within 7 days of slaughter.

Do not apply more than 1.6 pints of this product (0.75 lbs. a.e.) per acre per year.

FLAX

CROP	AMOUNT OF RHONOX PER ACRE	DIRECTIONS
FLAX	1/4 to 1/2 pint	Apply when flax is 2 to 8 inches tall. Do not spray once flax has reached the bud stage.

RESTRICTIONS AND LIMITATIONS FOR USE ON FLAX

Do not allow livestock animals to forage or graze treated areas within 7 days of slaughter.

Use a minimum of 8 gallons per acre for ground application and 3 gallons per acre for aerial application.

Do not apply more than 0.5 pint of this product (0.25 lb ae) per acre per year.

ESTABLISHED GRASS PASTURES AND RANGELAND

CROP	AMOUNT OF RHONOX PER ACRE	DIRECTIONS
Annual broadleaf weeds	1 to 2 pints	Apply when weeds are small and actively growing and prior to bud stage. Do not apply to newly seeded areas until grass is well established. Do not apply when grass is in the boot to milk stage if grass seed production is desired. Bentgrass and legumes may be injured by this treatment.
Perennial and biennial broadleaf weeds	2 to 3 pints	

RESTRICTIONS AND LIMITATIONS FOR USE ON PASTURES AND RANGELAND

Do not apply more than 3.2 pints of this product (1.5 lb ae) per acre per year.

Do not apply more than 2 applications per year with a minimum retreatment interval of 21 days.

For ground application, use a minimum of 10 gallons of water per acre.

For aerial application, use a minimum of 2 gallons of water per acre.

Do not graze animals on treated areas within 7 days after treatment.

GRASSES GROWN FOR SEED PRODUCTION

CROP	AMOUNT OF RHONOX PER ACRE	DIRECTIONS
Annual and perennial broadleaf weeds	1 to 2 pints	Apply to established stands in spring from tiller to early boot stage. Do not spray in boot stage. New spring seedings may be treated with the lower rate after grass seedlings have at least 5 leaves. Perennial weed regrowth may be treated in the fall.

RESTRICTIONS AND LIMITATIONS FOR USE ON GRASSES GROWN FOR SEED

For ground application use a minimum of 10 gallons of water per acre.

For aerial application use a minimum of 2 gallons of water per acre.

Do not apply more than 3.2 pints of this product (1.5 lb ae) per acre per year.

Do not apply more than 2 applications per year with a minimum retreatment interval of 21 days.

NON-CROP

(Roadsides, fencerows, vacant lots, rights-of-way and other similar areas)

CROP	AMOUNT OF RHONOX PER ACRE	DIRECTIONS
Annual, biennial and perennial broadleaf weeds	3.2 pints	Apply when annual broadleaf weeds are young and actively growing and perennial and biennial broadleaf weeds are actively growing and near the bud stage but before flowering. A second application may be required for best results on perennial and biennial weeds.

RESTRICTIONS AND LIMITATIONS FOR USE IN NON-CROP AREAS

For ground application use a minimum of 10 gallons of water per acre.

For aerial application use a minimum of 2 gallons of water per acre.

Do not apply more than 2 applications per year with a minimum retreatment interval of 21 days.

For Broadcast Application - Do not apply more than 3.2 pints of this product (1.5 lbs ae) per acre per year.

Do not apply more than 2 applications per year with a minimum retreatment interval of 21 days.

For Spot Treatment - Do not apply more than 6.4 pints of this product (3.0 lbs ae) per acre per year.

ORNAMENTAL TURF AREAS

(Golf Courses, Cemeteries and similar established grass areas)

CROP	AMOUNT OF RHONOX PER ACRE	DIRECTIONS
Annual broadleaf weeds	1 to 3 pints	Apply when annual broadleaf weeds are growing vigorously. Spring or fall applications will give the best results. Do not mow within 2 days before or after application. Do not use on lawns of creeping grasses such as bentgrass nor on freshly seeded turf until the grass has become well established.

RESTRICTIONS AND LIMITATIONS FOR USE IN TURF AREAS

Do not apply more than 6.4 pints of this product (3.0 lbs ae) per acre per year.

Do not apply more than 2 applications per year with a minimum retreatment interval of 21 days.

For ground application on turf areas use enough water for good coverage (minimum of 10 gallons per acre).

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container in a dry secured storage area. Keep container tightly closed when not in use.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law and may contaminate ground water. If these wastes can not be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA regional office for guidance.

CONTAINER DISPOSAL [HANDLING]:

NOTE: This product is available in multiple containers. Refer to the Net Contents section of this products labeling for the applicable "Nonrefillable" or "Refillable" designation. Follow the container disposal [handling] instructions below that apply to your container type/size.

Nonrefillable Containers 5 Gallons or Less: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

Nonrefillable containers larger than 5 gallons: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. If recycling or reconditioning not available, puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refillable containers larger than 5 gallons: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

OR

Refillable Container: Refill this container with pesticide only. Do not reuse this container for any other purpose. Close all openings and replace all caps. Contact Nufarm's Customer Service Department at 1-800-345-3330 to arrange for return of the empty refillable container.

Rhonox Crop Injury Warning

INJURY TO CROPS MAY OCCUR FROM THIS PESTICIDE. IF YOU ARE NOT PREPARED TO ACCEPT SOME DEGREE OF CROP INJURY, DO NOT USE THIS PRODUCT.

Crop varieties vary in response to MCPA, and some may be easily injured. Apply this herbicide only to varieties known to be tolerant to MCPA. If you are uncertain concerning tolerant varieties or local use situations that may affect crop tolerance to MCPA, contact your seed company or state Agricultural Extension Service for advice. If no information is available, limit the initial use to a small area or accept the potential for crop injury.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER, AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, CROP OR PLANT DAMAGE, OR LOSS OF YIELD, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE WEATHER, WIND, AND TEMPERATURE, OR THE MANNER OF USE OR APPLICATION, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER, AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER OR MANUFACTURER, AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY OTHER REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS. NO WARRANTIES SHALL BE CREATED BY COURSE OF DEALING, USAGE OF TRADE, OR COURSE OF PERFORMANCE. THERE ARE NO WARRANTIES THAT EXTEND BEYOND THE FACE HEREOF. THE SELLER OR MANUFACTURER ASSUMES NO RESPONSIBILITY THAT THE GOODS WILL BE FIT FOR ANY PARTICULAR PURPOSE FOR WHICH YOU MAY BE BUYING OR USING THE GOODS, EXCEPT AS OTHERWISE PROVIDED IN THE CONTRACT.

LIMITATION OF LIABILITY WARNING

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, NEITHER MANUFACTURER NOR SELLER SHALL BE LIABLE TO BUYER OR USER OR TO CUSTOMERS OF BUYER, IF ANY, FOR INDEMNIFICATION OR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE USE, MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY, SHALL BE FOR DAMAGES NOT EXCEEDING THE PURCHASE PRICE OF THE GOODS AND, IF BUYER OR USER WISHES, THE RETURN OF THE GOODS BY BUYER TO SELLER.

If you do not agree with or do not accept any of the directions for use, the warranty disclaimers, or limitations on liability, do not use the goods, and return it unopened to the Seller, and the purchase price will be refunded. *By using the goods, you expressly agree to all of the terms and conditions of this contract.*

USES WITH OTHER PRODUCTS (TANK MIXES) If the goods are used in combination with any other product except as specifically recommended in writing by Nufarm, then to the extent consistent with applicable law, Nufarm shall have no liability for any crop, plant, or other loss, damage, or injury arising out of its use in any such combination not so specifically recommended.

RV121114 [3]

Rhonox® is a registered trademark of Nufarm UK Limited.