



# 11-8-6

## Nutrient Spray

### GUARANTEED ANALYSIS

Total Nitrogen (N).....	11.00%
11.00% Urea Nitrogen	
Available Phosphorus (P <sub>2</sub> O <sub>5</sub> ).....	8.00%
Soluble Potash (K <sub>2</sub> O).....	6.00%
Boron (B).....	0.099%
Cobalt (Co).....	0.0005%
Copper (Cu).....	0.05%
0.05% Chelated Copper	
Iron (Fe).....	0.10%
0.10% Chelated Iron	
Manganese (Mn).....	0.05%
0.05% Chelated Manganese	
Molybdenum (Mo).....	0.0009%
Zinc (Zn).....	0.05%
0.05% Chelated Zinc	

DERIVED FROM: Urea, White Phosphoric Acid, Potassium Phosphate, Iron EDTA, Zinc EDTA, Copper EDTA, Manganese EDTA, Ammonium Molybdate, Cobalt Sulfate, and Boric Acid.

**WARNING:** This fertilizer contains boron. Its use on any other plants or under any other conditions other than those specified on the label may result in plant injury.  
**CAUTION:** This fertilizer is to be used only on crops which respond to molybdenum. Crops high in molybdenum are toxic to grazing animals (ruminants).

## KEEP OUT OF REACH OF CHILDREN PRECAUTION

Do not ingest. Avoid contact with skin, eyes, or clothing.  
 Avoid breathing dust, vapor or mist.

### FIRST AID

In all cases, call a poison control center or doctor for further treatment advice.

**IF SWALLOWED**, call a poison control center or doctor immediately. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything to an unconscious person. **IF ON SKIN**, take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. **IF INHALED**, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration. **IF IN EYES**, hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.

For chemical spills, leaks, fire or exposure, call CHEMTREC: (800) 424-9300.

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.

Store in a safe manner. Store in original container only and keep tightly sealed when not in use. Dispose of unused product and empty containers in accordance with federal, state and local regulations.

### GENERAL INFORMATION

**RNA NUTRIENT SPRAY 11-8-6** is a low salt, chlorine and sodium-free foliar fertilizer. **RNA NUTRIENT SPRAY 11-8-6** should be used in conjunction with a well balanced fertilizer program. To save on application cost, **RNA NUTRIENT SPRAY 11-8-6** may be applied with existing pesticide sprays. Rates listed under directions for use should be adjusted to fit weather, soil, and plant conditions.

Applications of **RNA NUTRIENT SPRAY 11-8-6** may begin as soon as the plant is large enough to absorb the foliar fertilizer. Applications of **RNA NUTRIENT SPRAY 11-8-6** may continue at 10 to 14-day intervals during the primary growth.

### DIRECTIONS FOR USE

- Do not apply **RNA NUTRIENT SPRAY 11-8-6** to water at a ratio of less than 1 to 10. (Example: 1 gallon of **RNA NUTRIENT SPRAY 11-8-6** to 10 gallons of water.)
- Avoid spray at full bloom. When spraying 10-50% blossoms, use lower rates.
- Do not use in high alkaline or acid solutions. Zinc EDTA, Manganese EDTA, Copper EDTA, Magnesium EDTA, Iron EDTA, and Calcium EDTA may be added to the recommendation if there is a specific need. **RNA NUTRIENT SPRAY 11-8-6** is especially beneficial for quick response in crops formerly under stress (i.e., disease, drought, hail, cold, water).

Crops	Rate Per Acre	Remarks
Fruit & Nut Crops	1-4 quarts	Apply with pre- and post-bloom sprays.
Cauliflower, Cabbage, Brussels Sprouts & other Brassica Crops	1-4 quarts	Apply at 7 to 10-day intervals using a minimum of 3 applications. A wetting will ensure good coverage. Consult your local PCS for recommendations on sensitive varieties.
Lettuce, Celery, Endive, and other Lettuce Crops	1-4 quarts	Begin application at 3 to 4-leaf stage. Continue applications at 7 to 10-day intervals through main growing season.
Wheat, Barley, and other Small Grains	1-4 quarts	Apply first application with broadleaf herbicide and the second application at root stage. A third application with fungicide may be required. Apply pre- and post-bloom.
Citrus	2-6 quarts	Apply pre- and post-bloom.

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.aapfco.org/metals.html>

Crops	Rate Per Acre	Remarks
Field, Row, and other Vegetable Crops	1-4 quarts	Apply after 2-leaf stage and before and during fruit growth.
Grapes	1-4 quarts	Apply with mildew spray and continue in 10-day intervals with pesticides and growth stimulants until bunch closing.
Hops	2-4 quarts	Apply with pesticides (2 applications before vines reach top of wire).
Peanuts	2-4 quarts	Apply early and continue applications with pesticide treatments.
Rice	2-4 quarts	Begin treatment before bloom and continue treatments (2) until head has formed.
Sugar Beets	2-4 quarts	Apply first application between 4 to 6-leaf stage. Continue applications at 14-day intervals.
Strawberries	2-4 quarts	Add 6 ounces per 10 gallons transplant water. Apply first application after transplanting and when growth begins. Continue up to harvest.
Tomatoes, Eggplant, and Potatoes	2-4 quarts	Add 6 ounces per 10 gallons transplant water. Apply first application with fungicide. Continue regular application until 3 weeks to harvest.
Lawn and Turf	2-4 quarts	Apply as growth begins and continue throughout growing season at 3-week intervals.

### COMPATIBILITY

Always jar test new combinations for compatibility prior to field mixing.

Conditions of Sale and Limitation of Warranty and Liability:

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of the product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of many different factors including, without limitation, manner of use or application, weather, combination with other products, or crop conditions. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Manufacturer and Seller harmless from any claims relating to such factors.

Seller warrants that this product conforms to the chemical description on the label. EXCEPT FOR THIS WARRANTY, THE PRODUCT IS FURNISHED "AS-IS," AND NEITHER SELLER NOR MANUFACTURER MAKES ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, WITH RESPECT TO THE SELECTION, PURCHASE OR USE OF THIS PRODUCT; SELLER AND MANUFACTURER SPECIFICALLY DISCLAIM ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. Buyer and User accept all risks arising from any use of this product, including without limitation uses contrary to label instructions, under abnormal conditions, or under conditions not reasonably foreseeable to (or beyond the control of) Seller or Manufacturer.

To the extent permitted by law, neither Manufacturer nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. THE EXCLUSIVE REMEDY OF THE BUYER OR USER, AND THE EXCLUSIVE LIABILITY OF MANUFACTURER AND SELLER, FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THIS PRODUCT, OR, AT THE ELECTION OF MANUFACTURER OR SELLER, THE REPLACEMENT OF THE PRODUCT.

These Conditions of Sale and Limitation of Warranty and Liability shall be interpreted in accordance with the laws of the State of California, excluding its conflicts of laws rules, and may not be amended by any oral or written agreement.

K-052717

Manufactured by: RNA • 22312 Railroad Avenue, San Joaquin, CA 93660

**NET CONTENTS:**   **GALLONS / Liter**  
**WEIGHT PER GALLON: 10.2 POUNDS at 68°F • pH 6.5 – 6.8**