

Roundup

FOR LAWNS,

**KILLS WEEDS ·
WON'T HARM LAWNS***

KILLS CRABGRASS · DANDELION · CLOVER · YELLOW NUTSEGE***

FAST ACTING FOR VISIBLE RESULTS IN HOURS

Keep Out Of Reach Of Children

CAUTION Stop! Read the entire label first. Observe all precautions and follow directions carefully. See back panel booklet for additional precautionary statements and first aid.

Mantener Fuera Del Alcance De Los Niños

PRECAUCIÓN ¡Atención! Lea primero toda la etiqueta. Observe todos los avisos de precaución y siga atentamente las instrucciones. También preste atención a los avisos de precaución y los primeros auxilios en la contraportada.

NET CONTENTS / CONTENIDO NETO: 1.25 GALLONS (1 GAL 32 FL OZ)

ACTIVE INGREDIENTS:
 MCPA, dimethylamine salt 0.257%
 Quinclorac 0.118%
 Dicamba, dimethylamine salt 0.029%
 Sulfentrazone 0.015%
OTHER INGREDIENTS: 98.581%
TOTAL 100.000%

THIS PRODUCT CONTAINS:
 0.018 lb 2-methyl-4-chlorophenoxyacetic acid equivalent per gallon or 0.210%
 0.010 lb 3,7-dichloro-8-quinolinecarboxylic acid per gallon or 0.118%
 0.002 lb 3,6-dichloro-o-anisic acid equivalent per gallon or 0.024%
 0.001 lb N-[2,4-dichloro-5-[4-(difluoromethyl)-4,5-dihydro-3-methyl-5-oxo-1H-1,2,4-triazol-1-yl]phenyl]methanesulfonamide per gallon or 0.015%

3 UP TO
MONTH
CONTROL**

PPN 1001101-04

2D
CODE

LB 5468

* When used as directed
 ** As listed
 *** See inside booklet for weed list

OPEN / ABRA

READY-TO-USE

FOR USE ON NORTHERN GRASSES

KILLS 253 TYPES OF WEEDS**

ONE STEP WEED CONTROL
FOR UP TO 3 MONTHS



For best results in non-lawn areas like garden beds, patios and driveways, use Roundup® Ready-To-Use Weed & Grass Killer III

† **GUARANTEED RESULTS:** If for any reason you are not satisfied after using this product, simply send us original proof of purchase and we will refund the purchase price.

📱 **Questions & Comments** Preguntas y comentarios
Call 1-800-246-7219 or Visit our Website at www.roundup.com

Produced for:
Monsanto Company
Lawn & Garden Products
800 N. Lindbergh Blvd.
St. Louis, MO 63167
© 2020 Monsanto Company

EPA Reg. No. 2217-917-71995
EPA Est. No. 239-IA-3

** As listed

PPN 1001101-05
2D Code
LBS18

12-digit UPC-A
(non-suppressed)
For Position Only
0 70183 50201 1

1005

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

RESTRICTIONS:



- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Do not allow people or pets to enter treated area until spray has dried.

PRECAUTIONS:

Roundup® For Lawns, is a ready-to-use (RTU) product intended for "spot" treatment of individual broadleaf and grass weeds by homeowners on residential lawns. This product may be applied to lawns containing the following species of grass: Kentucky bluegrass, perennial ryegrass, fescue species (including tall, red, and fine leaf varieties), bermudagrass¹, buffalograss and zoysiagrass.

¹ Application to bermudagrass may cause temporary yellowing or discoloration but full recovery can be expected. Do not apply to bermudagrass when daytime temperatures exceed 85° F.

Roundup® For Lawns, should not be used on lawns containing bentgrass, St Augustinegrass, bahiagrass, centipedegrass, or carpetgrass. Do not use on dichondra or lawns with desirable clovers or legumes, on vegetable gardens, fruits, or ornamentals (flowers, trees, groundcovers, hedges, landscape/flower beds and shrubs).

WEEDS CONTROLLED:

Roundup® For Lawns, will control or suppress the following broadleaf and grass weeds.

BROADLEAF WEEDS

Amaranth, purple	Bedstraw, northern	Black medic
Amaranth, slender	Bedstraw, pineywoods	Blood flower milkweed
Aster, heath	Beggarticks, hairy	Blue lettuce
Aster, white heath	Betony, Florida	Brassbuttons
Aster, purple	Bindweed, field	Bristly oxtongue
Aster, white prairie	Bindweed, hedge	Broomweed
Artichoke	Bitter Wintercress	Buckhorn
Austrian fieldcress	Bittercress, hairy	Bullnettle
Bedstraw, catchweed	Bitterweed	Burclover
Bedstraw, smooth	Black-eyed Susan	Burdock

Burweed, lawn (spurweed)	Cocklebur, spiny	Galinsoga, hairy
Buttercup, bulbous	Common mullein	Galinsoga, smallflower
Buttercup, creeping	Corn spurry	Goldenrod
Buttonweed, common (poorjoe)	Creeping beggarweed	Ground ivy
Buttonweed, Virginia	Creeping Jenny (creeping Charlie)	Groundsel
Carolina geranium (wild)	Cupid shaving brush	Gumweed
Carpetweed	Cudweed	Hairy fleabane
Catnip	Daisy, English	Hawkweed, meadow
Catsear, spotted (false dandelion)	Daisy, oxeye	Hawkweed, mouseear
Chamber bitter	Dandelion, common	Hawkweed, orange
Chickweed, common	Dandelion, false	Hawkweed, yellow
Chickweed, mouseear	Day flower	Heatall
Chickweed, sticky	Deadnettle	Heartleaf drymary
Chicory	Dichondra	Hemp
Cinquefoil	Dock, broadleaf	Henbit
Clover, crimson	Dock, curly	Hoary cress
Clover, alsike	Dogbane	Horsenettle
Clover, hop	Dogfennel	Horseweed
Clover, rabbit foot	Dollarweed (pennywort)	Jimsonweed
Clover, red	Doveweed	Johnny-Jumpup violet
Clover, strawberry	Elderberry	Knawel
Clover, yellow sweet	Falseflax	Knotweed, prostrate
Clover, white sweet	False sunflower	Kochia
Clover, white	Fiddleneck	Lambsquarters
Cockle	Filaree, redstem	Lespedeza
Cocklebur	Filaree, whitestem	Mallow, alkali
Cocklebur, oriental	Fleabane, daisy (annual)	Mallow, bristly
	Fleabane, rough	Mallow, common
		Mallow, dwarf
		Mallow, Venice
		Marcela

Matchweed	Pigweed, prostate	Ragweed, lanceleaf	Strawberry, Indian	Virginia creeper
Mexicanweed	Pigweed, redroot	Ragweed, western	mock	Western clematis
Moneywort	Pigweed, Russian	Redstem filaree	Tansy ragwort	Western salsify
Morningglory, bigroot	Pigweed, smooth	Rough cinquefoil	Tanweed	Wild aster
Morningglory, ivyleaf	Pigweed, tumble	Scarlet pimpernel	Thistle, blessed	Wild buckweed
Morningglory, Mexican	Pineappleweed	Shepherdspurse	Thistle, bull	Wild carrot
Morningglory, red	Plains coreopsis	Smartweed,	Thistle, Canada	Wild four-o'clock
Morningglory, tall	(tickseed)	ladysthumb	Thistle, floodman	Wild garlic
Morningglory, woolly	Plantain, blackseed	Smartweed, pale	(prairie)	Wild lettuce
Mugwort	Plantain, bracted	Smartweed,	Thistle, musk	Wild marigold
Mustard, ball	Plantain, broadleaf	Pennsylvania	Thistle, platte	Wild mustard
Mustard, black	Plantain, buckhorn	Smooth chaff-flower	Thistle, plumeless	(yellow rocket)
Mustard, blue	Plantain, hoary	Smooth dock	Thistle, Russian	Wild onion
Mustard, hedge	Plantain, narrowleaf	Sorrel, red (sheep)	Thistle, Scotch	Wild parsnip
Mustard, Indian,	Plantain, slender	Sowthistle, annual	Thistle, wavyleaf	Wild radish
Mustard, tansy	Plantain, woolly	Sowthistle, spiny	Thistle, yellow	Wild rape
Mustard, tumble	Poison ivy	Spanish needles	(horrible)	Wild strawberry
Mustard, white	Poison oak	Speedwell, bi-lobed	Trailing crownvetch	Wild sweet potato
Mustard, wormseed	Pokeweed	Speedwell, common	Velvetleaf	Wild violet
Nettle, stinging	Prairie sunflower	Speedwell, corn	Vervain, blue	Woodsorrel, creeping
(burning)	Prickly lettuce	Speedwell, ivyleaf	Vervain, hoary	Woodsorrel, violet
Nettle, tall	(compass plant)	Speedwell, Persian	Vervain, prostrate	Woodsorrel, yellow
Nettle, wood	Prickly sida	Speedwell, purslane	Vervain, tall	(Oxalis)
Parsley-piert	Prostrate knotweed	Speedwell, slender	Vetch, bird	Woolly croton
Parsnip	Puncturevine	Speedwell, snow	Vetch, hairy	Wormseed
Pearlwort	Purslane, common	Speedwell, thymeleaf	Vetch, milk	Yarrow, common
Pennycress	Pusley, Brazil	Spiny amaranth	Vetch, narrowleaf	Yarrow, western
(Frenchweed)	Pusley, Florida	Spurge, spotted	Vetch, wild	Yarrow, yellow
Pepperweed, perennial	Pusley, large flower	Spurge, prostrate	GRASSY WEEDS	
Pepperweed, Virginia	Ragweed, bur	Spurweed	Crabgrass	Foxtail (green, giant
(peppergrass)	Ragweed, common	Star of Bethlehem	(large and smooth)	and yellow) Yellow nutsedge

WHEN TO USE

Apply this product to actively-growing weeds during the early growth stages for the best results. Two or more applications at 14 day intervals throughout the season may be required for grass weeds. Apply when daytime temperatures are between 45° and 90° F.

SPRING AND EARLY SUMMER:

- **Crabgrass (large and smooth):** Apply this product from crabgrass emergence until the 6 leaf stage (2 tillers) or when the crabgrass is 3 to 4 inches tall. This period begins the first time that you can see crabgrass in your lawn. Use two applications with a 14-day interval between applications.
- **Foxtail (green, giant and yellow):** Apply this product to foxtail during the early growth stages or when the foxtail is 1 to 4 inches tall. Use two applications with a 14-day interval between applications.
- **Yellow nutsedge:** Two applications are required when the yellow nutsedge plants are 3 to 6 inches tall. Do not apply the second application until 14 days after the first application.
- **Dandelion, plantain, buttercup, speedwell, knotweed, pennywort (dollarweed) and other broadleaf weeds:** Apply when broadleaf weeds are young and actively growing with adequate soil moisture.

FALL:

- **Henbit, chickweed, white clover, ground ivy, lawn burweed, pearlwort, Carolina geranium and other broadleaf weeds:** Apply when broadleaf weeds are young and actively growing with adequate soil moisture for best results.

REFILL CONTAINER

Use this product to refill the empty Roundup® For Lawns, container by pouring product carefully and directly into the container. DO NOT add water.

HOW TO REFILL

Do not reuse or refill this container. Use this product to refill the empty Roundup® For Lawns, container by pouring product carefully and directly into the container. DO NOT add water.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE:	Store product in original container in a safe place away from direct sunlight. Keep from freezing. If frozen, allow to thaw and shake well before using.
PESTICIDE DISPOSAL AND CONTAINER HANDLING:	Nonrefillable container. Do not reuse or refill this container unless the directions for use allow a different concentrated or ready-to-use product to be diluted in or poured directly into the container. Reuse or refill this container according to the directions contained in the HOW TO REFILL section. If empty: Place in trash or offer for recycling, if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

FIRST AID	
IF IN EYES:	<ul style="list-style-type: none">• Hold eyes open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center for treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-246-7219 for emergency medical treatment advice.	

ENVIRONMENTAL HAZARDS

This product is toxic to fish, aquatic invertebrates, and aquatic plants. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift or runoff may be hazardous to aquatic organisms in water and to plants adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

NOTICE: To the extent consistent with applicable law, buyer assumes all responsibility for safety and use not in accordance with directions.

† GUARANTEED RESULTS: If for any reason you are not satisfied after using this product, simply send us original proof of purchase and we will refund the purchase price.

INSTRUCCIONES DE USO

Utilizar este producto de modo inconsistente con las declaraciones de su etiqueta constituye una violación de las leyes federales.

RESTRICCIONES:



- No aplique este producto de forma que entre en contacto con otras personas o mascotas, ya sea directamente o al ser arrastrado por el viento. No permita que las personas o las mascotas ingresen al área tratada hasta que se seque el rocío.

PRECAUCIONES:

Roundup® For Lawns, es un producto listo para usar (RTU) indicado para los propietarios de casas para el tratamiento “localizado” de grama y malezas de hoja ancha en céspedes. Este producto debe aplicarse a los céspedes con las siguientes especies de grama: azul de Kentucky, raigrás perenne, especies de cañuela o festuca (incluyendo las variedades alta, rubra y roja), Bermuda¹, búfalo y zoysia.

¹ La aplicación en el césped Bermuda puede causar amarillamiento o decoloración temporal, pero se puede esperar una recuperación completa. No aplique el producto a césped Bermuda cuando las temperaturas diurnas superen los 85° F.

Roundup® For Lawns, **no** debe utilizarse en céspedes de las variedades agrostis, San Agustín, bahía, ciempiés o alfombra. **No** lo utilice en dicondra ni en céspedes con tréboles o leguminosas que no desee eliminar, huertos de hortalizas, o plantas frutales u ornamentales (flores, árboles, cubiertas vegetales, setos, macizos de flores o jardines, y arbustos).

CUÁNDO USARLO

Para obtener los mejores resultados, aplique este producto a las malezas durante las primeras etapas de crecimiento activo. Pueda que se necesiten dos o más aplicaciones, a intervalos de 14 días, durante toda la temporada para eliminar las malezas gramíneas. Aplique cuando la temperatura diurna oscile entre los 45° y los 90° F.

EN LA PRIMAVERA Y A PRINCIPIOS DEL VERANO:

- **Para tratar el zacate mano de cangrejo (variedades lisa y sangradera ancha):** Aplique este producto a partir de la germinación del zacate mano de cangrejo hasta la etapa de 6 hojas (2 brotes) o cuando alcance una altura de 3 a 4 pulgadas. Este período comienza a partir del momento en el que observe zacate mano de cangrejo en el césped. Realice dos aplicaciones, dejando un intervalo de 14 días entre una y otra.
- **Para tratar el zacate cola de zorra (variedades verde, gigante y amarilla):** Aplique este producto al zacate cola de zorra durante las primeras etapas de crecimiento o cuando tenga entre 1 y 4 pulgadas de alto. Use dos aplicaciones, dejando un intervalo de 14 días entre una y otra.
- **Para tratar la juncia avellanada:** Es necesario hacer dos aplicaciones cuando las plantas de juncia avellanada tengan entre 3 y 6 pulgadas de alto. No realice la segunda aplicación hasta 14 días después de la primera.
- **Para tratar el diente de león, el lantén, la celidonia menor, la Verónica, el polígono, la centella asiática (acaricoba) y otras malezas de hoja ancha:** Aplique este producto cuando las malezas de hoja ancha sean pequeñas y estén creciendo de manera activa, con un suelo que tenga la humedad adecuada.

EN EL OTOÑO:

- **Para tratar la ortiga mansa, la estalaria, el trébol blanco, la hiedra rastretera, el codillo prieto del césped, la sagina, el geranio silvestre y otras malezas de hoja ancha:** A fin de obtener los mejores resultados, aplique este producto cuando las malezas de hoja ancha sean pequeñas y estén creciendo de manera activa, con un suelo que tenga la humedad adecuada.

RECARGA DE CONTENEDORES

Utilice este producto para rellenar el envase vacío del producto Roundup® For Lawns, vertiendo el producto con cuidado directamente en el contenedor. NO agregue agua.

CÓMO RELLENAR

No reutilice o rellene este envase. Utilice este producto para rellenar el envase vacío del producto Roundup® For Lawns, vertiendo el producto con cuidado directamente en el contenedor. NO agregue agua.

ALMACENAMIENTO Y ELIMINACIÓN

No contamine el agua, la comida ni el pienso para animales al almacenar o eliminar el producto.

ALMACENAMIENTO DEL PESTICIDA:	Guarde el producto en su envase original en un lugar seguro y alejado de la luz directa del sol. Proteja el producto del congelamiento. Si se congela, deje que se descongele y agítelo bien antes de usarlo.
ELIMINACIÓN DEL PESTICIDA Y MANEJO DEL ENVASE:	Este envase no se puede rellenar. No vuelva a usar o a rellenar este envase, a menos que las instrucciones de uso permitan que un producto concentrado o listo para usar distinto se pueda echar directamente o diluir en el envase. Este envase se puede reutilizar o rellenar de acuerdo con las instrucciones contenidas en la sección CÓMO RELLENAR. Si está vacío: Tírelo a la basura o reciclelo, de ser posible. Si está parcialmente lleno: Llame a su agencia local de desechos sólidos para obtener instrucciones sobre cómo eliminarlo. Nunca elimine el producto no utilizado en drenajes internos o externos.

AVISOS DE PRECAUCIÓN

RIESGOS PARA LOS SERES HUMANOS Y LOS ANIMALES DOMÉSTICOS

PRECAUCIÓN: Causa irritación moderada en los ojos. Evite el contacto con los ojos o la ropa. Lávese con abundante agua y jabón después de manipular el producto y antes de comer, beber, masticar chicle, consumir tabaco o usar el inodoro.

PRIMEROS AUXILIOS	
SI ENTRA EN LOS OJOS:	<ul style="list-style-type: none">Mantenga los ojos abiertos y enjuáguelos lentamente y con cuidado con agua por 15 a 20 minutos. Retire los lentes de contacto, de haberlos, después de transcurridos los primeros 5 minutos, y continúe enjuagando el ojo.Llame a un centro de control de envenenamientos para recibir instrucciones de tratamiento.
Tenga el envase o la etiqueta del producto con usted cuando llame al centro de envenenamientos o al médico, o al acudir a recibir tratamiento. También puede llamar al 1-800-246-7219 para obtener información sobre tratamiento médico de emergencia.	

RIESGOS MEDIOAMBIENTALES

Este producto es tóxico para los peces y los invertebrados acuáticos, así como para las plantas acuáticas. No lo aplique directamente al agua, en áreas donde haya agua presente en la superficie ni en zonas intermareales por debajo de la línea de la marea alta. El producto que se escurra o sea arrastrado por el viento puede ser peligroso para las plantas y para los organismos acuáticos de los cuerpos de agua adyacentes a las superficies tratadas. No contamine el agua al deshacerse de la que use para lavar o enjuagar los utensilios. Para proteger el medio ambiente, no permita que el pesticida penetre o se escurra en drenajes pluviales, zanjas de desagüe, cunetas o aguas superficiales. La aplicación de este producto en un clima tranquilo, cuando no se pronostiquen lluvias para las próximas 24 horas, ayudará a asegurar que ni el viento ni la lluvia arrastren o laven el pesticida fuera del área de tratamiento. Enjuagar el equipo usado para la aplicación sobre el área tratada ayudará a evitar escurrimientos hacia cuerpos de agua o sistemas de drenaje.

ADVERTENCIA: En la medida contemplada por las leyes correspondientes, el comprador asume toda la responsabilidad por la seguridad y el uso de este producto si no se siguen las instrucciones.

† **RESULTADOS GARANTIZADOS:** Si por algún motivo usted no queda satisfecho después de usar este producto, simplemente envíenos el comprobante de compra original y le reembolsaremos el precio de compra.



PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

FIRST AID	
IF IN EYES:	<ul style="list-style-type: none"> Hold eyes open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center for treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-246-7219 for emergency medical treatment advice.	

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE:	Store product in original container in a safe place away from direct sunlight. Keep from freezing. If frozen, allow to thaw and shake well before using.
PESTICIDE DISPOSAL AND CONTAINER HANDLING:	Nonrefillable container. Do not reuse or refill this container unless the directions for use allow a different concentrated or ready-to-use product to be diluted in or poured directly into the container. Reuse or refill this container according to the directions contained in the HOW TO REFILL section. If empty: Place in trash or offer for recycling, if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, aquatic invertebrates, and aquatic plants. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift or runoff may be hazardous to aquatic organisms in water and to plants adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.


Questions & Comments **Preguntas y comentarios**
 Call 1-800-246-7219 or Visit our Website at www.roundup.com

Produced for: EPA Reg. No. 2217-917-71995
 Monsanto Company EPA Est. No. 239-IA-3
 Lawn & Garden Products Roundup and Roundup & Design
 800 N. Lindbergh Blvd. are registered trademarks of
 St. Louis, MO 63167 Monsanto Technology LLC.
 © 2020 Monsanto Company



PPN 10011U-05 1005
 12-digit UPC-A
 (non-suppressed)
 For Position Only
 0 70183 150201 1

PRESS TO RESEAL. / PRESIONE PARA VOLVER A CERRAR.

