

RESTRICTED USE CLASSIFICATION

Due to Acute Fish Toxicity

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicators Certification.



Scourge®

Insecticide

with RESMETHRIN/PIPERONYL BUTOXIDE
18% + 54% MF FORMULA II

FOR USE ONLY BY FEDERAL, STATE, TRIBAL OR LOCAL GOVERNMENT OFFICIALS RESPONSIBLE FOR PUBLIC HEALTH AND VECTOR CONTROL, OR BY PERSONS CERTIFIED BY THE STATE OR TRIBAL LEAD PESTICIDE REGULATORY AGENCY TO PERFORM ADULT MOSQUITO CONTROL APPLICATION, OR BY PERSONS UNDER THEIR DIRECT SUPERVISION

- A SYNTHETIC PYRETHROID FOR EFFECTIVE ADULT MOSQUITO (INCLUDING ORGANOPHOSPHATE RESISTANT SPECIES), MIDGE (BITING AND NON-BITING), AND BLACKFLY CONTROL
- FOR CONTROL OF BITING AND NON-BITING FLIES
- FOR AERIAL OR GROUND APPLICATION
- CONTAINS 1.51 LB/GAL (181 g/L) OF RESMETHRIN AND 4.51 LB/GAL (540 g/L) OF PIPERONYL BUTOXIDE

ACTIVE INGREDIENTS

*Resmethrin	18.00%
**Piperonyl Butoxide Technical	54.00%
OTHER INGREDIENTS: ‡	28.00%
	100.00%

‡ Contains Petroleum Distillates

* Cis/trans isomers ratio max. 30% (±)cis and min. 70% (±)trans.

** Equivalent to min. 43.2% (butylcarbityl) (6-propylpiperonyl) ether and 10.8% related compounds.

EPA REG. NO. 432-667

EPA EST. NO. 432-TX-1

KEEP OUT OF REACH OF CHILDREN CAUTION

Refer to attached labels for complete First Aid, Precautionary Statements, Directions for Use and Storage and Disposal Instructions.

Net Contents

55 gal: 4192744

5 gal: 4192701

4494260D

100707AV1a

**BACKED
by BAYER™**

PRECAUCION AL CONSUMIDOR: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente. (TO THE USER: If you cannot read English, do not use this product until the label has been fully explained to you.)

FIRST AID	
IF SWALLOWED	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give any liquid to the person. • Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
[Synthetic Pyrethroid]	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
Note To Physician: Probable mucosal damage may contraindicate the use of gastric lavage.	

In case of Medical emergencies or health and safety inquiries or in case of fire, leaking or damaged containers, information may be obtained by calling 1-800-334-7577.

For Product information call Toll-Free: 1-800-331-2867.

PRECAUTIONARY STATEMENTS Hazards To Humans & Domestic Animals CAUTION

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Do not get on skin or clothing. Avoid contact with eyes. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

Environmental Hazards

This pesticide is extremely toxic to fish and aquatic invertebrates and shrimp. Runoff from treated areas or deposition of spray droplets into a body of water may be hazardous to fish and aquatic invertebrates. Do not apply over bodies of water (lakes, rivers, permanent streams, natural ponds, commercial fish ponds, swamps, marshes or estuaries), except when necessary to target areas where adult mosquitoes are present, and weather conditions will facilitate movement of applied material away from the water in order to minimize incidental deposition into the water body. Do not contaminate

water when disposing of equipment wash waters. This pesticide is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow drift when bees are actively visiting the treatment area, except when applications are made to prevent or control a threat to public and/or animal health determined by a state, tribal or local health or vector control agency on the basis of documented evidence of disease causing agents in vector mosquitoes, or the occurrence of mosquito-borne disease in animal or human populations, or if specifically approved by the state or tribe during a natural disaster recovery effort.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL FOR DIRECTIONS

This product is to be used for control of adult mosquitoes, (including organophosphate resistant species), midges (biting & non-biting) and blackflies by specially designed aircraft capable of applying ULTRA LOW VOLUME of finished spray formulation or by ground application with non-thermal or mechanical spray equipment that can deliver spray particles within the aerosol size range and at specified dosage levels.

NOTICE: This concentrate cannot be diluted in water. Mix well before using. Avoid storing excess diluted formulation in spray equipment tank beyond the period needed for application.

In treatment of corrals, feed lots, swine lots, and zoos, cover any exposed drinking water, drinking water fountains and animal feed before application.

Before making the first application in a season, it is advisable to consult with the state or tribal agency with primary responsibility for pesticide regulation to determine if permits or other regulatory requirements exist. Do not exceed 25 applications at 0.007 pounds of resmethrin per acre (not to exceed a total of 0.18 pounds of resmethrin per acre) in any given season.

More frequent applications may be made to prevent or control a threat to public and/or animal health determined by a state, tribal or local health or vector control agency on the basis of documented evidence of disease causing agents in vector mosquitoes or the occurrence of mosquito-borne disease in animal or human populations, or if specifically approved by the state or tribe during a natural disaster recovery effort.

GENERAL

Scourge is designed for application as an Ultra-Low Volume (ULV) aerosol to control adult mosquitoes and flies in residential, industrial, urban, recreational areas, and other areas where the labeled pests are a problem. Scourge provides effective control of adult mosquitoes, black flies, gnats, biting and non-biting midges, stable flies, horse flies, deer flies, sheep flies, horn flies, yellow flies and nuisance flying insects such as house flies in areas such as but not limited to industrial areas, urban areas, parks, campsites, woodlands, athletic fields, golf courses, playgrounds, recreational and overgrown waste areas, roadsides, and other public areas where adult mosquitoes and flies occur.

All types of applications should be conducted at temperatures of 50°F or higher. Apply when ground wind speeds are equal to or greater than 1 mph.

Scourge can be applied as undiluted or diluted with refined soybean oil, light mineral oil of 54 second viscosity or other suitable solvent or diluent.

Spray Droplet Size Determination

Ground Equipment:

Spray equipment must be adjusted so that the volume median diameter is less than 30 microns (Dv 0.5 < 30 µm) and that 90% of the spray is contained in droplets smaller than 50 microns (Dv 0.9 < 50µm). Directions from the equipment manufacturer or vendor, pesticide registrant or a test facility using a laser-based measurement instrument must be used to adjust equipment to produce acceptable droplet size spectra. Application equipment must be tested at least annually to confirm that pressure at the nozzle and nozzle flow rate(s) are properly calibrated.

Aerial Equipment:

Spray equipment must be adjusted so that the volume median diameter produced is less than 60 microns (Dv 0.5 < 60 µm) and that 90% of the spray is contained in droplets smaller than 100 microns (Dv 0.9 < 100 µm). The effects of flight speed and, for non-rotary nozzles, nozzle angle on the droplet size spectrum must be considered. Directions from the equipment manufacturer or vendor, pesticide registrant or a test facility using a wind tunnel and laser-based measurement instrument must be used to adjust equipment to produce acceptable droplet size spectra. Application equipment must be tested at least annually to confirm that pressure at the nozzle and nozzle flow rate(s) are properly calibrated.

GROUND APPLICATION

Vehicle-Mounted ULV Cold Aerosol Generators or Vehicle-Mounted Non-Thermal Aerosol (Cold Fog): Apply through non-thermal ULV application equipment and base acreage calculations on the equipment manufacturer's recommended swath width. Apply at a rate not to exceed 0.007 pounds of resmethrin per acre in any given 24 hour period. An optimum swath is created when Scourge is applied from a truck that is being driven perpendicular to the wind direction. Direct the spray head of equipment to ensure even distribution of the spray cloud throughout the area. For best results, apply when insects are most active and meteorological conditions are conducive to keeping the spray cloud in the air column close to the ground. An inversion of air temperatures and a light breeze is preferable. Application during the cooler hours of the night or early morning is recommended. Scourge may be applied through ULV Cold Aerosol Generators, or other equipment designed for non-thermal ULV aerosol applications. The desired application rate may be obtained under different conditions by altering the dilution rate of Scourge, the flow rate of the insecticide from the application equipment and the vehicle speed. Examples are given in the following Table.

imum altitude of 75 ft for rotary and 100 ft for fixed wing aircraft. Applications shall only be made when recommended by public health officials and trained personnel of mosquito abatement districts and other mosquito control programs. For best results, treat when insects are most active and meteorological conditions are conducive to keeping the spray cloud close to the ground. In order to compensate for windy conditions and ensure drift onto the target area aerial application with aircraft equipped with Global Positioning Systems (GPS) is recommended.

USE OF FIXED WING AIRCRAFT OR HELICOPTERS

UNDILUTED

Scourge may be applied undiluted at a treatment rate of 0.007 pounds of resmethrin per acre at a finished spray volume of 0.6 fl. oz. /acre.

DILUTED

For additional rates and dilutions see mixing instructions below.

MIXING INSTRUCTIONS FOR PRODUCT IN AIRCRAFT

USAGE

If using Scourge diluted follow the mixing instructions contained in the table for Aircraft usage.

FL oz/A of Finished Spray to be Applied	lb ai/A Wanted RESMETHRIN / PBO	Volumetric Dilution Needed	
		Product	Solvent
0.5	0.001/0.003 0.00175/0.00525 0.0035/0.0105	1 part	4.5 parts
		1 part	2.37 parts
		1 part	0.68 parts
1.0	0.001/0.003 0.00175/0.00525 0.0035/0.0105 0.007/0.021	1 part	10 part
		1 part	5.74 parts
		1 part	2.33 parts
		1 part	0.68 parts
3.0	0.001/0.003 0.00175/0.00525 0.0035/0.0105 0.007/0.021	1 part	34 parts
		1 part	19.2 parts
		1 part	9 parts
		1 part	4 parts
		1 part	4 parts

STORAGE AND DISPOSAL (continued)

Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with mineral oil and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill or incineration, or if allowed by State and Local authorities, by burning. If burned, stay out of smoke.

Rigid Non-refillable containers that are too large to shake (i.e., with capacities greater than 5 gallons or 50 lbs)

Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with mineral oil. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill or incineration, or if allowed by State and Local authorities, by burning. If burned, stay out of smoke.

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience LP. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE LP MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience LP is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE LP DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: TO THE EXTENT OF THE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE LP'S ELECTION, THE REPLACEMENT OF PRODUCT

Bayer (reg'd), the Bayer Cross (reg'd), Scourge® and Backed by Bayer™ are trademarks of Bayer.

Flow Rate (Fluid Ounces Per Minute)

Application Rate (lb resmethrin per Acre)	Vehicle Speed (MPH)	Flow Rate (Fluid Ounces Per Minute)				
		Undiluted	Diluted 1 + 1	Diluted 1 + 2	Diluted 1 + 4	Diluted 1 + 6
0.0015	10	0.77	1.56	2.32	3.90	5.40
	15	1.17	2.30	3.50	5.80	8.10
	20	1.54	3.10	4.62	7.70	10.80
0.00175	10	0.90	1.80	2.70	4.50	6.30
	15	1.35	2.70	4.05	6.75	9.45
	20	1.80	3.60	5.39	9.0	12.59
0.0035	10	1.80	3.60	5.39	9.00	12.59
	15	2.70	5.39	8.09	13.49	18.90
	20	3.60	7.20	10.80	17.98	25.17
0.007	10	3.59	7.19	10.79	17.98	25.17
	15	5.39	10.79	16.18	26.97	
	20	7.19	14.38	21.57		

In the State of New York: This product shall be applied at the rate of 0.001 to 0.007 pounds of resmethrin per acre with suitable diluent, not to exceed a finished spray solution of 16 fl. oz./acre.

Where dense vegetation is present, the use of the higher label rates is recommended.

AERIAL APPLICATION

Scourge may be applied either diluted or undiluted at rates of 0.0035 to 0.007 pounds resmethrin per acre by fixed wing or rotary aircraft that are capable of making a ULV application. Aerial application of Scourge should be made at a min-

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store product in its original container in a locked storage area

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal:

Rigid, Non-refillable containers small enough to shake (i.e., with capacities equal to or less than 5 gallons)



Produced for:
Bayer Environmental Science
A Division of Bayer CropScience LP
2 T. W. Alexander Drive
Research Triangle Park, NC 27709