



SEDGEHAMMER+ is a selective herbicide for the control of nutsedge and other weeds in turfgrass, ornamentals, landscaped areas and other specified non-crop areas.

| | % BY WT. |
|---|----------------------|
| ACTIVE INGREDIENT: | |
| Halosulfuron-methyl, methyl 3-chloro-5-(4,6-dimethoxyimidin-2-ylcarbamoylsulfamoyl) | |
| -1-methylpyrazole-4-carboxylate | 5.0% |
| OTHER INGREDIENTS: | 95.0% |
| | Total: 100.0% |

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se las explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

| FIRST AID | |
|---|---|
| IF IN EYES | <ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after 5 minutes, then continue rinsing eye. • Call poison control center or doctor for treatment advice. |
| HOT LINE NUMBER | |
| Have the product container or label with you when calling a poison control center, doctor, or going for treatment. For Medical emergencies involving this product, call toll free 1-888-478-0798. | |

PRECAUTIONARY STATEMENTS Hazards to Humans and Domestic Animals CAUTION

Causes moderate eye irritation. Avoid contact with eyes or clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- long-sleeved shirt and long pants
- shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls Statement: When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

| User Safety Recommendations: |
|--|
| Users should: <ul style="list-style-type: none"> • Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. • Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. |

ENVIRONMENTAL HAZARD SECTION OF PRECAUTIONARY STATEMENTS

GROUNDWATER ADVISORY

Halosulfuron-methyl is known to leach through soil into ground water under certain conditions as a result of label use. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow.

NET CONTENTS ____ OUNCES

EPA Reg. No. 81880-24-10163
EPA Est. No.

Distributed by:
Gowan Company
PO Box 5569
Yuma, AZ 85366-5569

SURFACE WATER ADVISORY

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of halosulfuron-methyl from runoff water and sediment. Runoff of this product will be greatly reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

WINDBLOWN SOIL PARTICLES

SEDGEHAMMER+ has the potential to move off-site due to wind erosion. Soils that are subject to wind erosion usually have a high silt and/or fine to very fine sand fractions and low organic matter. Other factors which can affect the movement of windblown soil include the intensity and direction of prevailing winds, vegetative cover, site slope, rainfall, and drainage patterns. Avoid applying SEDGEHAMMER+ if prevailing local conditions may be expected to result in off-site movement.

NON-TARGET ORGANISM ADVISORY

This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated area. Protect the forage and habitat of non-target organisms by minimizing spray drift.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water are:

- Coveralls,
- Shoes plus socks, and
- Chemical-resistant gloves made out of any waterproof material.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

- Keep people off treated areas until spray solution has dried.

PRODUCT INFORMATION

SEDGEHAMMER+ is a water dispersible granule (WDG) that selectively controls sedges such as purple and yellow nutsedge, Kyllinga species and certain broadleaf weeds (see weeds controlled chart). SEDGEHAMMER+ can be applied to commercial and residential turfgrass and on other non-crop sites including: airports, campgrounds, cemeteries, established woody ornamentals in landscape areas, fairgrounds, fallow areas, fuel storage areas, golf courses, landscaped areas, lumberyards, public recreation areas, race tracks, residential property, rights-of-ways, roadsides, school grounds, sports fields, tank farms and tennis courts.

WEED RESISTANCE STATEMENT

SEDGEHAMMER+ contains a Group 2 herbicide. Any weed population may contain or develop plants naturally resistant to (Group 2) Halosulfuron-methyl herbicides. Weed species with acquired resistance to (Group 2) Halosulfuron-methyl may eventually dominate the weed population if (Group 2) Halosulfuron-methyl herbicides are used repeatedly in the same field or in successive years as the primary method of control for targeted species. This may result in partial or total loss of control of those species by SEDGEHAMMER+ or other (Group 2) herbicides. Suspected herbicide-resistant weeds may be identified by these indicators:

- Failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds;
- A spreading patch of non-controlled plants of a particular weed species; and
- Surviving plants mixed with controlled individuals of the same species.

To delay herbicide resistance consider:

- Avoiding the consecutive use of SEDGEHAMMER+ or other target site of action Group 2 herbicides that have a similar target site of action, on the same weed species.
- Using tank mixtures or premixes with herbicides from different target site of action groups as long as the involved products are all registered for the same use, have different sites of action, and are both effective at the tank mix or prepack rate on the weed(s) of concern.
- Basing herbicide use on a comprehensive IPM program.
- Monitoring treated weed populations for loss of field efficacy.
 - Fields should be scouted prior to application to identify the weed species present and their growth stage to determine if the intended application will be effective.
 - Fields should be scouted after application to verify that the treatment was effective.

- Contacting your local extension specialist, certified crop advisors, and/or manufacturer for herbicide resistance management and/or integrated weed management recommendations for specific crops and resistant weed biotypes.
- For further information or to report suspected resistance, you may contact Gowan Company at 1-800-883-1844.

APPLICATION EQUIPMENT AND INSTRUCTIONS

Calibrate application equipment prior to application of this product. Application should be uniform and of sufficient volume to avoid streaking or skips. SEDGEHAMMER+ often works with 1 application, but depending on the size and age of the nutsedge a second treatment may be required 6 - 10 weeks after the initial treatment.

Herbicide symptoms are likely to show within 2 weeks as a necrotic ring at the base of the plant, even though the leaves and stems remain green and a deep leathery green in color. **Note:** This product already contains a non-ionic surfactant. No additional mixtures are necessary.

USE PRECAUTIONS

- For optimum results, do not mow turf for 2 days before or 2 days after application.
- Avoid applications when rainfall is forecasted to occur within 4 hours.
- This product may be used on established seeded, sodded, or sprigged turfgrass. Allow the turf to develop a good root system and uniform stand before application.
- Treated areas may be overseeded with annual or perennial ryegrass or bermudagrass, 2 weeks after application.
- Avoid application of SEDGEHAMMER+ when turfgrass or nutsedge is under stress since turf injury and poor nutsedge control may result.
- Avoid excessive overlap of the spray.
- Annual and perennial herbaceous ornamentals, including color plants, may be injured when transplanted into landscaped areas treated with SEDGEHAMMER+.

USE RESTRICTIONS

- Do not exceed 5 1/3 oz by weight of this product (0.25 lb active ingredient) per acre per 12 month period.
- No more than 4 applications can be made per year with a minimum of 14 days between applications.
- Do not store SEDGEHAMMER+ in solution. Once the product is mixed with water apply immediately.
- Do not apply as an over the top spray to desirable flowers, ornamentals, vegetables, shrubs or trees.
- Do not apply this product to golf course putting greens.
- Do not apply this product through any type of irrigation system.
- Do not apply this product by air

WEEDS CONTROLLED BY SEDGEHAMMER+

C = Control, S = Suppression, NA = No Activity

| WEED SPECIES | POSTEMERGENT ACTIVITY | WEED SPECIES | POSTEMERGENT ACTIVITY |
|--|-----------------------|--|-----------------------|
| Burcucumber <i>Sicyos angulatus</i> | S | Morningglory, tall & Ivyleaf <i>Ipomoea spp.</i> | S |
| Cocklebur, common <i>Xanthium strumarium</i> | C | Mustard, wild <i>Sinapis arvensis</i> | C |
| Corn Spurry <i>Spergula arvensis</i> | C | Nutsedge, yellow <i>Cyperus esculentus</i> | C |
| Flatsedge, rice <i>Cyperus iria</i> | C | Nutsedge, purple <i>Cyperus rotundus</i> | C |
| Fleabane, Philadelphia <i>Erigeron philadelphicus</i> | C | Pigweed, redroot <i>Amaranthus retroflexus</i> | C |
| Galinsoga <i>Galinsoga</i> | C | Pigweed, smooth <i>Amaranthus hybridus</i> | C |
| Groundsel, common <i>Senecio vulgaris</i> | NA | Pokeweed, common <i>Phytolacca Americana</i> | C |
| Jimsonweed <i>Datura stramonium</i> | NA | Purslane <i>Portulaca oleracea</i> | NA |
| Kyllinga, annual <i>Kyllinga sesquiflorus</i> | C | Radish, wild <i>Raphanus raphanistrum</i> | C |
| Kyllinga, green <i>Kyllinga breviflora</i> | C | Ragweed, common <i>Ambrosia artemisiifolia</i> | C |
| Kochia <i>Kochia scoparia</i> | S | Ragweed, giant <i>Ambrosia trifida</i> | C |
| Ladysthumb <i>Polygonum persicaria</i> | C | Shepherds purse <i>Capsella bursa-pastoris</i> | S |
| Lambsquarter, common <i>Chenopodium album</i> | NA | Smartweed, Pennsylvania <i>Polygonum pennsylvanicum</i> | C |
| Mallow, Venice <i>Hibiscus trionum</i> | C | Sunflower, common <i>Helianthus annuus</i> | C |
| Milkweed, common <i>Asclepias syriaca</i> | S | Velvetleaf <i>Abutilon theophrasti</i> | C |
| Milkweed, honeyvine <i>Ampelamus albidus</i> | S | | |

**SEDGEHAMMER+
RATES AND APPLICATION INSTRUCTIONS**

| CROP | RATE OZ/ 1000 SQ FT | DIRECTIONS FOR USE | | | | | | | | | | | | | | | | | | | | |
|--|--|--|---------------------|--|--|--|--|--|---|--|--|---------------------|--|--|--------------------------------------|---|---|---|--|---------------------------------------|--|---|
| TURFGRASS (INCLUDING COMMERCIAL AND RESIDENTIAL TURFGRASS, GOLF COURSES, ESTABLISHED LAWNS, AND ORNAMENTAL TURFGRASS) | 0.5 OZ (13.5 grams) or 1 pouch Mix in 1 gallon of water | Apply SEDGEHAMMER+ for postemergence control of purple or yellow nutsedge, Kyllinga ¹ and broadleaf weeds. Mix 0.5 oz (13.5 grams) of this product (1 pouch) in 1 gallon of water to treat 1000 sq ft of turfgrass. Mix or shake the solution to ensure ingredients are completely dispersed. Spray the target weeds thoroughly and wet the entire leaf surface of the undesirable plants. For best results, spray nutsedge after it has reached the 3 - 8 leaf stage of growth. ¹ For Optimum Kyllinga control 2 applications are recommended. | | | | | | | | | | | | | | | | | | | | |
| | 1 OZ (27 grams) or 2 pouches Mix in 1 gallon of water | For postemergence control of horsetail apply 1 oz (27 grams) of this product (2 pouches) in 1 gallon of water to treat 1000 sq ft after horsetail has leafed out. Note: This rate of product will control horsetail that is less than 6 inches tall and suppress horsetail that is greater than 6 inches tall. | | | | | | | | | | | | | | | | | | | | |
| | Apply SEDGEHAMMER+ as directed under the conditions described to the following established turfgrasses that are tolerant to this product: <table border="1" data-bbox="557 632 1360 1045" style="margin-left: auto; margin-right: auto;"> <thead> <tr> <th colspan="3" data-bbox="557 632 1360 667">Cool-Season Grasses</th> </tr> </thead> <tbody> <tr> <td data-bbox="557 667 789 732">Bentgrass, creeping <i>Agrostis stolonifera</i></td> <td data-bbox="789 667 1068 732">Fescue, Fine <i>Festuca arundinacea</i></td> <td data-bbox="1068 667 1360 732">Ryegrass, Perennial <i>Lolium perenne</i></td> </tr> <tr> <td data-bbox="557 732 789 793">Bluegrass, Kentucky <i>Poa pratensis</i></td> <td data-bbox="789 732 1068 793">Fescue, tall <i>Festuca arundinacea</i></td> <td data-bbox="1068 732 1360 793"></td> </tr> <tr> <th colspan="3" data-bbox="557 821 1360 856">Warm-Season Grasses</th> </tr> <tr> <td data-bbox="557 856 789 917">Bahigrass <i>Paspalum notatum</i></td> <td data-bbox="789 856 1068 917">Centipedegrass <i>Eremochioa ophiuroides</i></td> <td data-bbox="1068 856 1360 917">Kikuyugrass <i>Pennisetum clandestinum</i></td> </tr> <tr> <td data-bbox="557 917 789 978">Bermudagrass <i>Cynodon dactylon</i></td> <td data-bbox="789 917 1068 978">Seashore paspalum <i>Paspalum Vaginatum</i></td> <td data-bbox="1068 917 1360 978">Zoysiagrass <i>Zoysia Japonica</i></td> </tr> <tr> <td data-bbox="557 978 789 1039">Buffalograss <i>Bouteloua dactyloides</i></td> <td data-bbox="789 978 1068 1039">St. Augustine grass <i>Stenotaphrum secundatum</i></td> <td data-bbox="1068 978 1360 1039"></td> </tr> </tbody> </table> | | Cool-Season Grasses | | | Bentgrass, creeping <i>Agrostis stolonifera</i> | Fescue, Fine <i>Festuca arundinacea</i> | Ryegrass, Perennial <i>Lolium perenne</i> | Bluegrass, Kentucky <i>Poa pratensis</i> | Fescue, tall <i>Festuca arundinacea</i> | | Warm-Season Grasses | | | Bahigrass <i>Paspalum notatum</i> | Centipedegrass <i>Eremochioa ophiuroides</i> | Kikuyugrass <i>Pennisetum clandestinum</i> | Bermudagrass <i>Cynodon dactylon</i> | Seashore paspalum <i>Paspalum Vaginatum</i> | Zoysiagrass <i>Zoysia Japonica</i> | Buffalograss <i>Bouteloua dactyloides</i> | St. Augustine grass <i>Stenotaphrum secundatum</i> |
| Cool-Season Grasses | | | | | | | | | | | | | | | | | | | | | | |
| Bentgrass, creeping <i>Agrostis stolonifera</i> | Fescue, Fine <i>Festuca arundinacea</i> | Ryegrass, Perennial <i>Lolium perenne</i> | | | | | | | | | | | | | | | | | | | | |
| Bluegrass, Kentucky <i>Poa pratensis</i> | Fescue, tall <i>Festuca arundinacea</i> | | | | | | | | | | | | | | | | | | | | | |
| Warm-Season Grasses | | | | | | | | | | | | | | | | | | | | | | |
| Bahigrass <i>Paspalum notatum</i> | Centipedegrass <i>Eremochioa ophiuroides</i> | Kikuyugrass <i>Pennisetum clandestinum</i> | | | | | | | | | | | | | | | | | | | | |
| Bermudagrass <i>Cynodon dactylon</i> | Seashore paspalum <i>Paspalum Vaginatum</i> | Zoysiagrass <i>Zoysia Japonica</i> | | | | | | | | | | | | | | | | | | | | |
| Buffalograss <i>Bouteloua dactyloides</i> | St. Augustine grass <i>Stenotaphrum secundatum</i> | | | | | | | | | | | | | | | | | | | | | |
| ESTABLISHED WOODY ORNAMENTALS IN LANDSCAPED AREAS | 0.5 OZ (13.5 grams) or 1 pouch Mix in 1 gallon of water | Apply SEDGEHAMMER+ for postemergence control of purple or yellow nutsedge, Kyllinga ¹ and broadleaf weeds. Mix 0.5 oz (13.5 grams) of this product (1 pouch) in 1 gallon of water to treat 1000 sq ft. Spray around established woody ornamental species in landscaped areas. For transplanted woody ornamentals, allow 3 months after transplanting before applying this product. Mix or shake the solution to ensure ingredients are completely dispersed. Spray the target weeds thoroughly and wet the entire leaf surface of the undesirable plants. For best results, spray nutsedge after it has reached the 3 - 8 leaf stage of growth. ¹ For Optimum Kyllinga control 2 applications are recommended. | | | | | | | | | | | | | | | | | | | | |
| | 1 OZ (27 grams) or 2 pouches Mix in 1 gallon of water | For postemergence control of horsetail apply 1 oz (27 grams) of this product (2 pouches) in 1 gallon of water to treat 1000 sq ft after horsetail has leafed out. Note: This rate of product will control horsetail that is less than 6 inches tall and suppress horsetail that is greater than 6 inches tall. | | | | | | | | | | | | | | | | | | | | |
| FALLOW AREAS | 0.5 OZ (13.5 grams) or 1 pouch Mix in 1 gallon of water | Apply SEDGEHAMMER+ for postemergence control of purple or yellow nutsedge, Kyllinga ¹ and broadleaf weeds mix 0.5 oz (13.5 grams) of this product (1 pouch) in 1 gallon of water to treat 1000 sq ft. Spray on fallow areas prior to the establishment of turfgrass or woody ornamental plants. Allow 4 weeks between application and seeding or sodding of turfgrass, or transplanting woody ornamentals. Mix or shake the solution to ensure ingredients are completely dispersed. Spray the target weeds thoroughly and wet the entire leaf surface of the undesirable plants. For best results, spray nutsedge after it has reached the 3 - 8 leaf stage of growth. ¹ For Optimum Kyllinga control 2 applications are recommended. | | | | | | | | | | | | | | | | | | | | |
| | 1 OZ (27 grams) or 2 pouches Mix in 1 gallon of water | For postemergence control of horsetail apply 1 oz (27 grams) of this product (2 pouches) in 1 gallon of water to treat 1000 sq ft after horsetail has leafed out. Note: This rate of product will control horsetail that is less than 6 inches tall and suppress horsetail that is greater than 6 inches tall. | | | | | | | | | | | | | | | | | | | | |

| CROP | RATE OZ/ 1000 SQ FT | DIRECTIONS FOR USE |
|---|--|---|
| FENCE ROWS, FUEL STORAGE AREAS, LUMBERYARDS, TANK FARMS, RIGHT OF WAY, AND ROADSIDES | 0.5 OZ (13.5 grams) or 1 pouch Mix in 1 gallon of water | Apply SEDGEHAMMER+ for postemergence control of purple or yellow nutsedge, Kyllinga ¹ and broadleaf weeds mix 0.5 oz (13.5 grams) of this product (1 pouch) in 1 gallon of water to treat 1000 sq ft. Spray at specified rates on fence rows, fuel storage areas, lumberyards, tank farms, rights of ways, roadsides and other industrial sites. Mix or shake the solution to ensure ingredients are completely dispersed. Spray the target weeds thoroughly and wet the entire leaf surface of the undesirable plants. For best results, spray nutsedge after it has reached the 3 - 8 leaf stage of growth. This product may be tank mixed with Glyphosate herbicide. ¹ For Optimum Kyllinga control 2 applications are recommended. |
| | 1 OZ (27 grams) or 2 pouches Mix in 1 gallon of water | For postemergence control of horsetail apply 1 oz (27 grams) of this product (2 pouches) in 1 gallon of water to treat 1000 sq ft after horsetail has leafed out. Note: This rate of product will control horsetail that is less than 6 inches tall and suppress horsetail that is greater than 6 inches tall. |
| OTHER NON-CROP SITES (INCLUDING AIRPORTS, CAMPGROUNDS, CEMETERIES, FAIRGROUNDS, PUBLIC RECREATION AREAS, RACE TRACKS, RESIDENTIAL PROPERTY, SCHOOL GROUNDS, SPORTS FIELDS, AND TENNIS COURTS) | 0.5 OZ (13.5 grams) or 1 pouch Mix in 1 gallon of water | Apply SEDGEHAMMER+ for postemergence control of purple or yellow nutsedge, Kyllinga ¹ and broadleaf weeds mix 0.5 oz (13.5 grams) of this product (1 pouch) in 1 gallon of water to treat 1000 sq ft. Spray at specified rates on non-crop sites including: airports, campgrounds, cemeteries, fairgrounds, public recreation sites, race tracks, residential property, school grounds, sports fields, and tennis courts. |
| | 1 OZ (27 grams) or 2 pouches Mix in 1 gallon of water | For postemergence control of horsetail apply 1 oz (27 grams) of this product (2 pouches) in 1 gallon of water to treat 1000 sq ft after horsetail has leafed out. Note: This rate of product will control horsetail that is less than 6 inches tall and suppress horsetail that is greater than 6 inches tall. |

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store under cool, dry conditions (below 120°F). Do not store under moist conditions.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling, if available, or dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances.

DISPOSAL AUTHORITIES: If none of the foregoing procedures is permitted by state and local authorities, then contact your State Pesticide or Environmental control Agency, or your local Hazardous Waste Disposal office, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

FOR 24-HOUR EMERGENCY ASSISTANCE (SPILL, LEAK OR FIRE), CALL CHEMTREC® (800) 424-9300.

For more product information, call toll-free 1-800-883-1844.

NOTICE OF CONDITIONS OF SALE AND WARRANTY AND LIABILITY LIMITATIONS

Important: Read the entire Directions for Use and Notice of Conditions of Sale and Warranty and Liability Limitations before using this product. If terms are not acceptable return the unopened container for a full refund.

Our directions for use of this product are based on tests believed to be reliable. However, it is impossible to eliminate all risk associated with the use of this product. Crop injury, inadequate performance, or other unintended consequences may result due to soil or weather conditions, off target movement, presence of other materials, method of use or application, and other factors, all of which are beyond the control of Gowan Company. All such risks shall be assumed by the Buyer and User.

Gowan Company warrants that this product conforms to the specifications on the label when used in strict conformance with Direction for Use, subject to the above stated risk limitations. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, GOWAN COMPANY MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, GOWAN COMPANY'S EXCLUSIVE LIABILITY FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, OR ANY OTHER LEGAL THEORY IS STRICTLY LIMITED TO THE PURCHASE PRICE PAID OR REPLACEMENT OF PRODUCT, AT GOWAN COMPANY'S SOLE DISCRETION.

Chemtrec® is a registered trademark of American Chemistry Council, Inc.
SEDEHAMMER® is a registered trademark of Gowan Company.

01-R1219EPA