

EMULSIFIABLE CONCENTRATE

ACTIVE INGREDIENT	By Weight
Chlorpropham: Isopropyl-m-chlorocarbanilate	36.0%
OTHER INGREDIENTS	64.0%
TOTAL	100.0%

Contains 3 pounds active ingredient per gallon.

FOR AGRICULTURAL USE ONLY KEEP OUT OF THE REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if swallowed or absorbed through the skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Use only with adequate ventilation. Keep container closed when not in use.

EMERGENCY INFORMATION

For spill, leak, fire, exposure or accident call CHEMTREC $^{\! \odot}$ 1-800-424-9300. Outside the USA 1-703-527-3887

EPA Reg. No. 2749-70 EPA Est. No. 34704-MT-001

Batch Code: Net Contents: 5 Gallons

FIRST AID	
If Swallowed	Call a poison control center or doctor immediately for treatment advice.
	Have person sip a glass of water if able to swallow.
	Do not induce vomiting unless told to by a poison control center or doctor.
If in Eyes	Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.
	Remove contact lenses, if present, after the first 5 minutes, then continue
	rinsing eye.
	Call a poison control center or doctor for treatment advice.
If on Skin or	Take off contaminated clothing.
Clothing	Rinse skin immediately with plenty of water for 15 - 20 minutes.
	Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for	
treatment	

ENVIRONMENTAL HAZARD

This product is toxic to fish. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior

to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

PHYSICAL HAZARD AND CHEMICAL HAZARD

Do not store near heat or open flame. In case of fire, use water, spray, foam, dry chemical or CO2.

DIRECTIONS FOR USE

Notice to User: It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons either directly or through drift. Only protected handlers may be in the area during application.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category A on the EPA chemical resistance category selection chart.

Workers must wear: chemical resistant gloves (such as nitrile, butyl, neoprene, and/or barrier laminate) while handling potatoes which are wet.

Mixers, Loaders, Applicators and all other handlers must wear:

- Long sleeved shirt and long pants,
- Socks and shoes, and
- Chemical resistant gloves made of barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, polyvinyl chloride or viton.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

User Safety Recommendations:

Users should

- Follow manufacturer's instructions for cleaning/maintaining Personal Protective Equipment (PPE). If there are no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry
- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing.
 As soon as possible, wash thoroughly and change into clean clothing.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Avoid storage below freezing temperature. Prolonged storage may result in gel formulation. Warming contents and agitation will restore material to useable conditions, but do not heat above 250°F. Keep container closed. Do not contaminate water, food or feed by storage or disposal. This product inhibits germination of seed potatoes.

<u>PESTICIDE DISPOSAL</u>: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for quidance.


CONTAINER DISPOSAL:

Nonrefillable Containers 5 Gallons or Less: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke. Nonrefillable containers larger than 5 gallons: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

APPLICATION

SHIELD POTATO SPROUT INHIBITOR EC is applied at a 1% active ingredient level on washed and damp dried potatoes. The diluted material should be applied using spray nozzles placed evenly across the rolls on which the potatoes are being moved. The diluted solution can also be applied by a foaming system in conjunction with a suitable foaming agent. Do not apply this product using aerosol application equipment.

Do not apply more than one application of 0.0104 lbs of chlorpropham per 1000 lbs of potatoes.



DILUTION OF THE CONCENTRATE

Add 1 gallon of SHIELD POTATO SPROUT INHIBITOR EC to 35 gallons of water and mix well. A milky white emulsion will develop. Use this diluted emulsion at a rate of 1 gallon per 8,000 lbs of potatoes, (1 gallon for every 80 cwt. of potatoes). Do not reuse the diluted material that was prepared 24 hours or more earlier without mixing well.

NOTICE

SHIELD POTATO SPROUT INHIBITOR EC sprout inhibitor is used for the treatment of potatoes after harvest. Potatoes must be cured prior to treatment and, for maximum sprout inhibition, should be washed clean.

CONDITIONS OF SALE

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Aceto Life Sciences, L.L.C. To the extent consistent with applicable law, all such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, ACETO LIFE SCIENCES, L.L.C. MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Aceto Life Sciences, L.L.C. is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, ACETO LIFE SCIENCES, L.L.C. DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: To the extent consistent with applicable law, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID. OR AT ACETO LIFE SCIENCES. L.L.C. ELECTION. THE REPLACEMENT OF PRODUCT.

Manufactured For: Aceto Life Sciences L.L.C. 4 Tri Harbor Court Port Washington, NY 11050 Phone: 516-627-6000

Rev 05272020