



# ALLECTUS™ 0.225 G PLUS TURF FERTILIZER

## Insecticide

For fertilization and general insect control in turfgrass areas including athletic fields and parks and residential, commercial, industrial, institutional, and recreational lawns. Not for use on golf courses or sod farms.

**ACTIVE INGREDIENT:**

Imidacloprid*, 1-[(6-Chloro-3-pyridinyl)methyl]-N-nitro-2-imidazolidinimine .....	0.125%
Bifenthrin** .....	0.100%
<b>OTHER INGREDIENTS:</b> .....	99.775%
	100.000%

\* This product is covered by one or more of the following patents: U.S. Patent Nos. 4,742,060, 5,783,203, and 6,264,968. Other patents pending

\*\*Cis isomers 97% minimum, trans isomers 3% maximum.

Bifenthrin: U.S. Patent No. 4,238,505

**STOP - Read the label before use.**

## KEEP OUT OF REACH OF CHILDREN CAUTION

### FIRST AID

<b>If on skin:</b>	<ul style="list-style-type: none"> <li>Take off contaminated clothing.</li> <li>Rinse skin immediately with plenty of water for 15 to 20 minutes.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If in eyes:</b>	<ul style="list-style-type: none"> <li>Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.</li> <li>Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

**FOR A MEDICAL EMERGENCY INVOLVING THIS PRODUCT CALL: 1-866-944-8565.**

**Note To Physician:** No specific antidote is available. Treat the patient symptomatically.

This product contains a pyrethroid. If large amounts have been ingested, milk, cream and other digestible fats and oils may increase absorption and so should be avoided.

**EPA REG. NO. 432-1417-34704**

**EPA EST. NO. 84156-CAN-2 (Lot begins 2)**

**EPA EST. NO. 41124-OH-1 (Lot begins 0)**

**EPA Est. No. 41124-WI-1 (Lot begins W)**

**EPA EST. NO. 49991-TX-1 (Lot begins TX-1)**

**EPA EST. NO. 57131-WI-1 (Lot begins EC)**

**EPA EST. NO. 49991-TX-2 (Lot begins TX-2)**

**EPA EST. NO. 75527-PA-12 (Lot begins FS)**

**EPA EST. NO. 82757-MA-2 (Lot begins H)**

**EPA EST. NO. 82757-FL-1 (Lot begins S)**

**EPA EST. NO. 82757-OH-1 (Lot begins MF)**

**NET CONTENTS 50 LBS. (22.68 KG)**

IHT

C1110

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

### CAUTION

Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove contaminated clothing before reuse.

### ENVIRONMENTAL HAZARDS

This product is extremely toxic to fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Run-off may be hazardous to aquatic organisms in water adjacent to treated areas.

This product contains a chemical with properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirement specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**Do not formulate this product into other end-use products.**

### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If material is spilled for any reason or cause, carefully contain any spilled material to prevent non-target contamination. Do not walk through spilled material and dispose of as directed for pesticides above. Refer to Precautionary Statements on label for hazards associated with the handling of this material. In spill or leak incidents, keep unauthorized people away.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Dispose of empty bag in a sanitary landfill, by incineration, or if allowed by the state and local authorities, by burning. If burned, stay out of smoke. For help with any spill, leak, fire or exposure involving this material, call day or night CHEMTREC – 1-800-424-9300.

### GENERAL INFORMATION

#### APPLICATION TO TURFGRASS

The use of this product will provide plant nutrients as well as providing control of selected insect pests. ALLECTUS 0.225 PLUS TURF FERTILIZER can be used for the control of soil inhabiting pests of turfgrass, such as northern and southern masked chafers, *Cyclocephala borealis*, *C. immaculata*, and/or *C. lurida*; Asiatic garden beetle, *Maladera castanea*; black vine weevil, *Otiorynchus sulcatus*; European chafer, *Rhizotrogus majalis*; green June beetle, *Cotinis nitida*; May or June beetles, *Phyllophaga* spp.; Japanese beetle, *Popillia japonica*; oriental beetle, *Anomala orientalis*; *Euethola humulis rugiceps*; billbugs, *Sphenophorus* spp.; annual bluegrass weevil, *Hyperodes* spp.; black turfgrass ataenius, *Ataenius spretulus*; *Aphodius* spp.; European crane fly, *Tipula paludosa*; frit fly, *Oscinella frit*; chinch bugs, *Blissus* spp.; fire ants, *Solenopsis* spp.; cutworms, *Agrotis ipsilon*, *Peridroma saucia*, *Nephalodes menians*; armyworms, *Spodoptera* spp., *Pseudaletia* spp.; sod webworm, *Crambus* spp.; and mole crickets, *Scapteriscus* spp. ALLECTUS 0.225 PLUS TURF FERTILIZER can be used as directed on turfgrass in sites such as home lawns, business and office complexes, shopping complexes, multi-family residential complexes, airports, cemeteries, parks, playgrounds, and athletic fields.

The active ingredients ALLECTUS 0.225 PLUS TURF FERTILIZER have sufficient residual activity so that applications for control of subsurface feeders can be made preceding the egg laying activity. The need for an application for control of subsurface feeders can be based on historical monitoring of the site, previous records or experiences, current season adult trapping or other methods. Optimum control of subsurface feeders will be achieved when applications are made prior to egg hatch of the target pest, followed by sufficient irrigation or rainfall to move the active ingredient through the thatch.

The active ingredients in ALLECTUS 0.225 PLUS TURF FERTILIZER have sufficient knockdown and residual activity to provide curative and residual control of surface feeding pests. Applications for control of surface feeding pests can be made when infestations are anticipated based on historical monitoring of the site, previous records or experience, current season adult trapping or presence of insects at economic thresholds as deter-

# 25-0-0

GUARANTEED ANALYSIS:

Total Nitrogen (N) .....25.00%  
25.00% Urea Nitrogen\*

**DERIVED FROM:** Polymer Coated Sulfur Coated Urea, Urea.

\*6.69% Coated slow release nitrogen from Polymer Coated Sulfur Coated Urea.

**WARNING: THIS PRODUCT CONTAINS CHEMICALS KNOWN TO THE STATE OF CALIFORNIA TO CAUSE CANCER OR BIRTH DEFECTS OR OTHER REPRODUCTIVE HARM**

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.regulatory-info-lpi.com>  
For Golf Courses follow the Best Management Practice at: <http://www.dep.state.fl.us/water/nonpoint/docs/nonpoint/glfbmp07.pdf>  
Please follow the Best Management Practice for Florida Green Industries at: [http://www.dep.state.fl.us/central/Home/MeetingsTraining/FLGreen/BMP\\_Book\\_final.pdf](http://www.dep.state.fl.us/central/Home/MeetingsTraining/FLGreen/BMP_Book_final.pdf)

Product No. 8111130

F1548

Contains XCU Polymer Coated Sulfur Coated Urea, a trademark of Agrium Advanced Technologies



### GENERAL APPLICATION GUIDELINES:

Rate	lbs. Product per 1,000 sq ft	lbs. Nitrogen per 1,000 sq ft	lbs. A.I. per acre
<b>Low</b>	<b>3.70</b>	<b>0.93</b>	<b>0.36</b>
<b>Med</b>	<b>4.60</b>	<b>1.15</b>	<b>0.45</b>
<b>High</b>	<b>9.20</b>	<b>2.30</b>	<b>0.90</b>



DISTRIBUTED BY  
**LOVELAND PRODUCTS, INC.**  
P.O. BOX 1286, GREELEY, COLORADO 80632-1286

mined by scouting and/or recommendations of local State extension personnel or other qualified specialists.

## SIGNATURE ALLECTUS™ 0.225 G PLUS TURF FERTILIZER Insecticide EPA REG. NO. 432-1417-34704

Do not apply more than 200 lb. of ALLECTUS 0.225 PLUS TURF FERTILIZER (0.25 lb. of imidacloprid, 0.2 lb. bifenthrin) per acre per application on residential use sites (i.e., around private homes, apartment buildings, condominiums, non-agricultural outbuildings, non-commercial greenhouses, pre-schools or day care facilities). May be applied at up to 400 lb. of ALLECTUS 0.225 PLUS TURF FERTILIZER (0.5 lb. of imidacloprid, 0.4 lb. bifenthrin) per application on non-residential use sites (i.e., around institutional, public, commercial or industrial buildings; parks; recreational areas or athletic fields).

Applications should not be made when turfgrass areas are waterlogged or the soil is saturated with water. Adequate distribution of the active ingredient cannot be achieved when these conditions exist. The treated turf must be in such a condition that the rainfall or irrigation will penetrate vertically in the soil profile. Total amount applied cannot exceed 400 lb (0.5 lb of imidacloprid, 0.4 lb of bifenthrin) per acre per year.

**NOTE:** Not for use in commercial greenhouses or on grass grown for seed or on golf courses and sod farms. For use on plants intended for aesthetic purposes or climatic modification and being grown in interior landscapes, ornamental gardens or parks, or on lawns and grounds.

### APPLICATION EQUIPMENT

Apply ALLECTUS 0.225 PLUS TURF FERTILIZER uniformly over the area being treated with normally used granular application equipment. Both drop-type and rotary-type spreaders may be used to apply ALLECTUS 0.225 PLUS TURF FERTILIZER. Avoid the use of spreaders that apply the material in narrow, concentrated bands. Calibrate application equipment prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly and accurately.

### RECOMMENDED APPLICATIONS

(Be sure to read "APPLICATION EQUIPMENT" Section of this label)

CROP	PEST	DOSAGE ALLECTUS 0.225 PLUS TURF FERTILIZER
Turfgrasses	<b>Larvae of subsurface feeders:</b>	160 to 200 lb per acre (3.7 to 4.6 lb per 1000 ft sq)
	Asiatic garden beetle Black turfgrass ataenius Black vine weevil European chafer European crane fly	Green June beetle Japanese beetle Northern masked chafer Oriental beetle <i>Phyllophaga</i> spp. Southern masked chafer
	<b>Adults and/or larvae of subsurface and surface feeding pests:</b>	200 lb per acre (4.6 lb per 1000 ft sq) or for heavy infestations, a single annual application of 320 to 400 lb per acre (7.3 to 9.2 lb per 1000 sq ft) to non-residential turf
	Centipedes Fleas Millipedes Ticks	Imported fire ants <sup>1</sup> Mole crickets <sup>2</sup> Pillbugs <sup>3</sup> Sowbugs <sup>3</sup>
	<b>Adults and/or larvae of surface feeders:</b>	120 to 200 lb per acre (2.8 to 4.6 lb per 1000 ft sq)
	Annual bluegrass weevil Billbugs Chinch bugs	Frit fly Leafhoppers Mealybugs Nuisance ants
	<b>Larvae of surface feeders:</b>	80 to 200 lb per acre (1.8 to 4.6 lb per 1000 ft sq)
	Armyworm Cutworm	Sod webworms

Consult your local State Agricultural Experiment Station or State Extension Turf Specialists for more specific information regarding timing of application.

**NOTE:** For optimum control, irrigation or rainfall should occur within 24 hours after application to move the active ingredient through the thatch. Do not apply more than 400 lb product (0.5 lb of imidacloprid active ingredient, 0.4 lb bifenthrin) per acre per year. Do not apply more than 200 lb product (0.25 lb imidacloprid, 0.2 lb bifenthrin) per acre per application to residential turf. Avoid mowing turf or lawn area until after irrigation or rainfall has occurred so that uniformity of application will not be affected.

In New York State, this product may NOT be applied to any grass or turf area within 100 feet of a water body (lake, pond, river, stream, wetland, or drainage ditch).

In New York State, make a single repeat application of ALLECTUS 0.225 PLUS TURF FERTILIZER if there

are signs of renewed insect activity, but not sooner than two weeks after the first application.

<sup>1</sup>Imported Fire ants: ALLECTUS 0.225 G Plus Turf Fertilizer may be applied as a broadcast treatment in non-residential turf areas up to 9.2 lb. per 1000 sq. ft. For mound treatment, apply 3/4 cup ALLECTUS 0.225 G Plus Turf Fertilizer per mound and then drenching the mound with 1 to 2 gallons of water. The mounds should be treated with sufficient force to break their apex and allow the insecticide to flow into the ant tunnels.

<sup>2</sup>Mole crickets: For maximum effectiveness, apply ALLECTUS 0.225 PLUS TURF FERTILIZER as a broadcast treatment at 4.6 lb. per 1000 sq. ft. prior to or during the peak egg hatch period. For application when both adults and/or large nymphs are present and actively tunneling, irrigate prior to application if soil is dry and apply 4.6 lb. per 1000 sq. ft. Rainfall or additional irrigation within 24 hours after application will improve effectiveness of ALLECTUS 0.225 PLUS TURF FERTILIZER. For heavily infested mole cricket areas, use a single seasonal application rate of 320 to 400 lb per acre in non-residential turf.

<sup>3</sup>Pillbugs and sowbugs: For heavy infestation use 400 lbs per acre (9.2 lbs per 1000 sq. ft.) of ALLECTUS 0.225 G Plus Turf Fertilizer in non-residential turf.

### Spreader settings for ALLECTUS 0.225 G Plus Turf Fertilizer

The following chart can be used as a guide in establishing an accurate rate of ALLECTUS 0.225 PLUS TURF FERTILIZER. Application rates will vary depending upon speed, equipment wear and condition of spreader. Carefully calibrate application equipment prior to commercial use of this product.

Spreader	1.8 lbs. per 1000	3.7 lbs. per 1000	4.6 lbs. 1000
Andersons/Scotts AccuPro 1000	J	K	L <sup>3</sup> / <sub>4</sub>
Andersons/Scotts AccuPro 2000			
Cone 4	J	K	L <sup>3</sup> / <sub>4</sub>
Cyclone	3	3 <sup>3</sup> / <sub>4</sub>	4 <sup>1</sup> / <sub>2</sub>
Earthway 2400	14	16	18
Lely Models WTR, WRF, HR, 1250 (PTO @ 450 RPM)	4-III A	4 <sup>1</sup> / <sub>2</sub> -III A	5 <sup>1</sup> / <sub>4</sub> -III A
Scotts R8A			
Cone 4	H <sup>1</sup> / <sub>2</sub>	I <sup>1</sup> / <sub>2</sub>	J <sup>3</sup> / <sub>4</sub>
Spyker	3	3 <sup>3</sup> / <sub>4</sub>	4 <sup>1</sup> / <sub>2</sub>
Vicon 03 Series	20	24	30

### RESTRICTIONS

Avoid runoff or puddling of irrigation water following application. Avoid application of ALLECTUS 0.225 PLUS TURF FERTILIZER to areas which are waterlogged or saturated, which will not allow penetration into the root zone of the plant. Do not apply more than 400 lb (0.5 lb of imidacloprid and 0.4 lb of bifenthrin) per acre per year.

Treated areas may be replanted with any crop specified on an imidacloprid and a bifenthrin label, or with any crop for which a tolerance exists for the two active ingredients.

For crops not listed on any imidacloprid and a bifenthrin label, or for crops for which no tolerances for the active ingredients have been established, a 12-month plant-back interval should be observed.

Do not apply by air.

Not for use in nurseries or commercial greenhouses.

### IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of liability before using this product.

If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following conditions, disclaimer of warranties and limitations of liability.

CONDITIONS: The directions for use of this product are believed to be adequate and should be followed carefully. However, because of manner of use and other factors beyond Loveland Products, Inc.'s control it is impossible for Loveland Products, Inc. to eliminate all risks associated with the use of this product. As a result, crop injury or ineffectiveness is always possible. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: LOVELAND PRODUCTS, INC. MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Loveland Products, Inc. is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. Loveland Products, Inc. disclaims any liability whatsoever for special, incidental or consequential damages, resulting from the use or handling of this product.

LIMITATIONS OF LIABILITY: THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT LOVELAND PRODUCTS, INC.'S ELECTION, THE REPLACEMENT OF PRODUCT.

This Product contains imidacloprid, and it is specially formulated and sold under license from Bayer only for the uses set forth on this label. This active ingredient contained in this product, formulations containing this active ingredient, and/or methods of making or using this active ingredient or its formulations may be the subject of one or more Bayer patents, including but not limited to U.S. Patent No.4,742,060 5,783,203 and 6,264,968. The purchase and/or use of this product constitutes acceptance of a limited license to use this product only in strict accordance with the instructions set forth on this label. Any use of this product other than as expressly set forth on this label is prohibited and is not licensed through the purchase and/or use of this product and may result in the infringement of one or more Bayer patents and/or other intellectual property rights.