

KEEP OUT OF REACH OF CHILDREN

CAUTION

Si usted no entienda la etiqueta, busque a algulen para que se la explique a usted an detaile. (If you do not understand the label, find someone to explain it to you in detail).

FIRST AID

If swallowed

- Call poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor.

Do not give anything by mouth to an unconscious person.

If inhaled

mouth-to-mouth, if possible.

Move person to fresh air.

If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by

If in eyes

Call a poison control center or doctor for further treatment advice.

• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

Call a poison control center or doctor for treatment advice.

- If on skin or clothing
 - Take off contaminated clothing. Rinse skin immedialely with plenty of water for 15-20 minutes.

- Call a poison control center or doctor for treatment advice.

Exposed individuals and animals should receive emergency care as quickly as possible.

Poisoning Hotline

Seek medical care as soon as possible after exposure. For incidents involving animals, seek veterinary care as soon as possible after exposure. Have the product container or label with you when you call a poison control center or doctor or veterinarian, or when going for treatment. For information on this pesticide product (including health concerns and pesticide incidents), call the National Pesticide Information Center at 1-800-858-7378.

Note to Physician: Dust inhalation is usual route of accidental exposure; the product is rarely ingested. Treatment is symptomatic and supportive. Monitor severe exposures closely: seizures, electrolyte imbalances, acidosis, dehydration, and increased body temperature have been reported.

Metaldehyde is a systemic toxin. There is no antidote. The following instructions are not intended for laypeople. There is no home treatment.

Note to Veterinarian: Animals typically are exposed via ingestion. Treatment is symptomatic and supportive. If significant ingestion is suspected, and the animal is asymptomatic, consider induction of emesis or gastric lavage and administration of activated charcoal, cathartic and enema to

limit absorption. Monitor asymptomatic and symptomatic animals closely. Treat seizures, electrolyte imbalances, acidosis, dehydration, and increased body temperature. Monitor liver function post recovery.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Harmful if inhaled. Causes moderate eye irritation. Harmful if absorbed through the skin. Avoid contact with eyes, skin, or clothing. Avoid breathing dust. IMPORTANT: This product can be harmful to children and fatal to domestic animals when ingested. Children and dogs may be attracted to the product. Application of this product is prohibited unless children and domestic animals can be excluded from the treated area from start of the application until applied material is no longer visible.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Mixers, loaders, applicators and other handlers must wear:

- Long-sleeved shirt and long pants Shoes plus socks
- Chemical resistant gloves made of: barrier laminate, butyl rubber ≥14 mils, nitrile rubber ≥14 mils, neoprene rubber ≥ 14 mils, natural rubber ≥14 mils, polyethylene polyvinyl chloride ≥14 mils, or viton ≥14 mils.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

USER SAFETY RECOMMENDATIONS

Users should

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

exterior areas around schools and daycare facilities, and athletic fields, playgrounds, parks, recreation areas, etc.

Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to mollusks (e.g., clams, oysters, scallops, mussels). Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwaters or rinsate. This pesticide is toxic to birds and mammals. Granules on soil surface may be hazardous to terrestrial wildlife. Cover or collect any such materials spilled during loading.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. Read entire label before using. This product is solely for use on agricultural crops grown for commercial or research purposes, and in State and/or Federal invasive mollusk eradication

Broadcast applications and applications of this product directed to plant parts are prohibited except to grass (grown for seed and/or forage and hay), taro, corn and soybeans, wheat, and clover (grown for seed), and as specified for use in State and/or Federal invasive mollusk eradication operations.

operations. This product may not be applied to residential sites (i.e., in yards, gardens, around homes and apartments) and other sites such as the

FOR STATE AND/OR FEDERAL INVASIVE MOLLUSK ERADICATION OPERATIONS

This product may be used only on the following use sites in response to State and/or Federal invasive mollusk eradication operations.

Broadcast applications of this product may be made to the following use-sites for State and/or Federal invasive mollusk eradication operations: fallow land; barrier strips; uncultivated non-agricultural areas; recreational areas; non-food or non-feed brush; weed or dense vegetative areas; railroad, pipeline, highway, power and telephone rights of way and roadsides; guardrails and fences; lumberyards; storage areas; seaports; industrial facility sites, including yards and walkways around industrial buildings, parking areas, parks, golf courses, other public areas; and airports and similar industrial non-crop areas.

This product may be applied for State and/or Federal invasive mollusk eradication operations at an individual application rate of no more than 2 lb ai/A, and at no more than 6 applications/year.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) notification to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

DO NOT enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- chemical resistant gloves made of any water proof material,
- and shoes plus socks.

OR-CAL, INC. 29454 Meadowview Road Junction City, OR 97448

Phone: (541) 689-4413



NET WT. 40 lbs.



NET WT. 40 lbs.

SLUG & SNAIL BAIT

Phone: (541) 689-4413

Junction City, OR 97448 beoA weivwobseM 42492 OR-CAL, INC.

shall manufacturer be liable for consequential, special or indirect damage resulting from the use or handling of this product. the inherent risks referred to above. Manufacturer makes no other express or implied warranty. To the extent consistent with applicable law, in no case product conforms to the chemical description on the label and is reasonably fit for the specific purposes referred to in the Directions for Use, subject to Important: Read these entire Directions and Conditions of Sale before using Slugger 4.0 Slug & Snail Bait. Disclaimer: Manufacturer warrants that this

CONDITIONS FOR SALE

of empty bag in a sanitary landfill or by incineration. Container Disposal: Monrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then dispose

remaining product to a waste disposal facility or pesticide disposal program (often such programs are run by state or local governments or by Pesticide Disposal: To avoid wastes, use all material in this container by application according to label directions. If wastes cannot be avoided, offer ingest metaldehyde after opening or tearing packaging. Store this product in its original packaging, in a cool, secure location and out of reach of

Pesticide Storage: This product can be harmful to children and fatal to dogs and other domestic animals if ingested. Dogs have been known to Do not contaminate water, food or feed by storage and disposal.

STORAGE AND DISPOSAL

Soybean – http://crops.extension.iastate.edu/soybean/production_growthstages.html Corn – https://graincrops.ca.uky.edu/files/corngrowthstages_2011.pdf Growth stages are according to State University Crop Guides:

broadcast, directing the product to the soil.

pod formation. Apply as a perimeter treatment at planting (up to V4 growth stage) and then for subsequent applications (V4-R1 growth stage) by **Metaldehyde treated soybeans may not be fed to livestock and livestock may not graze in the treated fields. Applications must be completed prior to quent applications (V8-VT growth stage) by broadcast, directing the product to the soil. *Corn applications must be completed no later than flag stage. Apply as perimeter treatment at planting (up to V8 growth stage) and then for subse-

Plants	SZ	ιz	9	0	9
"Soybean (Only for use in the following states: Alabama, Arkansas, Connecticut, Delaware, Georgia, ouisiana, Maine, Maryland, Massachusetts, Mississippi, North Carolina, New Hampshire, New Jersey, lew York, Ohio, Pennsylvania, Rhode Island, South Carolina, Tennessee, Vermont, Virginia)	01	Ĺ	٤	0	٤
Corn (field and sweet)	SZ	۷	Þ	0	₽
aro (wetland): Apply by broadcast application under flooded conditions only. Prior to bait pplication, stop the water flow into and out of the lof (paddies) to be treated; water flow into and out of the lof (paddies) to be treated; water for a igh enough such that apple snails don't estivate. Apply the bait and holdimpound the water for exit the lof. Do not apply more than one application into 3 day before allowing water level in the lof. Do not apply more than one application er helding the beding the beding the satist percolation of water through the soil), such that apple snails begin to estivate, low additional water in the lof.	SZ	SE	Ĺ	L	Ĺ
fint (Peppermint, Spearmint, and their Oils)	57	Ιζ	Þ	ι	Þ
(Yest bne sgesor) szeri	50	₽l	Þ	12	Þ
(beez nof nwong) azsari	50	ιz	Þ	ız	Þ
Vheat (forage, grain, hay, straw) or use only in Oregon, Washington & Idaho	52	ÞÍ	ζ	ot roir9 to noitsemot bead baas	Þ
noegon (i or use أم Oregon, المراy for use in Oregon, المراه أمال (and tops) المراه أمال (and tops) والمراهبة	52	⊅l	ζ	50	Þ
ettuce, Cole Crops and the following Leafy Greens: broccoli, Brussels sprouts, cabbage, cauliflower, avalo, broccolo, collards, kale, kohlrabi, mizuna, mustard greens, spinach, rape greens	SZ	₽l	ε	0	9
seoseno	57	₽l	ε	0	9
orlebi & In Oregon, Washington & Idaho	52	₽l	7	0 1	Þ
rickly Pear cactus	70	98	ε	0	٤
eas and Bean (forage and hay)	52	L	٤	ι	9
ue geg pue sea	SZ	L	٤	Pre-Bloom	9
lops Only for use in Oregon, Washington & Idaho	SZ	₽Ĺ	Þ	09	Þ
биәѕиі	SZ	ız	9	ız	9
	sdl) ətsA Product/A)	ketreatment	Applications Per Season	INVETVAI	ber year applications

%00.00 f

%00'96

%00.₽

71-80-50-51-96017

17 52 Celery, Rhubarb, Swiss Chard, Chinese Celery, Celtuce, Cardoon, and Florence Fennel Only for use in Oregon, Washington & Idaho 10 70 Not for use in California Oι 70 Clover (grown for seed) 52 52 sea buckthorn), Berry-low growing (bearberry, bilberry, blueberry, cloudberry, muntries, lowberry, marionberry, olallieberry, red raspberry, wild raspberry, youngberry, Bushberries (aronia berry, blueberry, buffalo currant, Chilean guava, black currant, red currant, elderberry, European barberry, puneberry, cannerry, cannerry, slangonberry, salal, Berries: Caneberries (bingleberry, black raspberry, blackberry, boysenberry, dewberry, loganberry, 0 3 ħί 70 Only for use in Oregon, Washington & Idaho **CROP USE SITE**

For clover (forage and hay), apply as a perimeter treatment. Corn and Soybeans: Apply as a perimeter treatment at planting and for subsequent applications by broadcast, directing the product to the soil.

For clover (grown for seed), apply as a perimeter treatment at planting and then for the second application, by broadcast, directing the product to the

For grass (grown for seed and/or forage and hay), taro, and wheat apply by broadcast application. For use on citrus, apply in a band under tree canopy with commercial application equipment designed for banded application of granular products.

towards base of the plants with commercial application equipment designed for banded application of granular products.

ginseng, beans, peas, prickly pear cactus, tomatoes, lettuce, Cole crops and listed leafy greens and mint apply in a band, between rows and/or directed For use on ornamental plants, artichokes, listed berries, strawberries, celery, rhubarb, Swiss chard, Chinese celery, celtuce, cardoon, Florence fennel, day, but evening applications are preferred. will offer protection to gardens for several weeks. This product may be injurious to Daylilies and Clematis. Do not use on these plants. Apply any time of

plants, and grass (grown for seed and/or forage and hay). When applied as directed, Slugger 4.0 Slug & Snail Bait will withstand rain and sprinkling and Slugger 4.0 Slug & Snail Bait is a soil surface treatment for control of snails and slugs in citrus groves, berry crops, listed vegetable crops, ornamental

Do not use in or around Hanalei, Kaua'i (USFW Refuge)

- Do not re-water for 48 hours after application. Do not apply to dry soil. Make applications following irrigation.
- Grower is responsible for residues on crops. Use as band treatment between rows, unless otherwise directed on this label.
- Use only as a baiting treatment to be applied directly to the soil area beneath listed plants. Do not allow bait pellet to come in contact with plant parts.

directing the product to the soil (Not for use in California).

- directing the product to the soil. Corn and Soybeans: Apply as a perimeter treatment at planting and for subsequent applications by broadcast,
- Clover (grown for seed): Apply as a perimeter treatment at planting and for subsequent applications by broadcast, For greas grown for seed, greas grown for forage and hay, taro, and wheat: Apply by broadcast application. When broadcast applications are allowed, apply using standard broadcast spreaders.
 - specified for use in State and/or Federal invasive mollusk eradication operations. (grown for seed and/or forage and hay), taro, corn and soybeans, wheat, and clover (grown for seed), and as
- Broadcast applications and applications of this product directed to plant parts are prohibited except to grass recreation areas, etc. sites such as the exterior areas around schools and daycare facilities, and athletic fields, playgrounds, parks, This product may not be applied to residential sites (i.e., in yards, gardens, around homes and apartments) and other
- Federal invasive mollusk eradication operations. This product is solely for use on agricultural crops grown for commercial or research purposes, and in State and/or

RESTRICTIONS

houses. Do not enter or allow others to enter the treated area until applied material is no longer visible. pesticides 40 CFR Part 170. The WPS applies only when this product is used to produce agricultural plants on farms, forests, nurseries, or green-The requirements in this box apply to uses of the product that are NOT within the scope of the Worker Protection Standard, for agricultural

NON-AGRICULTURAL USE REQUIREMENTS

ACTIVE INGREDIENT:

Other Ingredients Metaldehyde

Controlled Release Pellet to Control Slugs and Snails. Around Listed Vegetables, Citrus, and Ornamentals. Controls Giant African Land Snail.

EPA Est. No. 66876-OR-001 EPA Reg. No. 71096-13 MATA CARACOLES Y BABOSAS SLUG & SNAIL BAIT C 2 C C E B + O IA) A(

CROP USE SITE