

STart ^{UP}™ METXL 480

Seed Treatment

ACTIVE INGREDIENT:

Metalaxyl: N-(2,6-dimethylphenyl)-N-(methoxyacetyl) alanine methyl ester 42.50%

OTHER INGREDIENTS: 57.50%

TOTAL: 100.00%

Contains 4.0 pounds Metalaxyl per gallon (480 grams per liter).

EPA Reg. No. 70506-356

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

FIRST AID

IF SWALLOWED	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have the person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800)-424-9300

For Medical Emergencies Only, Call Rocky Mountain Poison and Drug Safety: 1-866-673-6671

For Product Use Information Call 1-866-761-9397

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Some materials that are chemical-resistant to this product are listed below.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants,
- Shoes and socks,
- Chemical-resistant gloves (barrier laminate, butyl rubber \geq 14 mils, neoprene rubber \geq 14 mils, nitrile rubber \geq 14 mils, polyvinyl chloride (PVC) \geq 14 mils, or Viton \geq 14 mils).

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering controls statement:

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users Should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwater or rinsate.

Groundwater Advisory Statement: This chemical is known to leach through soil into ground water under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

Physical and Chemical Hazards

Do not mix or come into contact with oxidizing agents. Hazardous chemical reaction may occur.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard (WPS), 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statement of this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to users of this product that are covered by the WPS.

Do not enter or allow worker entry into treated areas during the REI of 48 hours.

Exception: If the product is soil-injected or soil-incorporated, the WPS, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is: coveralls, shoes plus socks and chemical-resistant gloves such as barrier laminate or nitrile rubber or neoprene rubber or Viton and protective eyewear.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the WPS for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural crops on farms, forests, nurseries, or greenhouses.

Do not enter or allow other to enter the treated area until sprays have dried.

USE PRECAUTIONS

STARTUP™ METXL 480 Seed Treatment is a systemic fungicide that controls certain diseases caused by members of the Oomycete class of fungi. When using on the same crop within the same season as mefenoxam, the maximum annual usage applies to the combination of both active ingredients when used on the same crop cycle in the same season.

Resistance Management Recommendations

For resistance management, STARTUP METXL 480 contains a Group 4 fungicide. Any fungal population may contain individuals naturally resistant to STARTUP METXL 480 and other Group 4 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of STARTUP METXL 480 or other Group 4 fungicides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicides from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
- For further information or to report suspected resistance contact UPL NA Inc. at 1-866-761-9397. You can also contact your pesticide distributor or university extension specialist to report resistance.

ROTATION (PLANTBACK) RESTRICTIONS

ROTATION CROP	PLANTING TIME FROM LAST APPLICATION (Days)
Any crop specified on this label	0
Bushberry Subgroup 13-07B	0
Caneberry Subgroup 13-07A	0
Corn	0
Cereal Grains (other than Corn)	14
Crops Not Intended for Food or Feed	0
All Other Crops Intended for Food or Feed	12 months

Replanting

If replanting is necessary, additional applications of this product may be made, provided that the total amount of metalaxyl applied does not exceed the maximum allowed for the specific crop.

SEED TREATMENT

This product is a systemic fungicide seed dressing specifically for control of systemic downy mildews, *Pythium* and *Phytophthora* spp. This product may be applied as a water-based slurry with other registered seed treatment insecticides and fungicides through standard slurry or mist-type commercial seed treatment equipment.

Precautions:

This product is a systemic fungicide having a specific mode of action. This product could be subject to development of resistant strains of fungi. Development of resistance cannot be predicted. Consult with your State Agricultural Experiment Station or Extension Service Specialist for guidance and ways to control any possible Metalaxyl resistant strains of fungi which may occur. FAILURE TO FOLLOW DIRECTIONS OF THIS LABEL MAY RESULT IN CROP INJURY OR POOR DISEASE CONTROL.

Restrictions:

DO NOT use with other seed treatment products unless previous experience assures compatibility.

In California: Only for use in commercial seed treating establishments. Not allowed for "on-farm" use.

On-Farm Seed Treatment: **DO NOT** apply this product through any type of irrigation system.

DO NOT use treated seed for food or feed purposes or process for oil. Treat only those seeds needed for immediate use, minimizing the interval between treatment and planting. **DO NOT** store excess treated seeds beyond planting time.

Seed treatment on agricultural establishment at or immediately before planting is within the scope of WPS, while commercial treatment of seeds is not within the scope.

Planter Box Treatment: Apply at the specified rate and premix with the seed directly in the planter box at planting. Place half the amount of seed into the planter box and apply half the required amount of pesticide to the seed. Stir the seed thoroughly with a paddle until the seed is uniformly coated. Place the remaining half of the seed into the planter box and add the remaining half of the pesticide. Again stir the seed with the paddle until all seed is thoroughly coated.

For Commercial Seed Treatment: Seed treated commercially or for sale must be dyed an unnatural color using an EPA approved dye before sale. Federal law requires that bags containing treated seed shall be labeled with the following information:

Tank Mixing: When tank mixing follow the most restrictive label instructions for any tank mix partner. **DO NOT** tank mix with any product that contains a prohibition against tank mixing.

SEED BAG LABELING REQUIREMENTS

Seed commercially treated with this product must be labeled in accordance with all applicable requirements of the Federal Seed Act. The user of this product is responsible for ensuring that the seed bag meets all requirements under the Federal Seed Act.

THE FEDERAL SEED ACT REQUIRES THAT BAGS CONTAINING TREATED SEEDS BE LABELED WITH THE FOLLOWING STATEMENTS:

This seed has been treated with STARTUP METXL 480 Seed Treatment, a fungicide containing Metalaxyl.

DO NOT use for feed, food, or oil purposes.

THE US ENVIRONMENTAL PROTECTION AGENCY REQUIRES THE FOLLOWING STATEMENTS ON CONTAINERS CONTAINING SEEDS TREATED WITH Metalaxyl (STARTUP METXL 480 SEED TREATMENT):

DO NOT allow children, pets, or livestock to have access to treated seeds.

Wear long pants, long-sleeved shirt, shoes plus socks, and chemical-resistant gloves when opening this bag or handling (e.g., loading, pouring) treated seed.

Treated seed exposed on soil surfaces may be hazardous to wildlife. Cover or collect treated seed spilled during loading and planting.

DO NOT contaminate bodies of water when disposing of planting equipment washwater. *If excess treated seed remains after planting, dispose of it by burying away from bodies of water.*

Store treated seed in cool, dry conditions.

Dispose of treated seed packaging in accordance with local requirements.

Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticides remain in the ethanol by-products that are used in agronomic practice.

DO NOT use the same auger for treating seed that will handle commercial or feed grains.

DO NOT plant any crop not listed on this label within 30 days of planting STARTUP METXL 480 Seed Treatment treated seed.

DO NOT harvest mustard greens. *Rape greens grown and harvested from STARTUP METXL 480 Seed Treatment treated seed must not be used for human and feed consumption. Rapeseed grown and harvested from STARTUP METXL 480 Seed Treatment treated seed is only for industrial uses and cannot be used for edible oil or any other human/feed consumption.*

DO NOT use Metalaxyl for control of *Pythium* in water-seeded rice.

After treated seed has been planted, DO NOT enter or allow worker entry into areas planted with treated seeds during the restricted-entry interval (REI) of 24 hours. EXCEPTION: Workers may enter these areas without restriction if there will be no worker contact with the soil/media subsurface.

DO NOT store leftover treated seed where there is a chance of it becoming mixed with untreated seed.

Seed treated with this product must be visually identifiable from untreated seed by the use of an approved colorant or dye to prevent accidental use of treated seed as food for humans or feed for animals. Refer to 21 CFR, Part 2.25. Any colorant or dye added to treated seed must be cleared for use in accordance with 40 CFR, Part 153.155(c).

DOMESTIC USE

CROP	DISEASE	RATE (Fluid Ounces per 100 Pounds of Seed)	
Brassica Leafy Vegetables Broccoli (Chinese and raab), Brussels sprout, Cabbage (Chinese bok choy, Chinese napa and mustard), Cauliflower, Cavalo broccoli, Collards, Kale, Kohlrabi, Mizuna, Mustard greens, Mustard Spinach, and Rape greens)	<i>Pythium</i> damping-off control	0.50	
Beets, Garden & Sugarbeets	<i>Pythium</i> damping-off control	0.50	
Canola* Mustard Seed* Rapeseed*	<i>Pythium</i> damping-off control	0.20 - 0.30	
	Restriction: DO NOT harvest mustard greens. Rape greens grown and harvested from STARTUP METXL 480 treated seed must not be used for human and feed consumption. Rapeseed grown and harvested from STARTUP METXL 480 treated seed is only for industrial uses and cannot be used for edible oil or any other human/feed consumption.		
Carrots*	<i>Pythium</i> damping-off control	0.50	
Cotton	<i>Pythium</i> seed rot and damping-off control	0.50 - 1.00	
Cucurbit Vegetables Chayote (fruit), Chinese waxgourd (Chinese preserving melon), Citron melon, Cucumber, Gherkin, Gourd (edible) (hyotan, cucuzza, hechima, Chinese okra), <i>Momordica</i> spp. (balsam apple, balsam pear, bittermelon, Chinese cucumber), Muskmelon (hybrids and/or varieties of <i>Cucumis melo</i> (true cantaloupe, cantaloupe, casaba, Santa Claus melon, crenshaw melon, honeydew melon, honey balls, Persian melon, golden pershaw melon, mango melon, pineapple melon, snake melon), Watermelon (hybrids and/or varieties of <i>Citrullus</i> spp.), Pumpkin, Squash (summer, winter including butternut squash, calabaza, hubbard squash, acorn squash, spaghetti squash).	<i>Pythium</i> damping-off control	0.50	
Forage Grasses Grasses grown for hay, grazing or silage, Corn (fodder or silage), Sorghum (hay or silage), and Small grain grown for hay, grazing or silage.	<i>Pythium</i> damping-off control	0.50	
Forage Legumes Alfalfa, Beans (for forage), Clover, Cowpeas (for forage), Cowpea hay, Lespedeza, Peanuts (for forage), Peanut hay, Peas (for forage), Pea vine hay, Soybeans (for forage), Soybean hay, Trefoil, Velvet beans (for forage) and Vetch.	<i>Pythium</i> damping-off and early season <i>Phytophthora</i> control	0.50 - 1.00	
Fruiting Vegetables Eggplant, Ground cherry (<i>Physalis</i> spp.), Pepino, Pepper (bell pepper, chili pepper, cooking pepper, pimento, sweet pepper), Tomatillo, Tomato.	<i>Pythium</i> damping-off control	0.50	
Grain Crops Barley, Buckwheat, Corn (field corn, sweet corn and popcorn), Millet, Milo, Oats, Rye, Triticale, and Wheat.	<i>Pythium</i> damping-off control	0.50	
	Systemic downy mildew control of sweet corn	1.00	
Use Rate Conversions for Field Corn			
PRODUCT FLUID OUNCES/			
100 lb seed†	1000 seeds	80,000 seed count	MG AI/SEED
0.50	0.0028	0.224	
1.00	0.0056	0.448	0.080
This product contains 14,195.5 mg metalaxyl (AI)/fluid ounce. †Based on standard number of 1786 seeds/lb (100,000 seeds/56 lb bu). NOTE: To maintain mg ai/seed, adjust fluid ounces/100 lb in proportion to the actual seeds/lb compared to the standard number of seeds/lb for that crop. For example if there are 1600 corn seeds/lb and the targeted rate is 0.040 mg ai/seed at a non-adjusted rate of 0.50 fl oz/100 lb of seed, use: (1600 ÷ 1786) x 0.50 = 0.45 fl oz/100 lb of seed.			
Leafy Vegetables Amaranth (Chinese spinach), Cardoon, Celery, Celery (Chinese), Celtuce, Chervil, Chrysanthemum (edible-leaved, garland), Corn salad, Cress (garden, upland), Dandelion, Dock (sorrel), Endive (escarole), Fennel (Florence), Lettuce (head and leaf), Orach, Parsley, Purslane (garden, winter), Radicchio (red chicory), Rhubarb, Spinach (including New Zealand and vine), Swiss chard.	<i>Pythium</i> damping-off control	0.50	
	Restriction: DO NOT apply this product to spinach which is destined to be grown in the greenhouse. Containers or bags containing STARTUP METXL 480 treated spinach seed must be labeled with this restriction.		

(continued)

DOMESTIC USE (continued)

CROP	DISEASE	RATE (Fluid Ounces per 100 Pounds of Seed)	
Legume Vegetables (except soybeans) Bean (<i>Lupinus</i> spp.) [grain lupin, sweet lupin, white lupin, white sweet lupin]; Bean (<i>Phaseolus</i> spp.) [field bean, kidney bean, lima bean (succulent or dry), navy bean, pinto bean, runner bean, snap bean, tepary bean, wax bean]; Bean (<i>Vigna</i> spp.) [adzuki bean, asparagus bean, blackeyed pea (succulent or dry), catjang, Chinese longbean, cowpea (succulent or dry), Crowder pea, moth bean (succulent or dry), mung bean, rice bean, southern pea (succulent or dry), urd bean, yardlong bean]; Broad bean (fava) (succulent or dry); Chickpea (garbanzo bean); Guar; Jackbean; Lablab bean (hyacinth); Lentil; Pea (<i>Pisum</i> spp.) [dwarf pea, edible-pod pea, English pea, field pea, garden pea, green pea, snow pea, sugar snap pea]; Pigeon pea (succulent or dry); Sword bean.	<i>Pythium</i> damping-off control	0.50	
	Early season <i>Phytophthora</i> control	0.50	
	Control of <i>Pythium</i> damping-off and systemic downy mildew of peas	1.00	
Onions (dry bulb and green)	<i>Pythium</i> damping-off control	0.50	
Peanuts*	<i>Pythium</i> seed rot and damping-off control	0.50	
Rice*	<i>Pythium</i> damping-off control	0.50 - 1.00	
	Restriction: If using STARTUP METXL 480 treated rice do not use Metalaxyl or mefenoxam for control of <i>Pythium</i> in water-seeded rice. Containers or bags containing STARTUP METXL 480 treated rice seed must be labeled with this restriction.		
Root and Tuber Vegetables (except potato) Arracacha, Arrowroot, Artichoke (Chinese, Jerusalem), Beet (garden, sugar), Burdock (edible), Canna (edible), Carrot, Cassava (bitter and sweet), Celeriac, Chayote (root), Chervil (turnip-rooted), Chicory, Chufa, Dasheen (taro), Ginger, Ginseng, Horseradish, Leren, Parsley (turnip-rooted), Parsnip, Radish (including oriental), Rutabaga, Salsify (black and Spanish), Skirret, Sweet potato, Tanier, Turmeric, Turnip, Yam bean, Yam (true).	<i>Pythium</i> damping-off control	0.50	
Sorghum	<i>Pythium</i> damping-off control	0.25 - 0.50	
	Systemic downy mildew (races susceptible to chemical control)	0.50 - 2.00	
	Apply 0.50 - 1.00 fluid ounces of this product on cultivars which possess sorghum downy mildew resistance. To those cultivars which do not possess resistance to a particular race of sorghum downy mildew in heavily infested areas with recent severe history, apply 1.00 - 2.00 fluid ounces of this product. Only high quality sorghum seed (90% germination and above) should be treated with this product.		
Soybeans	<i>Pythium</i> damping-off control and early season <i>Phytophthora</i> control	0.50 - 1.00	
Use Rate Conversions for Soybeans			
PRODUCT FLUID OUNCES/			
100 lb seed¹	1000 seeds	80,000 seed count	MG AI/SEED
0.50	0.0017	0.136	
1.00	0.0034	0.272	
This product contains 14,195.5 mg metalaxyl (AI)/fluid ounce. ¹ Based on standard number of 3000 seeds/lb (150,000 seeds/50 lb bu). NOTE: To maintain mg ai/seed, adjust fluid ounces/100 lb in proportion to the actual seeds/lb compared to the standard number of seeds/lb for that crop. For example if there are 2800 soybean seeds/lb and the targeted rate is 0.024 mg ai/seed at a non adjusted rate of 0.50 fl oz/100 lb of seed, use: (2800 ÷ 3000) x 0.50 = 0.467 fl oz/100 lb of seed.			
Spinach*	<i>Pythium</i> damping-off control and early season <i>Phytophthora</i> control	0.50 - 1.00	
	Restriction: DO NOT apply this product to spinach which is destined to be grown in the greenhouse. Containers or bags containing STARTUP METXL 480 treated spinach seed must be labeled with this restriction.		
Sunflowers	Systemic downy mildew control	1.00 - 2.00	
Turf Grasses Golf Courses, Home Lawns, Reclamation and Erosion Control Projects	<i>Pythium</i> damping-off control	1.00	

*Not registered for use in California

FOR EXPORT USE ONLY

For treatment of the following seed for export to countries requiring treatment with Metalaxyl before importation.

CROP	DISEASE	RATE (Fluid Ounces per 100 Pounds of Seed)
Brassica (Cole) Leafy Vegetables	<i>Pythium</i> damping-off control	0.50 - 1.00
Canola	<i>Pythium</i> damping-off control	0.30 - 1.10 (20 - 66 ml per 100 Kg of seed)
Corn	<i>Pythium</i> damping-off control	0.50 - 1.00
	Systemic downy mildew control	2.00
Cucurbit Vegetables	<i>Pythium</i> damping-off control	0.50
Fruiting Vegetables	<i>Pythium</i> damping-off control	0.50 - 1.00
Leafy Vegetables	<i>Pythium</i> damping-off control	0.50 - 1.00
Millet	<i>Pythium</i> damping-off control	0.50 - 1.00
	Systemic downy mildew control	1.70
Onions	<i>Pythium</i> damping-off control	0.50 - 1.00
Peas	<i>Pythium</i> damping-off control and early season <i>Phytophthora</i> control	2.30
Root and Tuber Vegetables (except potato)	<i>Pythium</i> damping-off control	0.50 - 1.00
	Restriction: DO NOT apply this product to potatoes.	
Sorghum	<i>Pythium</i> damping-off control	0.50 - 1.00
	Systemic downy mildew control	3.50
Sunflower	Systemic downy mildew control	6.00

Seed treated in accordance with the export use pattern must be labeled as follows:

FOR EXPORT ONLY

NOT FOR DOMESTIC SALES OR USE

DIRECTIONS FOR USE AT REDUCED RATES IN COMBINATION WITH OTHER FUNGICIDES

This product is a systemic fungicide seed dressing which may be applied in combination with other EPA-registered fungicides to aid in control of *Pythium*. This product may be applied as a water-based slurry through standard slurry of mist-type commercial seed treatment equipment.

CROP	DISEASE	RATE (Fluid Ounces per 100 Pounds of Seed)
Barley, Corn (field), Oats, Rye and Wheat	Aid in control of seed decay and damping-off caused by <i>Pythium</i>	0.07 - 0.25
Beans (field, kidney, navy and pinto), Lentils, Peas and Soybeans		
Peanuts*		
Rice*		
Sunflowers		

*Not registered for use in California

NOTE: Apply only in combination with EPA registered rates of broad-spectrum seed treatment fungicides.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store at temperatures above 40°F. Crystals may form at lower storage temperatures. If this occurs, place the product in a warm room (68°F or above) and roll or shake the container at frequent intervals until all crystals are dissolved.

PESTICIDE DISPOSAL: Open dumping is prohibited. Wastes resulting from the use of this product are acutely toxic. Improper disposal of unused pesticide, spray mixture or rinsate is a violation of Federal law. Pesticide, spray mixture or rinsate that cannot be used according to the label instruction must be disposed of according to Federal, state or local procedures. For guidance in proper disposal methods, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office.

CONTAINER DISPOSAL:

Nonrefillable Container: Do not reuse or refill this container. Triple rinse (or equivalent) promptly after emptying. Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Refillable Container: Refill this container with pesticide only. Do not reuse this container for any other purpose. Close all openings and replace all caps.

IMPORTANT INFORMATION READ BEFORE USING PRODUCT

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product reflect the opinion of experts based on field use and tests, and must be followed carefully. It is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of UPL NA Inc. or Seller. Handling, storage, and use of the product by Buyer or User are beyond the control of UPL NA Inc. and Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold UPL NA Inc. and Seller harmless for any claims relating to such factors.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, UPL NA INC. AND SELLER MAKE NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ON THIS LABEL.

To the extent consistent with applicable law, UPL NA Inc. or Seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product and **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF UPL NA INC. AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF UPL NA INC. OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

UPL NA Inc. and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by the duly authorized representative of UPL NA Inc.

UPL, the UPL logo and STARTUP are trademarks of a UPL Corporation Limited Group Company.

© 2020 UPL NA Inc. All rights reserved.

U-70506-356(041520-8704)