



| Active Ingredient | By Weight |
|-------------------------|-----------|
| Etoazole* | 72% |
| Other Ingredients | 28% |
| Total | 100% |

* 2-(2,6-difluorophenyl)-4-[4-(1,1-dimethylethyl)-2-ethoxyphenyl]-4,5-dihydrooxazole

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

| FIRST AID | |
|--|--|
| IF IN EYES: | <ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice. |
| IF ON SKIN OR CLOTHING: | <ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice. |
| IF SWALLOWED: | <ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to by a poison control center or doctor. • Do not give anything by mouth to an unconscious person. |
| IF INHALED: | <ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, Call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice. |
| EMERGENCY INFORMATION | |
| Have the product container or label with you when calling a poison control center or doctor, or going for treatment. FOR THE FOLLOWING EMERGENCIES, PHONE 24 HOURS A DAY: For Medical Emergencies phone:1-888-681-4261 For Transportation Emergencies, including spill, leak or fire, phone: CHEMTREC®1-800-424-9300 For Product Use Information phone: AMVAC®1-888-462-6822 | |

See inside booklet for First Aid, Precautionary Statements, and Directions for Use.

EPA Reg. No. 5481-650

EPA Est. No.: _____

Net Contents: _____



AMVAC Chemical Corporation
4695 MacArthur Court, Suite 1200
Newport Beach, CA 92660 U.S.A.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

Caution - Causes moderate eye irritation. Avoid contact with eyes, skin and clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear: long-sleeved shirt and long pants, shoes and socks, and chemical-resistant gloves made out of barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, natural rubber ≥ 14 mils, polyethylene, polyvinyl chloride ≥ 14 mils, or Viton® ≥ 14 mils.

Follow manufacturer's instructions for cleaning / maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to freshwater and marine/estuarine aquatic invertebrates, including oysters and shrimp. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwaters or rinsate.

PHYSICAL OR CHEMICAL HAZARDS

Do not mix or allow contact with oxidizing agents. Hazardous chemical reaction may occur.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is: long sleeved shirt and long pants, chemical-resistant gloves and shoes plus socks.

CHEMIGATION

RESTRICTION: Do not apply this product through any type of irrigation or chemigation system.

RESISTANCE MANAGEMENT RECOMMENDATIONS

This product is a Group 10B insecticide, and mite varieties that have developed resistance to Group 10B insecticides may eventually become dominant if Group 10B insecticides are repeatedly used as the primary control over successive years or in the same field. If this occurs, this product (as well as other Group 10B insecticides) may become partially or totally ineffective in controlling the resistance species.

To help prevent resistance to this product, users should:

Avoid using this product or other Group 10B insecticides with a similar mode of action in sequential applications to control the same mite species.

Use premixes or tank mixtures that include insecticides registered for this use that have different modes of action and are from different insecticide Groups.

Incorporate the use of this product into an Integrated Pest Management (IPM) program. Monitor application sites for reduced or failed efficacy.

To address site specific and resistant pest issues, contact your local crop advisor, extension specialist and/or the manufacturer for insecticide resistance management and/or IPM recommendations.

WATER-SOLUBLE PACKAGING INSTRUCTIONS

The packets within this bag are water-soluble and each contain 3 ounces of Stifle WP Miticide. Water Soluble Packages (WSP's) are designed to dissolve in water. Agitation may be used, if necessary, to help dissolve the WSP. Failure to follow handling and mixing instructions can increase your exposure to the pesticide products in WSP's. WSP's, when used properly, qualify as a closed mixing/loading system under the Agricultural Worker Protection standard [40 CFR 170.607(d)].

Handling Instructions

Follow these steps when handling pesticide products in WSPs

1. Mix in spray tank only
2. Handle WSP(s) in a manner that protects package from breakage and/or unintended release of contents. If package is broken, put on PPE required for clean-up and then continue with mixing instructions.
3. Keep the WSP(s) in outer packaging until just before use.
4. Keep the WSP dry prior to adding to the spray tank.
5. Handle with dry gloves and according to the label instructions for PPE.
6. Keep WSP intact. Do not cut or puncture WSP.
7. Reseal the WSP outer package to protect any unused WSP(s)

Mixing Instructions

To prepare a spray solution of Stifle WP Miticide, refer to the following table to determine the number of packets of Stifle WP Miticide to add to the spray tank:

| Application Rate (ounces/acre) | Acres Treated per 3 oz. Packet |
|-----------------------------------|-----------------------------------|
| 0.66 | 4.5 |
| 1.0 | 3.0 |
| 2.0 | 1.5 |
| 3.0 | 1.0 |
| 4.0 | 0.75 |

Follow the steps below when mixing this product, including if tank mixed with other pesticide products. If being tank mixed, the mixing directions 1 through 9 below take precedence over the mixing directions of the other tank mix products. WSPs may, in some cases, be mixed with other pesticide products so long as the directions for use of all mixed products do not conflict. Do not tank mix this product with products that prohibit tank mixing or have conflicting mixing directions.

1. If a basket or strainer is present in the tank hatch, remove prior to adding the WSP to the tank.
2. Fill tank with water to approximately one-third to one-half of the desired final volume of spray.
3. Stop adding water and stop any agitation.

4. Place intact/unopened WSP(s) into the tank.
5. Do not spray water from a hose or fill pipe to break or dissolve the WSP(s).
6. Start mechanical and recirculation agitation from the bottom of tank without using any overhead recirculation, if possible. If overhead recirculation cannot be turned off, close the hatch before starting agitation.
7. Dissolving the WSP(s) may take up to 5 minutes or longer, depending on water temperature, water hardness and intensity of agitation.
8. Stop agitation before tank lid is opened.
9. Open the lid to the tank, exercising caution to avoid contact with dusts or spray mix, to verify that the WSPs have fully dissolved and the contents have been thoroughly mixed into the solution.
10. Do not add other allowed products or complete filling the tank until the bags have fully dissolved and pesticide is thoroughly mixed.
11. Once the WSP have fully dissolved and any other products have been added to the tank, resume filling the tank with water to the desired level, close the tank lid, and resume agitation.
12. Use the spray solution when mixing is complete.
13. Maintain agitation of the diluted pesticide mix during transport and application.
14. It is unlawful to use any registered pesticide, including WSPs, in a manner inconsistent with its labeling.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

CROP SPECIFIC INSTRUCTIONS

| CANE BERRY (Subgroup 13-07A): Blackberry; Loganberry; Raspberry (Black, Red and Wild) and cultivars, varieties and hybrids of these crops. | | |
|---|--|---|
| PEST | APPLICATION RATE (Oz./A) | INSTRUCTIONS |
| European Red Spider Mite McDaniel Spider Mite Two-spotted Spider Mite Yellow Spider Mite | 2.0 – 3.0 0.090 – 0.135 pounds AI/A) | Apply using a full-coverage ground spray using sufficient water to ensure complete coverage of the entire plant (minimum 50 gallons / acre), including the reproductive parts of the plant that are targeted by mites. Choose the lower rate for light infestations and the higher rate for heavy infestations. Larger amounts of water should be used when treating older trees or plants that have dense and / or compact foliage. Best coverage is achieved when larger volumes of water are used in combination with properly calibrated spray equipment. Because Stifle WP Miticide is primarily a larvicide / ovicide, for best results apply when the mite pressure is still low. Be sure to read and follow the RESISTANCE MANAGEMENT RECOMMENDATIONS information at the beginning of this label. |
| RESTRICTIONS <ul style="list-style-type: none"> • Preharvest Interval (PHI): 0 days. • Do NOT make more than one (1) application of this product per calendar year. • Do NOT apply more than 3.0 ounces (0.135 pounds AI) of this product per acre per calendar year. • Do NOT apply less than 2.0 ounces / acre (0.09 pounds AI/A). Applications at a rate lower than 2.0 ounces / acre may result in poor control and potential development of mite resistance to the active ingredient in this product (etoxazole). | | |

| FIELD CORN, POPCORN, CORN GROWN FOR SEED | | |
|--|---|--|
| PEST | APPLICATION RATE (Oz./A) | INSTRUCTIONS |
| Banks Grass Mite ¹ Carmine Spider Mite Pacific Spider Mite Strawberry Spider Mite Two-spotted Spider Mite | 1.0 - 3.0 (0.045 - 0.135 pounds AI/A) | Complete coverage of the plants is essential to achieve good control, and applications should be made prior to tasseling to help with complete coverage. Because Stifle WP Miticide is primarily a larvicide / ovicide, for best results apply when the mite pressure is still low. Choose the lower rate for light infestations and the higher rate for heavy infestations. Aerial Applications ² : Apply using a minimum of 3 gallons of water per acre. Ground Applications: Apply using a minimum of 10 gallons of water per acre. When applying to Field Corn Grown for Seed, application of this product should be made at or before tasseling. If a second treatment is necessary, apply 14 days after the first application. Be sure to read and follow the RESISTANCE MANAGEMENT RECOMMENDATIONS information at the beginning of this label. |
| RESTRICTIONS <ul style="list-style-type: none"> • Preharvest Interval (PHI): 21 days. • Do NOT make more than two (2) applications of this product per calendar year. • Do NOT make an additional treatment within 14 days of the previous treatment. • Do NOT apply more than 6.0 ounces (0.27 pounds AI) of this product per acre per calendar year. • Do NOT apply less than 1.0 ounces / acre (0.045 pounds AI/A). Applications at a rate lower than 1.0 ounces / acre may result in poor control and potential development of mite resistance to the active ingredient in this product (etoxazole). | | |
| NOTES <ol style="list-style-type: none"> 1) When applying to control only Banks Grass Mite, use of the upper range of the rates listed (1.0 - 3.0 oz. / A) may be necessary, especially if the pest pressure is severe. In cases of severe Banks Grass Mite pest pressure in conjunction with extreme drought conditions, more than one miticide application may be necessary. 2) NOT APPROVED FOR AERIAL APPLICATION IN NEW YORK. | | |

| COTTON | | |
|---|--|--|
| PEST | APPLICATION RATE (Oz./A) | INSTRUCTIONS |
| Carmine Spider Mite Pacific Spider Mite Two-spotted Spider Mite | 0.66 - 1.0 (0.030 - 0.045 pounds AI/A) | Complete coverage of the plants is essential to achieve good control. Because Stifle WP Miticide is primarily a larvicide / ovicide, for best results apply when the mite pressure is still low. Choose the lower rate for light infestations and the higher rate for heavy infestations. Aerial Applications: Apply using a minimum of 3 gallons of water per acre. Ground Applications: Apply using a minimum of 10 gallons of water per acre. Be sure to read and follow the RESISTANCE MANAGEMENT RECOMMENDATIONS information at the beginning of this label. |
| RESTRICTIONS <ul style="list-style-type: none"> • Preharvest Interval (PHI): 28 days. • Do NOT make more than one (1) application of this product per calendar year. • Do NOT apply more than 1.0 ounce (0.045 pounds AI) of this product per acre per calendar year. Do NOT apply less than 0.66 ounces / acre (0.030 pounds AI/A). Applications at a rate lower than 0.66 ounces / acre may result in poor control and potential development of mite resistance to the active ingredient in this product (etoxazole). | | |

CUCURBIT VEGETABLES (Crop Group 9): Acorn Squash; Balsam Apple; Balsam Pear; Bittermelon; Butternut Squash; Calabaza Squash; Cantaloupe; Chayote (fruit); Chinese Cucumber; Chinese Okra; Chinese Waxgourd (Chinese Preserving Melon); Citron Melon; Cucumber; Cucuzza; Gherkin; Gourd, Edible; Hechima; Hubbard Squash; Hyotan; *Momordica* spp.; Muskmelon; Pumpkin; Spaghetti Squash; Summer Squash, Watermelon; Winter Squash

| PEST | APPLICATION RATE (Oz./A) | INSTRUCTIONS |
|---|--|---|
| Carmine Spider Mite Pacific Spider Mite Strawberry Spider Mite Two-spotted Spider Mite | 2.0 - 3.0 (0.090 - 0.135 pounds AI/A) | <p>Complete coverage of the plants is essential to achieve good control, including the reproductive parts of the plant that are targeted by mites. Choose the lower rate for light infestations and the higher rate for heavy infestations. Larger amounts of water should be used when treating mature plants or plants that have dense and / or compact foliage. Best coverage is achieved when larger volumes of water are used in combination with properly calibrated spray equipment. Because Stifle WP Miticide is primarily a larvicide / ovidicide, for best results apply when the mite pressure is still low.</p> <p>Aerial Applications¹: Apply using a minimum of 3 gallons of water per acre.</p> <p>Ground Applications: Apply using a minimum of 10 gallons of water per acre.</p> <p>Be sure to read and follow the RESISTANCE MANAGEMENT RECOMMENDATIONS information at the beginning of this label.</p> |

- RESTRICTIONS**
- Preharvest Interval (PHI): 7 days.
 - Do NOT make more than one (1) application of this product per calendar year.
 - Do NOT apply more than 3.0 ounces (0.135 pounds AI) of this product per acre per calendar year.
 - Do NOT apply less than 2.0 ounces / acre (0.090 pounds AI/A). Applications at a rate lower than 2.0 ounces / acre may result in poor control and potential development of mite resistance to the active ingredient in this product (etoxazole).

NOTES
 1) NOT APPROVED FOR AERIAL APPLICATION IN NEW YORK.

HOPS

| PEST | APPLICATION RATE (Oz./A) | INSTRUCTIONS |
|-------------------------|--|--|
| Two-spotted Spider Mite | 3.0 - 4.0 (0.135 - 0.180 pounds AI/A) | <p>Apply using a full-coverage ground spray using sufficient water to ensure complete coverage of the entire plant (minimum 50 gallons / acre). Choose the lower rate for light infestations and the higher rate for heavy infestations. Best coverage is achieved when larger volumes of water are used in combination with properly calibrated spray equipment. Because Stifle WP Miticide is primarily a larvicide / ovidicide, for best results apply when the mite pressure is still low.</p> <p>Be sure to read and follow the RESISTANCE MANAGEMENT RECOMMENDATIONS information at the beginning of this label.</p> |

- RESTRICTIONS**
- Preharvest Interval (PHI): 7 days.
 - Do NOT make more than one (1) application of this product per calendar year.
 - Do NOT apply more than 4.0 ounces (0.180 pounds AI) of this product per acre per calendar year.
 - Do NOT apply less than 3.0 ounces / acre (0.135 pounds AI/A). Applications at a rate lower than 3.0 ounces / acre may result in poor control and potential development of mite resistance to the active ingredient in this product (etoxazole).

LOW GROWING BERRY (Subgroup 13-07G): Bearberry; Bilberry; Blueberry, Lowbush; Cloudberry; Cranberry; Lingonberry; Muntries; Partridgeberry; Strawberry; and cultivars, varieties and/or hybrids of these crops.

| PEST | APPLICATION RATE (Oz./A) | INSTRUCTIONS |
|---|---|---|
| European Red Mite Pacific Spider Mite Two-spotted Spider Mite | 2.0 - 3.0 (0.090 - 0.135 pounds AI/A) | Apply using a full-coverage ground spray using airblast equipment and sufficient water to ensure complete coverage of the entire plant (minimum 100 gallons / acre). Choose the lower rate for light infestations and the higher rate for heavy infestations. Best coverage is achieved when larger volumes of water are used in combination with properly calibrated spray equipment. Because Stifle WP Miticide is primarily a larvicide / ovicide, for best results apply when the mite pressure is still low. NOTE: This product does not control Cyclamine Mite. If this mite species is present, a miticide registered for use to control Cyclamine Mite should be used. Be sure to read and follow the RESISTANCE MANAGEMENT RECOMMENDATIONS information at the beginning of this label. |
| Two-spotted Spider Mite | 2.0 - 3.0 (0.090 - 0.135 pounds AI/A) + Pyrethroid insecticide/Miticide (such as a product containing Fenpropathrin) - use at Labeled Rate | Apply using a full-coverage ground spray using sufficient water to ensure complete coverage of the entire plant (minimum 100 gallons / acre). Choose the lower rate for light infestations and the higher rate for heavy infestations. Best coverage is achieved when larger volumes of water are used in combination with properly calibrated spray equipment. If retreatment is necessary within 30 days of last treatment to comply with local IPM programs, treatments must be alternated with other non-pyrethroid insecticides. Be sure to follow all applicable directions, precautions and restrictions in the registered label for the pyrethroid tank mix partner Be sure to read and follow the RESISTANCE MANAGEMENT RECOMMENDATIONS information at the beginning of this label. |
| RESTRICTIONS <ul style="list-style-type: none"> • Preharvest Interval (PHI): 1 day. • Do NOT make more than one (1) application of this product per calendar year. • Do NOT apply more than 3.0 ounces (0.135 pounds AI) of this product per acre per calendar year. • Do NOT apply less than 2.0 ounces / acre (0.090 pounds AI/A). Applications at a rate lower than 3.0 ounces / acre may result in poor control and potential development of mite resistance to the active ingredient in this product (etoxazole). | | |

| MINT: Peppermint and Spearmint | | |
|--|--|--|
| PEST | APPLICATION RATE (Oz./A) | INSTRUCTIONS |
| Pacific Spider Mite Strawberry Spider Mite Two-spotted Spider Mite | 2.0 – 4.0 (0.090 – 0.180 pounds AI/A) | Complete coverage of the plants is essential to achieve good control. Best coverage is achieved when larger volumes of water are used in combination with properly calibrated spray equipment. Because Stifle WP Miticide is primarily a larvicide / ovicide, for best results apply when the mite pressure is still low. Choose the lower rate for light infestations and the higher rate for heavy infestations. Aerial Applications ¹ : Apply using a minimum of 10 gallons of water per acre. Ground Applications: Apply using a minimum of 50 gallons of water per acre. Be sure to read and follow the RESISTANCE MANAGEMENT RECOMMENDATIONS information at the beginning of this label. |
| RESTRICTIONS <ul style="list-style-type: none"> • Preharvest Interval (PHI): 7 days. • Do NOT make more than one (1) application of this product per calendar year. • Do NOT apply more than 4.0 ounces (0.180 pounds AI) of this product per acre per calendar year. • Do NOT apply less than 2.0 ounces / acre (0.090 pounds AI/A). Applications at a rate lower than 2.0 ounces / acre may result in poor control and potential development of mite resistance to the active ingredient in this product (etoxazole). | | |
| NOTES 1) NOT APPROVED FOR AERIAL APPLICATION IN NEW YORK. | | |

| PEPPER AND EGGPLANT (Subgroup 8-10B): African Eggplant; Bell Pepper; Eggplant; Martynia; Non-Bell Pepper; Okra; Pea Eggplant; Pepino; Roselle; Scarlet Eggplant and cultivars, varieties and/or hybrids of these crops. | | |
|--|--|--|
| PEST | APPLICATION RATE (Oz./A) | INSTRUCTIONS |
| Two-spotted Spider Mite | 2.0 – 3.0 (0.090 – 0.135 pounds AI/A) | Apply using a full-coverage ground spray using sufficient water to ensure complete coverage of the entire plant (minimum 20 gallons / acre), including the reproductive parts of the plant that are targeted by mites. Choose the lower rate for light infestations and the higher rate for heavy infestations. Larger amounts of water should be used when treating mature plants or varieties that have dense and / or compact foliage. Best coverage is achieved when larger volumes of water are used in combination with properly calibrated spray equipment. Because Stifle WP Miticide is primarily a larvicide / ovicide, for best results apply when the mite pressure is still low. Be sure to read and follow the RESISTANCE MANAGEMENT RECOMMENDATIONS information at the beginning of this label. |
| RESTRICTIONS <ul style="list-style-type: none"> • Preharvest Interval (PHI): 7 days. • Do NOT make more than one (1) application of this product per calendar year. • Do NOT apply more than 3.0 ounces (0.135 pounds AI) of this product per acre per calendar year. • Do NOT apply less than 2.0 ounces / acre (0.090 pounds AI/A). Applications at a rate lower than 2.0 ounces / acre may result in poor control and potential development of mite resistance to the active ingredient in this product (etoxazole). | | |

| POME FRUIT (Crop Group 11): Apple; Crabapple; Loquat; Mayhaw; Pear; Pear, Asian; Quince | | |
|---|--|---|
| PEST | APPLICATION RATE (Oz./A) | INSTRUCTIONS |
| European Red Mite McDaniel Spider Mite Pacific Spider Mite Two-spotted Spider Mite | 2.0 – 3.0 (0.090 – 0.135 pounds AI/A) | Apply using a full-coverage ground spray using airblast equipment and sufficient water to ensure complete coverage of the entire plant (minimum 100 gallons / acre). Choose the lower rate for light infestations and the higher rate for heavy infestations. Best coverage is achieved when larger volumes of water are used in combination with properly calibrated spray equipment. Because Stifle WP Miticide is primarily a larvicide / ovicide, for best results apply when the mite pressure is still low. NOTE: This product does not control Blister Mites or Rust Mites. If these mite species are present, a miticide registered for use to control these pests should be used. Be sure to read and follow the RESISTANCE MANAGEMENT RECOMMENDATIONS information at the beginning of this label. |
| RESTRICTIONS | | |
| <ul style="list-style-type: none"> • Preharvest Interval (PHI): 14 days. • Do NOT make more than one (1) application of this product per calendar year. • Do NOT apply more than 3.0 ounces (0.135 pounds AI) of this product per acre per calendar year. • Do NOT apply less than 2.0 ounces / acre (0.090 pounds AI/A). Applications at a rate lower than 2.0 ounces / acre may result in poor control and potential development of mite resistance to the active ingredient in this product (etoxazole). | | |

| SMALL FRUIT VINE CLIMBING, EXCEPT FUZZY KIWIFRUIT (Subgroup 13-07F): Amur River Grape; Gooseberry; Grape; Kiwifruit, hardy; Maypop; Schisandra Berry and cultivars, varieties and hybrids of these crops. | | |
|--|--|--|
| PEST | APPLICATION RATE (Oz./A) | INSTRUCTIONS |
| European Red Spider Mite Pacific Spider Mite Two-spotted Spider Mite Willamette Mite | 2.0 – 3.0 (0.090 – 0.135 pounds AI/A) | Apply using a full-coverage ground spray using sufficient water to ensure complete coverage of the entire plant (minimum 25 gallons / acre). Choose the lower rate for light infestations and the higher rate for heavy infestations. Larger amounts of water should be used when treating older trees or plants that have dense and / or compact foliage. Best coverage is achieved when larger volumes of water are used in combination with properly calibrated spray equipment. Because Stifle WP Miticide is primarily a larvicide / ovicide, for best results apply when the mite pressure is still low. NOTE: This product does not control Blister Mites or Rust Mites. If these mite species are present, a miticide registered for use to control these pests should be used. Be sure to read and follow the RESISTANCE MANAGEMENT RECOMMENDATIONS information at the beginning of this label. |
| RESTRICTIONS | | |
| <ul style="list-style-type: none"> • Preharvest Interval (PHI): 14 days. • Do NOT make more than one (1) application of this product per calendar year. • Do NOT apply more than 3.0 ounces (0.135 pounds AI) of this product per acre per calendar year. • Do NOT apply less than 2.0 ounces / acre (0.09 pounds AI/A). Applications at a rate lower than 2.0 ounces / acre may result in poor control and potential development of mite resistance to the active ingredient in this product (etoxazole). | | |

| STONE FRUIT (Crop Group 12): Apricot; Cherry (sweet and tart); Nectarine; Peach; Plum; Plum, Chicksaw; Plum, Damson; Plum, Japanese; Plumcot; Prune (fresh) | | |
|---|--|---|
| PEST | APPLICATION RATE (Oz./A) | INSTRUCTIONS |
| European Red Spider Mite Pacific Spider Mite Two-spotted Spider Mite | 2.0 - 3.0 (0.090 - 0.135 pounds AI/A) | Apply using a full-coverage ground spray using sufficient water to ensure complete coverage of the entire plant (minimum 50 gallons / acre). Choose the lower rate for light infestations and the higher rate for heavy infestations. Larger amounts of water should be used when treating older trees or plants that have dense and / or compact foliage. Best coverage is achieved when larger volumes of water are used in combination with properly calibrated spray equipment. Because Stifle WP Miticide is primarily a larvicide / ovicide, for best results apply when the mite pressure is still low. Be sure to read and follow the RESISTANCE MANAGEMENT RECOMMENDATIONS information at the beginning of this label. |
| RESTRICTIONS | | |
| <ul style="list-style-type: none"> • Preharvest Interval (PHI): 7 days. • Do NOT make more than one (1) application of this product per calendar year. • Do NOT apply more than 3.0 ounces (0.135 pounds AI) of this product per acre per calendar year. • Do NOT apply less than 2.0 ounces / acre (0.09 pounds AI/A). Applications at a rate lower than 2.0 ounces / acre may result in poor control and potential development of mite resistance to the active ingredient in this product (etoxazole). | | |

| NON-BEARING FRUIT TREES | | |
|--|--|---|
| PEST | APPLICATION RATE (Oz./A) | INSTRUCTIONS |
| European Red Mite McDaniel Spider Mite Pacific Spider Mite Two-spotted Spider Mite | 2.0 - 3.0 (0.090 - 0.135 pounds AI/A) | Apply using a full-coverage ground spray using airblast equipment and sufficient water to ensure complete coverage of the entire plant. Best coverage is achieved when larger volumes of water are used in combination with properly calibrated spray equipment. Choose the lower rate for light infestations and the higher rate for heavy infestations. Because Stifle WP Miticide is primarily a larvicide / ovicide, for best results apply when the mite pressure is still low. Be sure to read and follow the RESISTANCE MANAGEMENT RECOMMENDATIONS information at the beginning of this label. |
| RESTRICTIONS | | |
| <ul style="list-style-type: none"> • Do NOT harvest nuts or fruit from trees that have been treated within one (1) year of application. • Do NOT make more than one (1) application of this product per calendar year. • Do NOT apply more than 3.0 ounces (0.135 pounds AI) of this product per acre per calendar year. • Do NOT apply less than 2.0 ounces / acre (0.090 pounds AI/A). Applications at a rate lower than 2.0 ounces / acre may result in poor control and potential development of mite resistance to the active ingredient in this product (etoxazole). | | |

| CHRISTMAS TREES [(Not For Use in the State of California)] | | |
|---|---|---|
| PEST | APPLICATION RATE (Oz./A) | INSTRUCTIONS |
| McDaniel Spider Mite Pacific Spider Mite Spruce Spider Mite Two-spotted Spider Mite | 2.0 – 3.0 (0.090 – 0.135 pounds AI/A) | Complete coverage of the plants is essential to achieve good control. Best coverage is achieved when larger volumes of water are used in combination with properly calibrated spray equipment. Because Stifle WP Miticide is primarily a larvicide / ovicide, for best results apply when the mite pressure is still low. Choose the lower rate for light infestations and the higher rate for heavy infestations. Aerial Applications: Apply using a minimum of 10 gallons of water per acre. Ground Applications: Apply as a full coverage spray using a minimum of 50 gallons of water per acre. Be sure to read and follow the RESISTANCE MANAGEMENT RECOMMENDATIONS information at the beginning of this label. |
| RESTRICTIONS | | |
| <ul style="list-style-type: none"> • [Not for use on Christmas Trees in the State of California] • Do NOT make more than one (1) application of this product per calendar year. • Do NOT apply more than 3.0 ounces (0.135 pounds AI) of this product per acre per calendar year. • Do NOT apply less than 2.0 ounces / acre (0.090 pounds AI/A). Applications at a rate lower than 2.0 ounces / acre may result in poor control and potential development of mite resistance to the active ingredient in this product (etoxazole). | | |

| TREE NUTS: Almond; Beech Nut; Brazil Nut; Butternut; Cashew; Chestnut; Chinquapin; Filbert (Hazelnut); Hickory Nut; Macadamia Nut; Pecan; Pistachio; Walnut, Black and English | | |
|---|---|---|
| PEST | APPLICATION RATE (Oz./A) | INSTRUCTIONS |
| European Red Mite Pacific Spider Mite Two-spotted Spider Mite | 2.0 – 3.0 (0.090 – 0.135 pounds AI/A) | Apply using a full-coverage ground spray using airblast equipment and sufficient water to ensure complete coverage of the entire plant. Choose the lower rate for light infestations and the higher rate for heavy infestations. Best coverage is achieved when larger volumes of water are used in combination with properly calibrated spray equipment. Because Stifle WP Miticide is primarily a larvicide / ovicide, for best results apply when the mite pressure is still low. Be sure to read and follow the RESISTANCE MANAGEMENT RECOMMENDATIONS information at the beginning of this label. |
| RESTRICTIONS | | |
| <ul style="list-style-type: none"> • Preharvest Interval (PHI): 28 days. • Do NOT make more than one (1) application of this product per calendar year. • Do NOT apply more than 3.0 ounces (0.135 pounds AI) of this product per acre per calendar year. • Do NOT apply less than 2.0 ounces / acre (0.090 pounds AI/A). Applications at a rate lower than 2.0 ounces / acre may result in poor control and potential development of mite resistance to the active ingredient in this product (etoxazole). | | |

| TROPICAL AND SUB-TROPICAL FRUIT (INEDIBLE PEEL): Avocado; Canistel; Mango; Papaya; Sapodilla; Sapote, Black and Mamey; Star Apple | | |
|---|--|---|
| PEST | APPLICATION RATE (Oz./A) | INSTRUCTIONS |
| Avocado Brown Mite Persea Mite Two-spotted Spider Mite | 2.0 - 3.0 (0.090 - 0.135 pounds AI/A) | Complete coverage of the plants is essential to achieve good control, including the reproductive parts of the plant that are targeted by mites. Larger amounts of water should be used when treating mature plants or plants that have dense and / or compact foliage. Best coverage is achieved when larger volumes of water are used in combination with properly calibrated spray equipment. Because Stifle WP Miticide is primarily a larvicide / ovicide, for best results apply when the mite pressure is still low. Choose the lower rate for light infestations and the higher rate for heavy infestations. Aerial Applications ¹ : Apply using a minimum of 20 gallons of water per acre. Ground Applications: Apply using a minimum of 50 gallons of water per acre. Be sure to read and follow the RESISTANCE MANAGEMENT RECOMMENDATIONS information at the beginning of this label. |
| RESTRICTIONS | | |
| <ul style="list-style-type: none"> • Preharvest Interval (PHI): 1 day. • Do NOT make more than one (1) application of this product per calendar year. • Do NOT apply more than 3.0 ounces (0.135 pounds AI) of this product per acre per calendar year. • Do NOT apply less than 2.0 ounces / acre (0.090 pounds AI/A). Applications at a rate lower than 2.0 ounces / acre may result in poor control and potential development of mite resistance to the active ingredient in this product (etoxazole). | | |
| NOTES | | |
| 1) NOT APPROVED FOR AERIAL APPLICATION IN NEW YORK. | | |

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.

PESTICIDE STORAGE

Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers. Store in a cool, dry place. Do not store diluted spray. For help with any spill, leak, fire or exposure involving this material, call day or night (800) 424-9300.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Open dumping is prohibited.

CONTAINER HANDLING

Nonrefillable outer bag. Do not reuse or refill the outer bag. Completely empty the bag into application equipment, then dispose of empty bag in a sanitary landfill or by incineration.

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants (a) that this product conforms to the chemical description on the label; and (b) that the directions, warnings, and other statements on this label are based upon responsible experts' evaluations of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants and residues on food crops, and upon reports of field experience. Tests have not been made on all varieties of food crops and plants, or in all states or under all conditions. THIS WARRANTY DOES NOT EXTEND TO THE USE OF THIS PRODUCT CONTRARY TO LABEL INSTRUCTIONS, OR UNDER CONDITIONS NOT REASONABLY FORESEEABLE.

THERE ARE NO EXPRESS WARRANTIES OTHER THAN THOSE SET FORTH HEREIN. TO THE EXTENT CONSISTENT WITH

APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE, TO MAKE ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OF FITNESS FOR A PARTICULAR PURPOSE, OR ANY WARRANTY OF QUALITY OR PERFORMANCE. THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS OR CAUTIONS.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN TORT, CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

AMVAC offers this product, and Buyer accepts it, subject to the foregoing Limited Warranty which may be varied only by agreement in writing signed by an authorized representative of AMVAC.

© 2020 AMVAC Chemical Corporation is a wholly owned subsidiary of American Vanguard Corporation. All Rights Reserved. AMERICAN VANGUARD, AMVAC, and their respective logos are trademarks owned by AMVAC Chemical Corporation. CHEMTREC is a service mark of the American Chemistry Council, Inc. Viton is a trademark of the Chemours Company FC, LLC.

AMVAC Chemical Corporation
4695 MacArthur Court, Suite 1200
Newport Beach, CA 92660 U.S.A.

SPECIMEN