

ETOXAZOLE GROUP 10B INSECTICIDE

Suremite™ Miticide

Miticide for use on Caneberry, Christmas Trees, Cotton, Cucurbit Vegetables, Field Corn, Popcorn, Corn (Grown for Seed), Hops, Low Growing Berry, Mint, Non-Bearing Fruit Trees, Pepper and Eggplant, Pome Fruit, Small Fruit Vine Climbing (except Fuzzy Kiwifruit), Stone Fruit, Sweet Corn, Tree Nuts, and Tropical and Sub-Tropical Fruit (Inedible Peel).

ACTIVE INGREDIENT: etoxazole	72.0%
OTHER INGREDIENTS:	28.0%
TOTAL	100.0%

EPA Reg. No. 2749-592 EPA Est. No. 065387-AR-002

Read "LIMIT OF WARRANTY AND LIABILITY" before buying or using. If terms are not acceptable, return at once unopened.

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

<p>FIRST AID If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. If swallowed: Call poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.</p>	<p>If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance; then give artificial respiration, preferably by mouth to-mouth, if possible. Call a poison control center or doctor for further treatment advice. If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. Emergency Assistance: Have the product container or label with you when calling a poison control center or doctor or going for treatment. FOR EMERGENCY MEDICAL HELP, CALL CHEMTREC TOLL FREE 800-424-9300 or 1-703-527-3887.</p>
---	---

FOR CHEMICAL SPILL, LEAK, FIRE, EXPOSURE OR MEDICAL EMERGENCY INVOLVING THIS PRODUCT CALL CHEMTREC® TOLL FREE (800) 424-9300 OR (703) 527-3887.

12 oz (4x3 oz WSBs)
NET CONTENTS

Manufactured for: Aceto Life Sciences, L.L.C., 4 Tri Harbor Court, Port Washington, NY 11050



PRECAUTIONARY STATEMENTS**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION. Causes moderate eye irritation. Avoid contact with eyes or clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical resistant gloves made out of: barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, natural rubber ≥ 14 mils, polyethylene, polyvinyl chloride ≥ 14 mils, or viton ≥ 14 mils.
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exists, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS

Water soluble bags, when used correctly, qualify as a closed mixing/loading system under the Worker Protection Standard [40 CFR 170.607(d)]. Mixers and loaders handling this product while it is enclosed in intact water-soluble bags may elect to wear reduced PPE of long-sleeved shirt, long pants, shoes, socks, a chemical resistant apron, and chemical resistant gloves. When reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for "applicators and other handlers" and have such PPE immediately available for use in an emergency, such as a spill or equipment break-down.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. The wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product, Wash the outside of the gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to freshwater and marine/estuarine aquatic invertebrates, including oysters and shrimp. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high watermark. Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwaters or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that it will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Long-sleeved shirt and long pants
- Chemical resistant gloves made out of: barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, natural rubber ≥ 14 mils, polyethylene, polyvinyl chloride ≥ 14 mils, or viton ≥ 14 mils.
- Shoes plus socks

PRODUCT INFORMATION

This product contains etoxazole, a contact acaricide/ovicide and a member of the diphenyloxazoline class of insecticide/miticide. This product kills eggs and nymphs and prevent adults from laying viable eggs, it inhibits the molting process in juvenile mites, but does not kill mite adults.

Evidence of activity may be slower than typical contact insecticides as treated susceptible pests may remain alive on the plant for 3-7 days; however, pests have stopped feeding and any feeding damage during this time is typically very low.

USE RESTRICTIONS

- Chemigation for field corn, popcorn and corn (grown for seed production) only.

RESISTANCE MANAGEMENT

For resistance management, Suremite™ Miticide contains a Group 10B acaricide. Any mite population may contain individuals naturally resistant to Suremite™ Miticide and other Group 10B acaricides. The resistant individuals may dominate the mite population if this group of acaricides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay acaricide resistance, take the following steps:

- Rotate the use of Suremite™ Miticide or other Group 10B acaricides within a growing season, or among growing seasons, with different groups that control the same pests.
- Use tank mixtures with acaricides from a different group that are equally effective on the target pest when such use is permitted. Do not rely on the same mixture repeatedly for the same pest population. Consider any known cross-resistance issues (for the targeted pests) between the individual components of a mixture. In addition, consider the following recommendations provided by the Insecticide Resistance Action Committee (IRAC):
 - > Individual insecticides selected for use in mixtures should be highly effective and be applied at the rates at which they are individually registered for use against the target species.
 - > Mixtures with components having the same IRAC mode of action classification are not recommended for insect resistance management.

RESISTANCE MANAGEMENT *(continued)*

- > When using mixtures, consider any known cross-resistance issues between the individual components for the targeted pest(s).
- > Mixtures become less effective if resistance is already developing to one or both active ingredients, but they may still provide pest management benefits.
- > The insect resistance management benefits of an insecticide mixture are greatest if the two components have similar periods of residual insecticidal activity. Mixtures of insecticides with unequal periods of residual insecticide activity may offer an insect resistance management benefit only for the period where both insecticides are active.
- Adopt an integrated pest management program for acaricides use that includes scouting, uses historical information related to pesticide use, crop rotation, record keeping, and which considers cultural, biological and other chemical control practices.
- Monitor after application for unexpected target pest survival. If the level of survival suggests that the presence of resistance, consult with your local university specialist or certified pest control advisor.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.
- For further information or to report suspected resistance contact Aceto Life Sciences, LLC at Aceto@aceto.com.

CHEMIGATION (For Use on Field Corn, Popcorn and Corn Grown for Seed Production Only)

Suremite™ Miticide alone or in combination with other products which are registered for application through sprinkler irrigation may be applied through irrigation systems. Apply this product only through micro-irrigation (individual spaghetti tube), drip irrigation, overhead irrigation or motorized calibrated irrigation equipment. Do not apply through any other type of irrigation system. Lack of effectiveness can result from non-uniform distribution of treated water. If you have questions about calibration, contact your State Extension Service specialist, equipment manufacturer or other experts. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments.

Using Water from Public Water Systems

- Do not apply Suremite™ Miticide through any irrigation system physically connected to a public water system.

Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days per year. Suremite™ Miticide may be applied through irrigation systems which may be supplied by a public water system only if the water from the public water system is discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and to top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe. Before beginning chemigation, always make sure that the air gap exists and that there is no blockage of the overflow of the reservoir tank.

Any irrigation system using water supplied from a public water system must also meet the following requirements:

Operating Instructions for All Specified Types of Irrigation Systems

1. The system must be calibrated to uniformly apply the rates specified. If you have questions about calibration, contact your State Extension Service specialist, equipment manufacturer or water management experts.
2. The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
3. The pesticide injection pipeline must contain a functional, automatic, quick closing check valve to prevent the flow of fluid back toward the injection pump.

4. The pesticide injection pipeline must also contain a functional, normally closed, solenoid operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
6. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
7. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
8. Do not apply when wind speed favors drift beyond the area intended.

Calibration and Application Instructions

Apply Suremite™ Miticide under the schedule specified in the specific crop rates/instructions, not according to the irrigation schedule, unless the events coincide. In general, set the equipment to apply the minimum amount of water per acre. Run the system at 95-100 % of the manufacturer's maximum rated travel speed.

The following calibration and application techniques are provided for user reference, but do not constitute a warranty of fitness for application through sprinkler irrigation equipment. Check with State and local regulatory agencies for potential use restrictions before applying any agricultural chemical through sprinkler irrigation equipment.

Center Pivot Irrigation Equipment

1. Use only drive systems that provide uniform water distribution.
2. Do not use end guns when chemigating Suremite™ Miticide through center pivot systems because of non-uniform application.
3. Plug the first nozzle closest to the well head to protect the water source.
4. Determine the size of the area to be treated.
5. Determine the time required to apply 0.10 to 0.15 inches of water over the area to be treated when the system and injection equipment are operated at normal pressures as recommended by the equipment manufacturer. Run the system at 95 to 100% of the manufacturer's rated maximum travel speed.
6. Using water, determine the injection pump output when operated at normal line pressure.
7. Determine the amount of Suremite™ Miticide, and any tankmix partners, required to treat the area covered by the irrigation system.
8. Add the required amount of Suremite™ Miticide, and any tankmix partners, and sufficient water to meet the injection time requirements to the solution tanks. (See "Mixing Instructions" section of this label.)
9. Make sure the system is fully charged with water before starting injection of the Suremite™ Miticide solution. Time the injection to last at least as long as it takes to bring the system to full pressure.
10. Maintain constant agitation in the solution tank during the injection period.
11. Inject the specified amount of Suremite™ Miticide per acre continuously for one complete revolution of the system.
12. Stop the injection equipment after treatment is complete. Continue to operate the system until the Suremite™ Miticide solution has cleared all of the sprinkler heads.
13. Allow time for all lines to flush the pesticide through all nozzles before turning off irrigation water.

Solid Set, Hand Move and Moving Wheel Irrigation Equipment

1. Determine the acreage covered by the sprinklers.
2. Fill injector solution tank with plain water and calibrate the flow rate of the system to deliver the contents of the tank over a 30 to 50 minute time interval at the end of the irrigation cycle.
3. Determine the amount of Suremite™ Miticide required to treat the area covered by the irrigation system.
4. Add the required amount of Suremite™ Miticide, and any other tankmix partners, into the same quantity of water used to calibrate the injection period. (See "Mixing Instructions" section of this label.)
5. Operate the system at the same pressure and time interval established during the calibration.
6. Inject specified amount of Suremite™ Miticide per acre for either a 30 to 50 minute period at the end of a regular irrigation set, or as a 30 to 50 minute injection as a separate application not associated with a regular irrigation to maximize retention of the insecticide by the foliage.
7. Stop injection equipment after treatment is completed. Continue to operate the system until the Suremite™ Miticide solution has cleared the last sprinkler head. To ensure lines are flushed and free from remaining pesticides, a dye indicator may be injected into the lines to mark the end of the application period.

APPLICATION DIRECTIONS

Suremite™ Miticide is a wettable powder packaged in a water soluble bag (WSB) designed to be diluted with water at the rates listed in the specific crop use directions. Applications should be made immediately after the spray solution is prepared. Thorough spray coverage is essential for effective control. Applications may be made with high or low volume spray equipment that provides thorough coverage of the plant. Apply with properly calibrated spray equipment. For best results, apply when pest populations are beginning to build, before reaching economic thresholds.

Since this product forms a suspension in water, it is important to maintain good agitation during mixing and spraying. If the spray suspension is allowed to settle for a short period of time, be sure to agitate the spray suspension for a minimum 10 minutes.

Apply spray solutions within 24 hours after mixing.

Use Rate Equivalency

Since this product is packaged in a water soluble bag (WSB) unit dose, e.g. 3.0 oz per WSB, then the number of areas treated at various use rates per WSB are listed below.

Rate oz./ acre	Acres Treated	Number of 3 oz. WSB
0.66	4.55	1
1.0	3.0	1
1.1	2.73	1
2.0	1.5	1
2.2	1.36	1
3.0	1.0	1
4.0	0.75	1

INSTRUCTIONS FOR USING WATER SOLUBLE BAGS

Water Soluble Bags (WSBs) are designed to dissolve in water. Agitation may be used, if necessary, to help dissolve the WSB.

Failure to follow handling and mixing instructions can increase your exposure to the pesticide products in WSBs. WSBs, when used properly, qualify as a closed mixing/loading system under the Agricultural Worker Protection Standard [40 CFR 170.607(d)].

Handling Instructions

Follow these steps when handling pesticide products in WSBs.

1. Mix in spray tank only.
2. Handle WSB(s) in a manner that protects package from breakage and/or unintended release of contents. If package is broken, put on PPE required for clean-up and then continue with mixing instructions.
3. Keep the WSB(s) in outer packaging until just before use.
4. Keep the WSB dry prior to adding to the spray tank.
5. Handle with dry gloves and according to the label instructions for PPE.
6. Keep WSB intact. Do not cut or puncture WSB.
7. Reseal the WSB outer packaging to protect any unused WSB(s).

Mixing Instructions

Follow the steps below when mixing this product, including if tank mixed with other pesticide products. If being tank mixed, the mixing directions 1 through 9 below take precedence over the mixing directions of the other tank mix products. WSBs may, in some cases, be mixed with other pesticide products so long as the directions for use of all mixed products do not conflict. Do not tank mix this product with products that prohibit tank mixing or have conflicting mixing directions.

1. If a basket or strainer is present in the tank hatch, remove prior to adding the WSB to the tank.
2. Fill tank with water to approximately one-third to one-half of the desired final volume of spray.
3. Stop adding water and stop any agitation.
4. Place intact/unopened WSB(s) into the tank.
5. Do not spray water from a hose or fill pipe to break or dissolve the WSB(s).
6. Start mechanical and recirculation agitation from the bottom of tank without using any overhead recirculation, if possible. If overhead recirculation cannot be turned off, close the hatch before starting agitation.
7. Dissolving the WSB(s) may tank up to 5 minutes or longer, depending on water temperature, water hardness and intensity of agitation.
8. Stop agitation before tank lid is opened.
9. Open the lid to the tank, exercising caution to avoid contact with dusts or spray mix, to verify that the WSBs have fully dissolved and the contents have been thoroughly mixed into the solution.
10. Do not add other allowed products or complete filling the tank until the bags have fully dissolved and pesticide is thoroughly mixed.
11. Once the WSB have fully dissolved and any other products have been added to the tank, resume filling the tank with water to the desired level, close the tank lid, and resume agitation.
12. Use the spray solution when mixing is complete.
13. Maintain agitation of the diluted pesticide mix during transport and application.
14. It is unlawful to use any registered pesticide, including WSBs, in a manner inconsistent with its label.

TANK MIXTURES

To improve this product's effectiveness, apply in combination with other pesticide products that are registered for the same crop and application techniques. For current information on the best tank mixture partner in your area, consult with the local dealer, distributor or State Agricultural Extension service.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

If this product is to be tank mixed with other pesticides, conduct a compatibility test prior to mixing. Use a small container and mix all components in a small amount, usually 0.5 to 1qt. of spray. Mix this product as indicated above and then combine all products in the same ratio and order of addition as in the proposed spray mixture. Observe the mixture for indication of incompatibility which usual occurs in 10 to 30 minutes after mixing. If incompatibility is observed, try changing the order of addition of the components. The guideline on tank mixture partners is driven by formulation type. Start with water dispersible granules (WP's), wettable powders (WP's) including water soluble bags (WSB's), suspension concentrated (SC's) or flowable (F's), all with very good agitation. Next follow with water miscible concentrates and emulsifiable concentrates (EC's) before adding drift control additives, nonionic surfactants (NIS's) or crop oil concentrates (COC's). After vigorous agitation, there must be a homogeneous suspension. Let the final tank mixture stand and observe for any rapid settling or floating of components. If any indications of physical incompatibility develop, do not use this mixture for spraying.

SPRAY COVERAGE: All parts of the crop must receive uniform spray coverage or else desired result may not occur. Higher water volumes and increased spray pressure generally provide better coverage. Consult your local agricultural specialist for specific information on the best rates, timings, and spray volumes for your region.

Suremite™ Miticide Crop/Use Site Index

Crop	Page
Caneberry	9
Christmas Trees	10
Cotton	11
Cucurbit Vegetables	12
Field Corn, Popcorn, Corn (Grown for Seed Production)	13
Hops	14
Low Growing Berry	15
Mint (Peppermint and Spearmint)	16
Non-Bearing Fruit Trees	17
Pepper and Eggplant	18
Pome Fruit	19
Small Fruit Vine Climbing (except Fuzzy Kiwifruit)	20
Stone Fruit	21
Sweet Corn	22
Tree Nuts	23
Tropical and Sub-Tropical Fruit (Inedible Peel)	24

The use rate for this product is expressed in terms of the oz. of this product per acre as Rate oz. /acre and Rate lb. a.i. acre. The pre-harvest interval (PHI) is the required number of days between the last application of this product and the harvesting of the crop.

Caneberry - Subgroup 13-07 A:**Blackberry; loganberry; raspberry, black and red; wild raspberry; cultivars, varieties, and/or hybrids of these.**

Pest	Rate oz/acre	Rate lb a.i./acre	Application Instructions
European Red Spider Mite McDaniel Spider Mite Two-spotted Spider Mite Yellow Spider Mite	2.0 – 3.0	0.09 – 0.135	<p>Apply by ground as a full coverage spray in a minimum spray solution of 50 gallons per acre.</p> <p>Use a sufficient volume of water to ensure thorough coverage of foliage, reproductive parts and fruit. This prevents mites from dispersal, feeding and reproduction.</p> <p>This product is predominately an ovicide/larvicide. Use when mite populations are low.</p> <p>Apply before the population levels reaches economic threshold, Consult local and state agricultural authorities for specific details.</p> <p>Select the lower rate for light infestations and the higher rate for heavy infestations.</p> <p>Adjust volume based on size of the bush canopy. Use higher volumes for older and varieties that have more compact and dense foliage.</p> <p>Good spray coverage is essential. Orient nozzles to assure good coverage.</p> <p>Use a higher volume of spray will assure better coverage especially under adverse conditions, such as hot, dry weather and/or a dense canopy.</p>

Restrictions:

Do not apply more than 1 application per year.

Do not use more than 3 oz. of this product per acre per year (0.135 lb. a.i. per acre per year).

Pre-Harvest Interval (PHI) is 0 day.

Do not use below 2.0 oz. of this product per application (0.09 lb. a.i. per application), as this may result in poor control and contribute to development of resistance to etoxazole among mite populations.

Christmas Trees

Pest	Rate oz/acre	Rate lb a.i./acre	Application Instructions
McDaniel Spider Mite Pacific Spider Mite Spruce Spider Mite Two-spotted Spider Mite	2.0 – 3.0	0.09 – 0.135	Apply by ground as full coverage spray or in a minimum spray solution of 10 gallons per acre by air. This product is predominately an ovicide/larvicide. Use when mite populations are low. Apply early in the life cycle of mites before the population levels reaches economic threshold, Consult local and state agricultural authorities for specific details. Select the lower rate for light infestations and the higher rate for heavy infestations.

Restrictions:

Do not apply more than 1 application per year.

Do not use more than 3.0 oz. of this product per acre per year (0.135 lb. a.i. per acre per year).

Do not use below 2.0 oz. of this product per application (0.09 lb. a.i. per application), as this may result in poor control and contribute to development of resistance to etoxazole among mite populations.

Cotton

Pest	Rate oz/acre	Rate lb a.i./acre	Application Instructions
Carmine Spider Mite Pacific Spider Mite Two-spotted Spider Mite	0.66 – 1.0	0.03 – 0.045	<p>Apply a minimum spray solution of 10 up to 50 gallons per acre by ground.</p> <p>Apply a minimum spray solution of 3 up to 10 gallons per acre by air.</p> <p>This product is predominately an ovicide/larvicide. Use when mite populations are low.</p> <p>Use a sufficient volume of water to ensure thorough coverage of foliage.</p> <p>Apply when the mite populations are low and beginning to build up on the plants before the population levels reaches economic threshold.</p> <p>Consult local and state agricultural authorities for specific details.</p> <p>Select the lower rate for light infestations and the higher rate for heavy infestations.</p> <p>Adjust volume based on size of the foliage.</p> <p>Good spray coverage is essential. Orient nozzles to assure good coverage.</p> <p>Use a higher volume of spray will assure better coverage especially under adverse conditions, such as hot, dry weather and/or a dense canopy.</p>

Restrictions:

Do not apply more than 1 application per year.

Do not use more than 1.0 oz. of this product per acre per year (0.045 lb. a.i. per acre per year).

Pre-Harvest Interval (PHI) is 28 days.

Do not use below 0.66 oz. of this product per application (0.03 lb. a.i. per application), as this may result in poor control and contribute to development of resistance to etoxazole among mite populations.

Cucurbit Vegetables - Crop Group 9:

Chayote (fruit); Chinese waxgourd (Chinese preserving melon); citron melon; cucumber; gherkin; gourd, edible (includes hyotan, cucuzza, hechima, Chinese okra); Momordica spp. (includes balsam apple, balsam pear, bitter melon, Chinese cucumber); muskmelon (includes cantaloupe); pumpkin; squash, summer; squash, winter (includes butternut squash, calabaza, hubbard squash, acorn squash, spaghetti squash); watermelon.

Pest	Rate oz/acre	Rate lb a.i./acre	Application Instructions
Carmine Spider Mite Pacific Spider Mite Strawberry Spider Mite Two-spotted Spider Mite	2.0 – 3.0	0.09 – 0.135	<p>Apply a minimum spray solution of 10 gallons per acre by ground.</p> <p>Apply a minimum spray solution of 3 gallons per acre by air**.</p> <p>This product is predominately an ovicide/larvicide. Use when mite populations are low.</p> <p>Use a sufficient volume of water to ensure thorough coverage of crop's foliage and reproductive parts. This prevents mites from dispersal, feeding and reproduction.</p> <p>Apply when the mite populations are low and beginning to build up on the plants before the population levels reaches economic threshold.</p> <p>Consult local and state agricultural authorities for specific details.</p> <p>Select the lower rate for light infestations and the higher rate for heavy infestations.</p> <p>Adjust volume based on size of the foliage. Use higher volumes on more mature plants and varieties that have more compact and dense foliage.</p> <p>Good spray coverage is essential. Orient nozzles to assure good coverage.</p> <p>Use a higher volume of spray will assure better coverage especially under adverse conditions, such as hot, dry weather and/or a dense canopy.</p>

Restrictions:

** Do not use aerial applications in New York

Do not apply more than 1 application per year.

Do not use more than 2.0 oz. of this product per acre per year (0.09 lb. a.i. per acre per year), as this may result in poor control and contribute to development of resistance to etoxazole among mite populations.

Pre-Harvest Interval (PHI) is 7 day.

Field Corn, Popcorn Corn, Corn (Grown for Seed Production)

Pest	Rate oz/acre	Rate lb a.i./acre	Application Instructions
Banks Grass Mite ^A Carmine Spider Mite Pacific Spider Mite Strawberry Spider Mite Two-spotted Spider Mite	1.0 – 3.0	0.045 – 0.135	<p>Apply a minimum spray solution of 10 gallons per acre by ground. Apply a minimum spray solution of 3 gallons per acre by air^{**}.</p> <p>This product is predominately an ovicide/larvicide. Use when mite populations are low.</p> <p>Use a sufficient volume of water to ensure thorough coverage of foliage.</p> <p>Apply when the mite populations are beginning to build up on the plants before the population levels reaches economic threshold. Also apply this product before the tasseling growth stage. Consult local and state agricultural authorities for specific details.</p> <p>Select the lower rate for light infestations and the higher rate for heavy infestations.</p> <p>Adjust volume based on size of the foliage.</p> <p>Good spray coverage is essential. Orient nozzles to assure good coverage.</p> <p>Use a higher volume of spray will assure better coverage especially under adverse conditions, such as hot, dry weather and/or a dense canopy.</p> <p>For corn (grown for seed) use this product before tasseling growth stage.</p> <p>For applications using chemigation refer to chemigation section.</p> <p>^A Applications targeting exclusively Banks grass mites may require higher rates within the rate range of 1.0 to 3.0 oz/acre particularly if populations have exceeded the established threshold in the fields to be treated with this product. Also, under extreme drought conditions and higher populations of Banks grassmites may require an additional application of a non-etoxazole miticide.</p>

Restrictions:

****** Do not use aerial applications in New York

Do not apply more than 1 application per year.

Do not use more than 3.0 oz. of this product per acre per year (0.135 lb. a.i. per acre per year).

Pre-Harvest Interval (PHI) is 21 days.

Do not use below 1.0 oz. of this product per application (0.045 lb. a.i. per application), as this may result in poor control and contribute to development of resistance to etoxazole among mite populations.

Hops

Pest	Rate oz/acre	Rate lb a.i./acre	Application Instructions
Two-spotted Spider Mite	3.0 – 4.0	0.135 – 0.18	<p>Apply a minimum spray solution of 50 gallons per acre by ground.</p> <p>Use a sufficient volume of water to ensure thorough coverage of foliage.</p> <p>This product is predominately an ovicide/larvicide. Use when mite populations are low.</p> <p>Apply before the population levels reaches economic threshold, Consult local and state agricultural authorities for specific details.</p> <p>Select the lower rate for light infestations and the higher rate for heavy infestations.</p> <p>Adjust volume based on size of the crop. Use higher volumes for more compact and dense foliage.</p> <p>Good spray coverage is essential. Orient nozzles to assure good coverage.</p> <p>Use a higher volume of spray will assure better coverage especially under adverse conditions, such as hot, dry weather and/or a dense canopy.</p>

Restrictions:

Do not apply more than 1 application per year.

Do not use more than 4.0 oz. of this product per acre per year (0.18 lb. a.i. per acre per year).

Pre-Harvest Interval (PHI) is 7 days.

Do not use below 3.0 oz. of this product per application (0.135 lb. a.i. per application), as this may result in poor control and contribute to development of resistance to etoxazole among mite populations.

Low-growing Berry Subgroup - Subgroup 13-07 G:

Bearberry; bilberry; blueberry, lowbush; cloudberry; cranberry; lingonberry; muntries; partridgeberry; strawberry; cultivars, varieties, and/or hybrids of these.

Pest	Rate oz/acre	Rate lb a.i./acre	Application Instructions
European Red Mite Pacific Spider Mite Two-spotted Spider Mite	2.0 – 3.0	0.09 – 0.135	<p>Apply a minimum spray solution of 100 gallons per acre by ground. Use a sufficient volume of water to ensure thorough coverage of foliage, reproductive parts and fruit. This prevents mites from dispersal, feeding and reproduction.</p> <p>This product is predominately an ovicide/larvicide. Use when mite populations are low.</p> <p>Apply early in the life cycle of mites before the population levels reaches economic threshold, Consult local and state agricultural authorities for specific details.</p> <p>Select the lower rate for light infestations and the higher rate for heavy infestations.</p> <p>This product will not control Cyclamine Mites. Use another miticide registered for this pest if these mites are a problem.</p> <p>Adjust volume based on size of the bush canopy. Use higher volumes for older and varieties that have more compact and dense foliage.</p> <p>Good spray coverage is essential. Orient nozzles to assure good coverage.</p> <p>Use a higher volume of spray will assure better coverage especially under adverse conditions, such as hot, dry weather and/or a dense canopy.</p>
Two-spotted Spider Mite	2.0 – 3.0	0.09 – 0.135	<p>Apply a minimum spray solution of 100 gallons per acre by ground. Alternate with other non-pyrethroid insecticides if retreatment is needed in less than 30 days to comply with local IPM programs.</p>

Restrictions:

Do not apply more than 1 application per year.

Do not use more than 3.0 oz. of this product per acre per year (0.135 lb. a.i. per acre per year).

Pre-Harvest Interval (PHI) is 1 day.

Do not use below 2.0 oz. of this product per application (0.09 lb. a.i. per application), as this may result in poor control and contribute to development of resistance to etoxazole among mite populations.

Mint (Peppermint and Spearmint)

Pest	Rate oz/acre	Rate lb a.i./acre	Application Instructions
Pacific Spider Mite Strawberry Spider Mite Two-spotted Spider Mite	2.0 - 4.0	0.09 – 0.18	<p>Apply a minimum spray solution of 50 gallons per acre by ground. Apply a minimum spray solution of 10 gallons per acre by air**.</p> <p>This product is predominately an ovicide/larvicide. Use when mite populations are low.</p> <p>Use a sufficient volume of water to ensure thorough coverage of foliage.</p> <p>Apply when the mite populations are beginning to build up on the plants before the population levels reaches economic threshold. Consult local and state agricultural authorities for specific details.</p> <p>Select the lower rate for light infestations and the higher rate for heavy infestations.</p> <p>Adjust volume based on size of the foliage.</p> <p>Good spray coverage is essential. Orient nozzles to assure good coverage.</p> <p>Use a higher volume of spray will assure better coverage especially under adverse conditions, such as hot, dry weather and/or a dense canopy.</p>

Restrictions:

** Do not use aerial applications in New York

Do not apply more than 1 application per year.

Do not use more than 4.0 oz. of this product per acre per year (0.18 lb. a.i. per acre per year).

Pre-Harvest Interval (PHI) is 7 days.

Do not use below 2.0 oz. of this product per application (0.09 lb. a.i. per application), as this may result in poor control and contribute to development of resistance to etoxazole among mite populations.

Non-Bearing Fruit Trees

Pest	Rate oz/acre	Rate lb a.i./acre	Application Instructions
European Red Spider Mite McDaniel Spider Mite Pacific Spider Mite Two-spotted Spider Mite	2.0 - 3.0	0.09 – 0.135	Apply as a full coverage spray by ground with airblast equipment. This product is predominately an ovicide/larvicide. Use when mite populations are low. Apply early in the life cycle of mites before the population levels reaches economic threshold, Consult local and state agricultural authorities for specific details. Select the lower rate for light infestations and the higher rate for heavy infestations.

Restrictions:

Do not harvest fruit from treated trees within one (1) year of application.

Do not apply more than 1 application per year.

Do not use more than 3.0 oz. of this product per acre per year (0.135 lb. a.i. per acre per year).

Do not use below 2.0 oz. of this product per application (0.09 lb. a.i. per application), as this may result in poor control and contribute to development of resistance to etoxazole among mite populations.

Pepper/Eggplant - Subgroup 8-10B:

African eggplant, eggplant, pea eggplant, pepino, roselle, scarlet eggplant, bell pepper, martynia, non-bell pepper, okra, cultivars, varieties and/or hybrids of these.

Pest	Rate oz/acre	Rate lb a.i./acre	Application Instructions
Two-spotted Spider Mite Broad Mite (Polyphagotarsonemus latus)	2.0 - 3.0	0.09 - 0.135	<p>Apply a minimum spray solution of 20 gallons per acre by ground. Use a sufficient volume of water to ensure thorough coverage of crop's foliage and reproductive parts.</p> <p>This product is predominately an ovicide/larvicide. Use when mite populations are low.</p> <p>Apply early in the life cycle of mites before the population levels reaches economic threshold, Consult local and state agricultural authorities for specific details.</p> <p>Select the lower rate for light infestations and the higher rate for heavy infestations.</p> <p>Adjust volume based on size of the bush canopy. Use higher volumes on more mature plants and varieties that have more compact and dense foliage.</p> <p>Good spray coverage is essential. Orient nozzles to assure good coverage.</p> <p>Use a higher volume of spray will assure better coverage especially under adverse conditions, such as hot, dry weather and/or a dense canopy.</p>

Restrictions:

Do not apply more than 1 application per year.

Do not use more than 3.0 oz. of this product per acre per year (0.135 lb. a.i. per acre per year).

Pre-Harvest Interval (PHI) is 7 days.

Do not use below 2.0 oz. of this product per application (0.09 lb. a.i. per application), as this may result in poor control and contribute to development of resistance to etoxazole among mite populations.

Pome Fruit - Crop Group 11: Apple; crabapple; loquat; mayhaw; pear; pear, Asian; quince

Pest	Rate oz/acre	Rate lb a.i./acre	Application Instructions
European Red Spider Mite McDaniel Spider Mite Pacific Spider Mite Two-spotted Spider Mite	2.0 - 3.0	0.09 – 0.135	<p>Apply a minimum spray solution of 100 gallons by ground as a full coverage spray with airblast equipment.</p> <p>This product is predominately an ovicide/larvicide. Use when mite populations are low.</p> <p>Apply early in the life cycle of mites before the population levels reaches economic threshold, Consult local and state agricultural authorities for specific details.</p> <p>Select the lower rate for light infestations and the higher rate for heavy infestations.</p> <p>Use a sufficient volume of water to ensure thorough coverage of fruit and leaf surfaces. Adjust volume based on size of the tree canopy, Use higher volumes for dense canopies.</p> <p>This product will not control Rust Mites or Blister Mites. Use another miticide registered for these pests if these mites are a problem.</p>

Restrictions:

Do not apply more than 1 application per year.

Do not use more than 3.0 oz. of this product per acre per year (0.135 lb. a.i. per acre per year).

Pre-Harvest Interval (PHI) is 14 days.

Do not use below 2.0 oz. of this product per application (0.09 lb. a.i. per application), as this may result in poor control and contribute to development of resistance to etoxazole among mite populations.

Small Fruit Vine Climbing Subgroup, except Fuzzy Kiwifruit - Subgroup 13-07 F:
Amur river grape; gooseberry; grape; kiwifruit, hardy; maypop; schisandra berry; cultivars, varieties, and/or hybrids of these.

Pest	Rate oz/acre	Rate lb a.i./acre	Application Instructions
European Red Spider Mite Pacific Spider Mite Two-spotted Spider Mite Willamette Mite	2.0 - 3.0	0.09 - 0.135	<p>Apply a full coverage spray with a minimum spray solution of 25 gallons per acre by ground.</p> <p>This product is predominately an ovicide/larvicide. Use when mite populations are low.</p> <p>Apply early in the life cycle of mites before the population levels reaches economic threshold, Consult local and state agricultural authorities for specific details.</p> <p>Select the lower rate for light infestations and the higher rate for heavy infestations.</p> <p>This product will not control Rust Mites or Blister Mites. Use another miticide registered for these pests if these mites are a problem.</p> <p>Use a sufficient volume of water to ensure thorough coverage of fruit and leaf surfaces. Adjust volume based on size of the tree canopy, Use higher volumes for dense canopies.</p>

Restrictions:

Do not apply more than 1 application per year.

Do not use more than 3.0 oz. of this product per acre per year (0.135 lb. a.i. per acre per year).

Pre-Harvest Interval (PHI) is 14 days.

Do not use below 2.0 oz. of this product per application (0.09 lb. a.i. per application), as this may result in poor control and contribute to development of resistance to etoxazole among mite populations.

Stone Fruit - Crop Group 12-12:

Apricot; cherry, sweet; cherry, tart; nectarine; peach; plum; plum, Chickasaw; plum, Damson; plum, Japanese; Plumcot, Prune (fresh)

Pest	Rate oz/acre	Rate lb a.i./acre	Application Instructions
European Red Spider Mite Pacific Spider Mite Two-spotted Spider Mite	2.0 - 3.0	0.09 - 0.135	<p>Apply a minimum spray solution of 50 gallons per acre by ground. This product is predominately an ovicide/larvicide. Use when mite populations are low.</p> <p>Apply early in the life cycle of mites before the population levels reaches economic threshold, Consult local and state agricultural authorities for specific details.</p> <p>Select the lower rate for light infestations and the higher rate for heavy infestations.</p> <p>Use a sufficient volume of water to ensure thorough coverage of fruit and leaf surfaces. Adjust volume based on size of the tree canopy, Use higher volumes for dense canopies.</p>

Restrictions:

Do not apply more than 1 application per year.

Do not use more than 3.0 oz. of this product per acre per year (0.135 lb. a.i. per acre per year).

Pre-Harvest Interval (PHI) is 7 days.

Do not use below 2.0 oz. of this product per application (0.09 lb. a.i. per application), as this may result in poor control and contribute to development of resistance to etoxazole among mite populations.

Sweet Corn (Except in CA)

Pest	Rate oz/acre	Rate lb a.i./acre	Application Instructions
Banks Grass Mite ^A Carmine Spider Mite Pacific Spider Mite Strawberry Spider Mite Two-spotted Spider Mite	1.0 - 3.0	0.045 - 0.135	<p>Apply a minimum spray solution of 10 gallons per acre by ground.</p> <p>Apply a minimum spray solution of 3 gallons per acre by air^{**}.</p> <p>This product is predominately an ovicide/larvicide. Use when mite populations are low.</p> <p>Use a sufficient volume of water to ensure thorough coverage of foliage.</p> <p>Apply when the mite populations are beginning to build up on the plants before the population levels reaches economic threshold.</p> <p>Also apply this product before the tasseling growth stage to allow good spray penetration and better control through the plant's canopy. Consult local and state agricultural authorities for specific details.</p> <p>Select the lower rate for light infestations and the higher rate for heavy infestations.</p> <p>Adjust volume based on size of the foliage.</p> <p>Good spray coverage is essential. Orient nozzles to assure good coverage.</p> <p>Use a higher volume of spray will assure better coverage especially under adverse conditions, such as hot, dry weather and/or a dense canopy.</p> <p>^A Applications targeting exclusively Banks grass mites may require higher rates within the rate range of 1.0 to 3.0 oz/acre particularly if populations have exceeded the established threshold in the fields to be treated with this product. Also, under extreme drought conditions and higher populations Banks grass mites may require an additional application of a non-etoxazole miticide.</p>

Restrictions:

^{**} Do not use aerial applications in New York

Do not apply more than 1 application per year.

Do not use more than 3.0 oz. of this product per acre per year (0.135 lb. a.i. per acre per year).

Pre-Harvest Interval (PHI) is 21 days.

Do not use below 1.0 oz. of this product per application (0.045 lb. a.i. per application), as this may result in poor control and contribute to development of resistance to etoxazole among mite populations.

Tree Nuts - Crop Group 14-12: almond; beechnut; Brazil nut; butternut; cashew; chestnut; chinquapin; hazelnut(filbert); hickory nut; macadamia nut; pecan; pistachio; walnut, black; walnut, English.

Pest	Rate oz/acre	Rate lb a.i./acre	Application Instructions
European Red Spider Mite Pacific Spider Mite Pecan Leaf Scorch Mite (Eotetranychus hicoriae) (Except in CA) Two-spotted Spider Mite	2.0 - 3.0	0.09 - 0.135	Apply as a full coverage spray by ground with airblast equipment. This product is predominately an ovicide/larvicide. Use when mite populations are low. Apply early in the life cycle of mites before the population levels reaches economic threshold, Consult local and state agricultural authorities for specific details. Select the lower rate for light infestations and the higher rate for heavy infestations.

Restrictions:

Do not apply more than 1 application per year.

Do not use more than 3.0 oz. of this product per acre per year (0.135 lb. a.i. per acre per year).

Pre-Harvest Interval (PHI) is 7 days.

Do not use below 2.0 oz. of this product per application (0.09 lb. a.i. per application), as this may result in poor control and contribute to development of resistance to etoxazole among mite populations.

Tropical and Sub-Tropical (Inedible Peel):**Avocado, papaya, star apple, black sapote, mango, sapodilla, canistel, mamey sapote.**

Pest	Rate oz/acre	Rate lb a.i./acre	Application Instructions
Avocado Brown Mite Persea Mite Two-spotted Spider Mite	2.0 - 3.0	0.09 - 0.135	<p>Apply a minimum spray solution of 50 gallons per acre by ground.</p> <p>Apply a minimum spray solution of up to 10 gallons per acre by air**.</p> <p>This product is predominately an ovicide/larvicide. Use when mite populations are low.</p> <p>Use a sufficient volume of water to ensure thorough coverage of crop's foliage and reproductive parts. This prevents mites from dispersal, feeding and reproduction.</p> <p>Apply when the mite populations are low and beginning to build up on the plants before the population levels reaches economic threshold.</p> <p>Consult local and state agricultural authorities for specific details.</p> <p>Select the lower rate for light infestations and the higher rate for heavy infestations.</p> <p>Adjust volume based on size of the foliage. Use higher volumes on more mature plants and varieties that have more compact and dense foliage.</p> <p>Good spray coverage is essential. Orient nozzles to assure good coverage.</p> <p>Use a higher volume of spray will assure better coverage especially under adverse conditions, such as hot, dry weather and/or a dense canopy.</p>

Restrictions:

**Do not use aerial applications in New York.

Do not apply more than 1 application per year.

Preharvest interval (PHI) is 1 day.

Do not use more than 3.0 oz. of this product per acre per year (0.135 lb. a.i. per acre per year).

Do not use below 2.0 oz. of this product per application (0.09 lb. a.i. per application), as this may result in poor control and contribute to development of resistance to etoxazole among mite populations.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

PESTICIDE STORAGE: Store in original container in a cool, dry and secure location.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Open dumping is prohibited.

CONTAINER HANDLING: Nonrefillable outer bag. Do not reuse or refill outer bag. Completely empty bag into application equipment, then dispose of the empty bag in a sanitary landfill or by incineration.

Batch Code: _____

WARRANTY DISCLAIMER AND NOTICE**IMPORTANT: READ BEFORE USE**

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Aceto Life Sciences, L.L.C. To the extent consistent with applicable law all such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, ACETO LIFE SCIENCES, L.L.C. MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Aceto Life Sciences, L.L.C. is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, ACETO LIFE SCIENCES, L.L.C. DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: To the extent consistent with applicable law, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT ACETO LIFE SCIENCES, L.L.C.'S ELECTION, THE REPLACEMENT OF PRODUCT.

All trademarks are proprietary of their respective owners.

Active ingredient made in China, formulated in USA and package in USA.

6.0000

6.0000

6.0000

6.0000

6.0000

6.0000

6.0000

ETOXAZOLE GROUP 10B INSECTICIDE

Suremite™ Miticide

Miticide for use on Caneberry, Christmas Trees, Cotton, Cucurbit Vegetables, Field Corn, Popcorn, Corn (Grown for Seed), Hops, Low Growing Berry, Mint, Non-Bearing Fruit Trees, Pepper and Eggplant, Pome Fruit, Small Fruit Vine Climbing (except Fuzzy Kiwifruit), Stone Fruit, Sweet Corn, Tree Nuts, and Tropical and Sub-Tropical Fruit (Inedible Peel).

ACTIVE INGREDIENT: etoxazole	72.0%
OTHER INGREDIENTS:	28.0%
TOTAL	100.0%

EPA Reg. No. 2749-592

EPA Est. No. 065387-AR-002

See inside booklet for Precautionary Statements and Directions for Use.

Read "LIMIT OF WARRANTY AND LIABILITY" before buying or using. If terms are not acceptable, return at once unopened.

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If swallowed: Call poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance; then give artificial respiration, preferably by mouth to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Emergency Assistance: Have the product container or label with you when calling a poison control center or doctor or going for treatment. FOR EMERGENCY MEDICAL HELP, CALL CHEMTREC TOLL FREE 800-424-9300 or 1-703-527-3887.

FOR CHEMICAL SPILL,
LEAK, FIRE, EXPOSURE
OR MEDICAL EMERGENCY
INVOLVING THIS
PRODUCT CALL
CHEMTREC® TOLL FREE
(800) 424-9300 OR
(703) 527-3887.

12 oz (4x3 oz WSBs)

NET CONTENTS

Manufactured for: Aceto Life Sciences, L.L.C., 4 Tri Harbor Court, Port Washington, NY 11050



PEEL BACK BOOK HERE AND RESEAL AFTER OPENING

File #: 87300 Art Order #: 394043 Item#: 205076-209
 folder: file://blmemart1/PRODUCTION/Products/800000/87300
 Size: Vert: 6.0000" X Horz: 6.7500" __ _Date: 1/27/2022 Time: 08:40:20
 colors: Black dieline
 alt by: joew _____ DocuProof---Checked by: _____
 code: No Code