

**GROUP 2 HERBICIDE**

# Rapport® BroadSpec

## Herbicide

WATER DISPERSIBLE GRANULE

FOR USE ON WHEAT, BARLEY, OATS, TRITICALE AND FALLOW

### ACTIVE INGREDIENTS:

Thifensulfuron-methyl

Methyl 3-[[[(4-methoxy-6-methyl-1,3,5-triazin-2-yl)amino]carbonyl]amino]sulfonyl]-2-thiophenecarboxylate ..... 25.0%

Tribenuron-methyl

Methyl 2-[[[(4-methoxy-6-methyl-1,3,5-triazin-2-yl)methylamino]carbonyl]amino]sulfonyl]benzoate ..... 25.0%

OTHER INGREDIENTS: ..... 50.0%

TOTAL: ..... 100.0%

### KEEP OUT OF REACH OF CHILDREN CAUTION

See Inside Booklet for FIRST AID and additional PRECAUTIONARY STATEMENTS

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand this label, find someone to explain it to you in detail).

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300

For Medical Emergencies Only, Call (877) 325-1840

EPA REG. NO. 71368-79

Manufactured for  
Nufarm, Inc.  
11901 S. Austin Avenue  
Alsip, IL 60803



**Nufarm**

Grow a better tomorrow.

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
CAUTION**

Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing.

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

Some materials that are chemical resistant to this product are listed below. If you want more options follow the instructions for Category A on an EPA chemical resistance category selection chart.

**Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material such as natural rubber
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

**USER SAFETY RECOMMENDATIONS**

**Users Should:**

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible wash thoroughly and change into clean clothing.

**FIRST AID**

**IF IN EYES:**

- Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

**HOT LINE NUMBER**

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.

**ENVIRONMENTAL HAZARDS**

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

**PESTICIDE HANDLING**

- Calibrate sprayers only with clean water away from the well site.
- Make scheduled checks of spray equipment.
- Ensure that all operation employees accurately measure pesticides.
- Mix only enough product for the job at hand.
- Avoid overfilling of spray tank.
- Do not discharge excess material on the soil at a single spot in the field, grove, or mixing/loading station.
- Dilute and agitate excess solution and apply at labeled rates or uses.
- Avoid storage of pesticides near well sites.
- When triple-rinsing the pesticide container, be sure to add the rinsate to the spray mix.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

## **AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

This product is for use on wheat, barley, oats, triticale and fallow in many states. Check with your state extension or Dept. of Agriculture before use, to be certain this product is registered in your state. Nufarm will not be responsible for losses or damages resulting from the use of this product in any manner not in accordance with instructions on this label.

## **USE INFORMATION**

This product is a water dispersible granule that is used for selective postemergence weed control in wheat (including durum), barley, oat, triticale and fallow. The best control is obtained when this product is applied to young, actively growing weeds. The specified use rate will depend on weed spectrum and size of weed at time of application. The degree and duration of control may depend on the following:

- weed spectrum and infestation intensity
- weed size at application
- environmental conditions at and following treatment

This product is noncorrosive, nonflammable, nonvolatile, and does not freeze. This product should be mixed, and completely dissolved in water and applied as a uniform broadcast spray.

## **ENVIRONMENTAL CONDITIONS AND BIOLOGICAL ACTIVITY**

This product is absorbed primarily through the foliage of plants, rapidly inhibiting the growth of susceptible weeds. One to three weeks after application to weeds, leaves of susceptible plants appear chlorotic, and the growing point subsequently dies.

This product provides the best control in vigorously growing crops that shade competitive weeds. Weed control in areas of thin crop stand or seeding skips may not be as satisfactory. However, a crop canopy that is too dense at application can intercept spray and reduce weed control.

The herbicidal action of this product may be affected in crops stressed from adverse environmental conditions (such as extreme temperatures or moisture), abnormal soil conditions, cultural practices, or variations in crop variety. In warm, moist conditions, the expression of herbicide symptoms is accelerated; in cold, dry conditions, expression of herbicide symptoms is delayed. In addition, weeds hardened-off by drought stress are less susceptible to this product.

## **RESTRICTIONS**

Injury to or loss of adjacent sensitive crops, desirable trees or vegetation may result from failure to observe the following:

- Do not apply, drain or flush equipment on or near desirable trees or other plants or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots.
- Do not use on lawns, walks, driveways, tennis courts, or similar areas. Prevent drift of spray to desirable plants.
- Do not apply this product by air in the state of New York.
- Do not apply this product through any irrigation system.
- Do not exceed the maximum application rate of 1 ounce per acre per crop season.
- Do not apply to wheat, barley, oats and triticale crops underseeded with another crop.
- Do not harvest wheat or barley sooner than 45 days after the last application of this product.
- When using this product in tank mixes or sequential applications with other products containing thifensulfuron-methyl and/or tribenuron-methyl, do not exceed the following limits:

Use	Active Ingredient	Maximum oz ai per Single Application	Maximum oz ai per Use Period
<b>Wheat, Barley, Triticale</b>	Thifensulfuron-methyl	0.45	0.75
	Tribenuron-methyl	0.25	0.25
<b>Oats</b>	Thifensulfuron-methyl	0.30	0.30
	Tribenuron-methyl	0.10	0.10
<b>Fallow, Pre-Plant, Burndown, Post Harvest, Crop Preemergence</b>	Thifensulfuron-methyl	0.45	0.75
	Tribenuron-methyl	0.25	0.25

## PRECAUTIONS

Take all necessary precautions to avoid all direct or indirect contact (such as spray drift) with non-target plants or areas.

Carefully observe all sprayer cleanup instructions both prior to and after using this product, as spray tank residue may damage crops other than wheat, barley, oats and triticale.

Wheat, barley, oats and triticale may differ in their response to various herbicides. Nufarm recommends that you first consult your state experiment station, university, or extension agent as to sensitivity to any herbicide. If no information is available, limit the initial use of this product to a small area.

Under certain conditions, such as heavy rainfall, prolonged cold weather (daily high temperature less than 50°F), or wide fluctuations in day/night temperatures prior to or soon after this product's application, temporary discoloration and/or crop injury may occur. To reduce the potential of crop injury, tank mix this product with 2,4-D (ester formulations perform best—see Tank Mixtures section of this label) and apply after the crop is in the tillering stage of growth.

This product should not be applied to wheat, barley, oats and triticale that is stressed by severe weather conditions, drought (including low levels of subsoil moisture), low fertility, water saturated soil, disease, or insect damage, as crop injury may result. Risk of injury is greatest when crop is in the 2- to 5-leaf stage. Severe winter stress, drought, disease, or insect damage following application also may result in crop injury.

Dry, dusty field conditions may result in reduced control in wheel track areas.

## APPLICATION INFORMATION USE RATE

Apply this product at a rate of 0.4 to 1.0 ounce per acre. When applying 0.4 to 0.6 ounce per acre, this product must be used in a tank-mix combination with other registered herbicides.

### WHEAT (INCLUDING DURUM), BARLEY AND TRITICALE

Apply 0.4 to 1.0 ounce of this product per acre to wheat (including durum), barley or triticale. The total amount of this product cannot exceed 1.0 ounce per acre per crop season.

### OAT (SPRING AND WINTER)

Apply 0.4 ounce per acre to oat for control of light broadleaf weed infestations and do not make more than one application (or more than 0.1 ounces of active ingredient tribenuron-methyl) per season. Additional broadleaf herbicide tank mixes labeled for oat are recommended for control of moderate to heavy infestations of broadleaf weeds.

### FALLOW

Apply 0.4 to 1.0 ounce of this product per acre to fallow. The total amount of this product cannot exceed 1.0 ounce per acre per crop season. This product should be applied in combination with other suitable registered fallow herbicides such as glyphosate plus 2,4-D (ester formulations work best) or glyphosate plus dicamba.

When this product is applied at a rate of 0.4 to 0.6 ounce per acre, this product must be used in a tank mix combination with other registered fallow herbicides.

### PRE-PLANT OR CROP PREEMERGENCE BURNDOWN

Apply 0.4 to 1.0 ounce of this product per acre as a burndown treatment prior to, or shortly after planting (prior to crop emergence). The total amount of this product cannot exceed 1.0 ounce per acre per crop season.

## POST HARVEST

Apply this product at 0.4 to 1.0 ounce per acre to crop stubble after harvest. Use the 1.0 ounce per acre rate when weed infestation is heavy and predominantly consists of those weeds listed under the “WEEDS PARTIALLY CONTROLLED” section of this label or when application timing and environmental conditions are marginal. (See the “APPLICATION TIMING” section of this label for restriction on planting intervals). This product should be applied in combination with other suitable registered burndown herbicides (See the “TANK MIXTURES” section of this label for additional information). Sequential treatments of this product may also be made provided the total amount of this product applied during one fallow/pre plant cropland season does not exceed 1.0 ounce per acre.

## APPLICATION TIMING

Since this product has very little or no soil activity, it controls only those weeds that have germinated; therefore, apply this product when all or most of the weeds have germinated. Annual broadleaf weeds should be past the cotyledon stage, actively growing, and less than 4” tall or wide. Rainfall immediately after treatment can wash this product off of weed foliage, resulting in reduced weed control. Four hours of dry weather are needed to allow this product to be sufficiently absorbed by weed foliage.

### WHEAT (INCLUDING DURUM), BARLEY, WINTER OATS AND TRITICALE

Make applications after the crop is in the 2-leaf stage, but before the flag leaf is visible.

### SPRING OATS

Make applications to spring planted oat after the crop has reached the 3 leaf stage but prior to jointing. Some oat varieties may be sensitive to this product including Ogle, Porter, and Premier. Not all varieties have been tested for sensitivity, therefore, it is best to consult local recommendations regarding oat varietal sensitivity prior to use.

### FALLOW

This product may be used as a fallow treatment, in the spring, summer or fall when the majority of weeds have emerged and are actively growing.

### PRE-PLANT OR CROP PREEMERGENCE BURNDOWN

Apply this product as a burndown treatment to wheat (including durum), barley and triticale to control emerged weeds prior to, or shortly after planting (prior to emergence). Make applications when the majority of weeds have emerged and are actively growing. Wheat, barley, rice and triticale may be replanted anytime after the application of this product. See “CROP ROTATION” for the time interval required before planting.

## POST HARVEST

This product may be used as a burndown treatment to crop stubble when the majority of weeds have emerged and are actively growing. See the “CROP ROTATION” section of this label for additional information.

## CROP ROTATION

Labeled crops may be planted at specified time intervals following application of labeled rates of this product. Use the time intervals listed below to determine the required time interval before planting. Refer to individual product labels to determine rotational crop restrictions when tank mixtures are used.

### Time Interval Before Planting\* (days after treatment with this product)

Crop	Days
Barley, Rice, Triticale, and Wheat (including durum)	0
Soybeans	7 <sup>^</sup> *
Cotton, Field Corn, and Grain Sorghum	14*
Sugarbeets, Winter Rape, and Canola	60
Any other crop	45

<sup>^</sup> When this product is applied at 0.5 ounce per acre or less the time interval for soybeans is 1 day.

\* When this product is used on light textured soils, such as sands and loamy sands, extend time to planting by 7 additional days.

Where this product is used on high pH soils (>7.9), extend time to planting by 7 additional days.

## WEEDS CONTROLLED

This product effectively controls the following weeds when used according to label directions:

Annual knawel	Curly dock	Redroot pigweed
Annual sowthistle	False chamomile	Russian thistle*
Black mustard	Field chickweed	Scentless chamomile/ mayweed
Blue/Purple mustard	Field pennycress	Shepherd's-purse
Broadleaf dock	Filaree (redstem, Texas)	Slimleaf lambsquarters
Bur buttercup	Flixweed	Smallflower buttercup
Bushy wallflower/ Treacle mustard	Green smartweed	Smallseed falseflax
Canada thistle *	Henbit	Stinking chickweed
Clasping pepperweed	Kochia *	Stinking mayweed/ dogfennel
Coast fiddleneck	Ladysthumb	Sunflower
Common buckwheat	Lanceleaf sage *	Swinecress
Common chickweed	London rocket	Tansymustard
Common cocklebur *	Marshelder	Tarweed fiddleneck
Common groundsel	Mayweed chamomile	Tumble/Jim Hill mustard
Common lambsquarters	Miners lettuce	Volunteer canola
Common ragweed *	Narrowleaf lambsquarters	Volunteer lentils
Common sunflower	Nightflowering catchfly	Volunteer peas
Corn chamomile	Pennsylvania smartweed	White cockle
Corn gromwell *	Pineappleweed	Wild buckwheat*
Corn spurry	Prickly lettuce*	Wild chamomile
Cowcockle	Prostrate knotweed	Wild mustard
Cress (mouse-ear)	Prostrate pigweed	
	Redmaids	

## WEEDS PARTIALLY CONTROLLED\*\*

This product partially controls the following weeds when used according to label directions:

Catchweed bedstraw  
Mallow (common, little)  
Marestail  
Nightshade (cutleaf, hairy)

\* See SPECIFIC WEED PROBLEMS for more information.

\*\* Partial control: A visual reduction of weed population as well as a significant loss of vigor. For better results, use the highest rate of this product per acre and include a tank mix partner such as 2,4-D, MCPA, Buctril/Maestro 2EC or Banvel/Clarity or Diablo/Clash (refer to TANK MIXTURES).

## SPECIFIC WEED PROBLEMS

**Canada thistle:** For control in wheat and barley, use 0.8 ounce per acre plus surfactant when all thistles are 4" to 8" with 2" to 6" of new growth. Make the application in the spring. Control will be improved by using this product in combination with 2,4-D or dicamba (refer to TANK MIXTURES).

**Common cocklebur, Common ragweed, Lanceleaf sage:** In wheat and barley, apply this product at 0.4 to 0.8 ounce per acre in combination with 2,4-D at rates from 1/4 to 3/8 pound active ingredient (ester formulations work best) when weeds are small and actively growing. When using 1/4 pound active ingredient of 2,4-D, be sure to add surfactant at the rate of 1/4 to 1/2 quart per 100 gallons of spray solution (0.06 to 0.125% v/v--use the higher rate under stress conditions).

**Corn gromwell, Wild buckwheat:** For control in wheat and barley, use 0.8 ounce this product per acre plus surfactant.

**Resistant biotypes including but not exclusive to (Kochia, Russian thistle, Prickly lettuce):** Naturally occurring resistant biotypes of these weeds are known to occur. For best results, use this product in a tank mix with a herbicide partner that is labeled for control of the resistant weed biotype in question.

## SPRAY ADJUVANTS

Always include a spray adjuvant with applications of this product. In addition to a spray adjuvant, an ammonium nitrogen fertilizer may be used.

Consult your Ag dealer or applicator, local Nufarm fact sheets, technical bulletins, and service policies prior to using an adjuvant system. If another herbicide is tank mixed with this product, select adjuvants authorized for use with both products. Products must contain only EPA-exempt ingredients.

## NONIONIC SURFACTANT (NIS)

- Apply 0.06% to 0.50% volume/volume (1/2 to 4 pints per 100 gallon of spray solution).
- Surfactant products must contain at least 60% nonionic surfactant with a hydrophilic/lipophilic balance (HLB) greater than 12.

See the Tank Mixtures section of this label for additional information.

## PETROLEUM CROP OIL CONCENTRATE (COC) OR MODIFIED SEED OIL (MSO)

- Apply at 1% volume/volume (1 gallon per 100 gallon spray solution) or 2% volume/volume under arid conditions.
- Oil adjuvants must contain at least 80% high quality, petroleum (mineral) or modified vegetable seed oil with at least 15% surfactant emulsifiers.

## BASIC BLEND ADJUVANTS

- Apply at 1% volume/volume (1 gallon per 100 gallon spray solution).]

## SPECIAL ADJUVANT TYPES

- Combination adjuvant products may be used at doses that provide the required amount of NIS, COC, MSO and/or ammonium nitrogen fertilizer. Consult product literature for use rates and restrictions.
- In addition to the adjuvants specified above, other adjuvant types may be used if they provide the same functionality and have been evaluated and approved by Nufarm product management. Consult separate Nufarm technical bulletins for detailed information before using adjuvant types not specified on this label.

## AMMONIUM NITROGEN FERTILIZER

- Use 2 quarts per acre of a high-quality urea ammonium nitrate (UAN), such as 28%N or 32%N, or 2 pounds per acre of a spray-grade ammonium sulfate (AMS). Use 4 quarts per acre UAN or 4 pounds per acre AMS under arid conditions.

## GROUND APPLICATION

For ground application 5 to 20 gallon per acre (GPA) should be used, for optimum spray distribution and thorough coverage. Select nozzle types that deliver medium to coarse droplets for best coverage and drift control. Set pressure according to the nozzle manufacturer's guidelines. Avoid use of nozzles and pressure that deliver fine droplets.

Raindrop RA nozzles are not recommended for this product applications, as weed control performance may be reduced.

Use screens that are 50-mesh or larger.

## AERIAL APPLICATION

Use nozzle types and arrangements that provide optimum spray distribution and maximum coverage.

- Use 2 to 5 GPA
- Use at least 3 GPA in Idaho, Oregon, or Utah.

Do not apply this product by air in the state of New York.

See the **Spray Drift Management** section of this label.

## CHEMIGATION

Do not apply this product through any irrigation system.

## PRODUCT MEASUREMENT

This product is measured using this product's volumetric measuring cylinder. The degree of accuracy of this cylinder varies by +/- 7.5%. For more precise measurement, use scales calibrated in ounces.

## TANK MIXTURES

This product may be tank mixed with other suitable registered herbicides to control weeds listed as suppressed, weeds resistant to this product or weeds not listed under **Weeds Controlled**. Read and follow all manufacturers' label instructions for the companion herbicide. If those instructions conflict with this label, do not tank mix the herbicide with this product.

This product can also be mixed with registered fungicides, insecticides, or liquid fertilizer for use on wheat, barley, or fallow.

## WITH 2,4-D (AMINE OR ESTER) OR MCPA (AMINE OR ESTER)

This product may be tank mixed with the amine or ester formulations of 2,4-D or MCPA herbicides for use on wheat and barley. For best results in the Red River Valley and adjacent areas of North Dakota and Minnesota, add the ester formulations of 2,4-D or MCPA herbicides to the tank at 3/8 pound active ingredient (such as 3/4 pint of a 4 lb/gal product, or 1/2 pint of a 6 lb/gal product). No additional surfactant is needed with this mixture.

For best results in other areas, add the ester formulations of 2,4-D or MCPA herbicides to the tank at 1/4 to 3/8 pound active ingredient (such as 1/2 to 3/4 pint of a 4 lb/gal product, or 1/3 to 1/2 pint of a 6 lb/gal product). Surfactant may be added to the mixture at 1/2 to 1 quart per 100 gallons of spray solution (0.125 to 0.25% v/v); however, adding surfactant may increase the potential for crop injury, especially at the higher phenoxy rates.

Higher rates of 2,4-D or MCPA may be used, but do not exceed the highest rate allowed by those respective labels. Read and follow all label instructions on timing, precautions, and warnings for these herbicides before using these tank mixtures.

## **WITH DICAMBA (SUCH AS BANVEL®/CLARITY®, DIABLO®/CLASH®)**

This product may be tank mixed with 1/16 to 1/8 pound active ingredient dicamba (such as 2 to 4 fluid ounces Banvel/Diablo, or 2 to 4 fluid ounces Clarity/Clash). Use higher rates when weed infestation is heavy. Surfactant may be added to the mixture at 1/2 to 1 quart per 100 gallons of spray solution (0.125 to 0.25% v/v); however, adding surfactant may increase the potential for crop injury. Refer to the specific dicamba label for application timing and restrictions. Tank mixes of this product plus dicamba may result in reduced control of some broadleaf weeds.

## **WITH 2,4-D (AMINE OR ESTER) AND BANVEL/CLARITY, DIABLO/CLASH**

This product may be applied in a 3-way tank mix with formulations of dicamba and 2,4-D. Make application of this product + 1/16 to 1/8 pound active ingredient dicamba (such as 2 to 4 fluid ounces Banvel/Diablo, or 2 to 4 fluid ounces Clarity/Clash) + 1/4 to 3/8 pound active ingredient of 2,4-D ester or amine per acre. Use higher rates when weed infestation is heavy. Surfactant may be added to the mixture at 1/2 to 1 quart per 100 gallons of spray solution (0.125 to 0.25% v/v); however, adding surfactant may increase the potential for crop injury. Consult the specific 2,4-D label, dicamba label, or local recommendations for more information and restrictions.

Apply this 3-way combination to winter wheat after the crop is tillering and prior to jointing (first node). In Spring Wheat (including Durum) apply after the crop is tillering and before it exceeds the 5-leaf stage.

In Spring Barley, apply after the crop is tillering and before it exceeds the 4-leaf stage.

## **WITH BROMOXYNIL (SUCH AS BUCTRIL®/MAESTRO® 2EC/BRONATE®/MAESTRO MA)**

This product may be tank mixed with bromoxynil containing herbicides registered for use on wheat, barley, or fallow. For best results, add bromoxynil containing herbicides to the tank at 3/16 to 3/8 pound active ingredient per acre (such as Bronate/Maestro MA/Buctril/Maestro 2EC at 3/4 to 1 1/2 pint per acre).

Read and follow all label instructions on timing, precautions, and warnings for these herbicides before using these tank mixtures. Follow the most restrictive labeling. Tank mixes of this product plus Buctril/Maestro 2EC may result in reduced control of Canada thistle.

## **WITH STARANE®/COMET®, STARANE®+SWORD®, STARANE+SALVOTM OR STARANE NXT**

This product may be tank mixed with fluroxypyr containing herbicides registered for use on wheat, barley or fallow. For improved control of kochia (2 to 4" tall), Russian thistle, mustard species, and wild buckwheat this product may be tank mixed with 1/3 to 1-1/3 pints per acre of Starane/Comet, 2/3 to 2-2/3 pints per acre of Starane+Salvo or 3/4 to 2-3/4 pints per acre of Starane+Sword. Additional 2,4-D or MCPA can be added based on local recommendations (refer to 2,4-D and MCPA labels for maximum amount that can be applied to the crop).

For improved control of kochia less than 2" tall this product may be used in combination with Starane NXT at 10 to 14 fluid ounces per acre, for improved control of kochia 2 to 4" tall this product may be used in combination with Starane NXT at 14 to 21 fluid ounces per acre. Add 1 to 2 pints NIS per 100 gallons of spray solution in tank mixes of Starane NXT with this product (See the "SPRAY ADJUVANTS" section of this label for additional information).

Refer to the Starane/Comet, Starane+Salvo, Starane+Sword or Starane NXT label for information regarding use restrictions, labeled crops, rotational cropping recommendations, sprayer cleanup, use precautions and other information. The most restrictive provisions on any label will apply. Do not use the tank mix if any restrictions on the labels conflict with instructions on this product label.

## **WITH CLEANWAVE® HERBICIDE**

For improved control of kochia and other broadleaf weeds in wheat (including durum) this product may be tank mixed with CleanWave herbicide. Add 1 to 2 pints of NIS per 100 gallons of spray solution. Refer to the tank mix product label for information regarding use restrictions, labeled crops, rotational cropping recommendations, sprayer cleanup, use precautions and other information. The most restrictive provisions on any label will apply. Do not use the tank mix if any restrictions on the labels conflict with instruction on this product label.

## **WITH WIDEMATCH®, COLT®, TRUSLATE® OR OTHER CLOPYRALID + FLUROXYPYR CONTAINING HERBICIDES**

For improved control of kochia (2 to 4" tall) and Canada thistle this product may be tank mixed with clopyralid + fluroxypyr containing herbicides in wheat, durum wheat, barley and oats. Additional 2,4-D or MCPA can be added for enhanced control of resistant Russian thistle or kochia and large mustard species. Add 1 to 2 pints of NIS per 100 gallons of spray solution. Refer to the tank mix product label for information regarding use restrictions, labeled crops, rotational cropping recommendations, sprayer cleanup, use precautions and other information. The most restrictive provisions on any label will apply. Do not use the tank mix if any restrictions on the labels conflict with instruction on this product label.

## **With Axial Star, or Rimfire® herbicides**

For improved control of wild oats and other grasses, 0.4 to 0.8 ounces per acre of this product may be tank mixed with Axial Star, or Rimfire in wheat and barley. Add 1 to 4 pints NIS per 100 gallons of spray solution in tank mixes of Rimfire with this product (see SPRAY ADJUVANTS). Refer to Axial Star label for specific adjuvant instructions. Read and follow all label instructions on use restrictions, labeled crops, rotational cropping intervals, sprayer cleanup, use precautions and other information. The most restrictive provisions on any label will apply.



## **WITH HUSKIE®**

For improved control of kochia and other broadleaf weeds this product may be tank mixed with Huskie herbicide in wheat, durum wheat, barley and triticale. No additional surfactant is required when tank mixing with Huskie herbicide. Refer to the Huskie product label for information regarding use restrictions, labeled crops, rotational cropping recommendations, sprayer cleanup, use precautions and other information. The most restrictive provisions on any label will apply. Do not use the tank mix if any restrictions on the labels conflict with instruction on this product label.

## **WITH WOLVERINE®**

This product may be tank mixed with «Wolverine» at 27.2 fluid ounces per acre in wheat, durum, or barley for control of broadleaf weeds, including kochia (less than 2" in height). The most restrictive provisions on any label will apply. Do not use the tank mix if any restrictions on the labels conflict with instruction on this product label.

## **WITH HOELON® HERBICIDE**

This product may be used in combination with Hoelon 3EC and Buctril/Maestro 2EC herbicides in accordance with the Hoelon 3EC label. For best results, use the three-way tank mix of this product at 0.4 ounce per acre plus Hoelon 3EC at 2 2/3 pint per acre plus Buctril/Maestro 2EC at 1-1/2 pints per acre. Apply only to winter wheat. This tank mix should only be used under good soil conditions when wild oat is in the 1- to 4-leaf stage. If conditions are not ideal for the performance of Hoelon 3EC, wild oat control may be reduced. Be sure to follow all warnings and cautions on the Hoelon 3EC and Buctril/Maestro 2EC labels.]

## **WITH EVEREST® 2.0 HERBICIDE**

This product may be used in combination with Everest 2.0 herbicide for broadspectrum grass and broadleaf control in wheat and durum wheat only. Tank mixing this product with Everest 2.0 herbicide will provide additional broadleaf weed control and increase the activity on grassy weeds like yellow foxtail. Consult the Everest 2.0 label for use rates and instruction. Additional broadleaf tank mix partners are required to control ALS resistant weeds. Additional tank mix partners include 2,4-D ester, MCPA ester, bromoxynil products (including Huskie) and Widematch (clopyralid+fluroxypyr). Add NIS at 2-4 pints per 100 gallons of spray solution plus ammonium nitrogen fertilizer or use a basic blend adjuvant at 1% v/v. Refer to the tank mix product label for information regarding use restrictions, labeled crops, rotational cropping recommendations, sprayer cleanup, use precautions and other information. The most restrictive provisions on any label will apply. Do not use the tank mix if any restrictions on the labels conflict with instruction on this product label.

## **WITH RAZE® HERBICIDE**

This product may be used in combination with Raze herbicide for broadspectrum grass and broadleaf control in wheat and durum wheat only. Tank mixing this product with Raze herbicide will provide additional broadleaf weed control and increase the activity on grassy weeds like yellow foxtail. Consult the Raze label for use rates and instruction. 2,4-D ester, MCPA ester or bromoxynil can be tank mixed for additional broadleaf control. Add NIS at 2-4 pints per 100 gallons of spray solution plus ammonium nitrogen fertilizer or use a basic blend adjuvant at 1% v/v. Refer to the tank mix product label for information regarding use restrictions, labeled crops, rotational cropping recommendations, sprayer cleanup, use precautions and other information. The most restrictive provisions on any label will apply. Do not use the tank mix if any restrictions on the labels conflict with instruction on this product label.

## **WITH AXIAL XL HERBICIDE**

This product may be used in combination with Axial XL herbicide for broadspectrum grass and broadleaf control in wheat (not including durum wheat) and barley only. Tank mixing this product with Axial XL herbicide will provide additional broadleaf weed control and control wild oat, yellow foxtail and Italian ryegrass as well as suppression of green foxtail. Consult the Axial XL label for use rates and instruction. No additional surfactant is required when mixing with Axial XL herbicide. Tank mixing additional broadleaf herbicides may reduce grass control. Refer to the tank mix product label for information regarding use restrictions, labeled crops, rotational cropping recommendations, sprayer cleanup, use precautions and other information. The most restrictive provisions on any label will apply. Do not use the tank mix if any restrictions on the labels conflict with instruction on this product label.

## **WITH ASSERT® HERBICIDE**

This product can be tank mixed with Assert. When tank mixing this product with Assert, always include another broadleaf weed herbicide with a different mode of action (for example: 2,4-D ester, MCPA ester, Buctril/Maestro 2EC, or Bronate/Maestro MA). Tank-mixed applications of this product plus Assert may cause temporary crop discoloration, stunting, or injury when heavy rainfall occurs shortly after application.

## **WITH NEXTSTEP® NG HERBICIDE**

This product can be tank mixed with 12.8 -16 fl oz/A of NEXTSTEP NG herbicide for control of wild oat and broadleaf weeds. Green and yellow foxtail control from NEXTSTEP NG will be reduced. No additional surfactant is required when mixing with NEXTSTEP NG herbicide. Refer to the tank mix product label for information regarding use restrictions, labeled crops, rotational cropping recommendations, sprayer cleanup, use precautions and other information. The most restrictive provisions on any label will apply. Do not use the tank mix if any restrictions on the labels conflict with instruction on this product label.

## **WITH OTHER GRASS CONTROL PRODUCTS**

Tank mixtures of this product and grass control products not listed above may result in poor grass control. Nufarm recommends that you first consult the grass control product label, your state experiment station, university, or extension agent, Agricultural dealer, or Nufarm representative as to the potential for antagonism before using the mixture. If no information is available, limit the initial use of this product and the grass product to a small area.

Do not tank mix with Achieve Herbicide.

## **WITH INSECTICIDES OR FUNGICIDES**

This product may be tank mixed or used sequentially with insecticides (or fungicides) registered for use on cereal grains. However, under certain conditions (drought stress, or if the crop is in the 2- to 4-leaf stage), tank mixes or sequential applications of this product with organophosphate insecticides (such as parathion) may produce temporary crop yellowing or, in severe cases, crop injury. Test these mixtures in a small area before treating large areas. However, review all insecticide and fungicide labels for restrictions.

**Do not use this product plus Malathion**, as crop injury will result.

## **WITH LIQUID NITROGEN SOLUTION FERTILIZER**

Liquid nitrogen fertilizer solutions may be used as a carrier in place of water. Run a tank mix compatibility test before mixing this product in fertilizer solution. Do not add this product directly to liquid nitrogen fertilizer; the granules will not dissolve. This product must be thoroughly mixed with clean water before it is added to liquid nitrogen fertilizer. If granules remain when the mixture is poured out, add more clean water and mix until all granules have disappeared. Ensure that the agitator is running when this product is added. Use of this mixture may result in temporary crop yellowing and stunting.

If using low rates of liquid nitrogen fertilizer in the spray solution (less than 50% of the spray solution volume), the addition of surfactant is necessary. Add surfactant at 1/4 to 1 quart per 100 gallon of spray solution (0.06 to 0.25% v/v) based on local recommendations. When using high rates of liquid nitrogen fertilizer solution in the spray solution, adding surfactant increases the risk of crop injury. Consult your agricultural dealer, consultant, fieldman, or Nufarm representative for a specific recommendation before adding an adjuvant to these tank mixtures.

If 2,4-D or MCPA is included with this product and fertilizer mixture, ester formulations tend to be more compatible (see manufacturer's label). Additional surfactant is not needed when using this product in tank mix with 2,4-D ester or MCPA ester and liquid nitrogen fertilizer solutions.

Do not use low rates of liquid nitrogen fertilizer solution as a substitute for a surfactant.

Do not use with liquid fertilizer solutions with a pH less than 3.0.

## **TANK MIXTURES IN FALLOW**

This product may be used as a fallow treatment, and may be tank mixed with other herbicides that are registered for use in fallow.

Read and follow all manufacturer's label instructions for the companion herbicide. If those instructions conflict with this label, do not tank mix the herbicide with this product.

## **TANK MIXTURES IN PRE-PLANT OR CROP PREEMERGENCE BURNDOWN**

This product may be used as a pre-plant burndown or crop preemergence treatment alone or tank mixed with other herbicides that are registered for use as a pre-plant burndown or crop preemergence product, such as glyphosate plus 2,4-D (ester formulations work best) or glyphosate plus dicamba or Pre-Pare® Herbicide.

## **WITH PRE-PARE HERBICIDE**

This product may be used in combination with Pre-Pare herbicide and glyphosate herbicide for broadspectrum grass and broadleaf burndown activity in wheat (not including durum wheat) only. Tank mixing this product with Pre-Pare herbicide will provide additional broadleaf weed control and grassy weed residual activity. Consult the Pre-Pare herbicide label for use rates and instruction. Refer to the tank mix product label for information regarding use restrictions, labeled crops, rotational cropping recommendations, sprayer cleanup, use precautions and other information. The most restrictive provisions on any label will apply. Do not use the tank mix if any restrictions on the labels conflict with instruction on this product label.

Read and follow all manufacturer's label instructions for the companion herbicide. If those instructions conflict with this label, follow the most restrictive labeling (such as planting interval after application), or do not tank mix the herbicide with this product.

## **TANK MIXTURES IN POST HARVEST APPLICATIONS**

This product may be used as a post harvest treatment to crop stubble, and may be tank mixed with other herbicides that are registered for use in fallow.

## MIXING INSTRUCTIONS

Do not use with spray additives that alter the pH of the spray solution below pH 5.0 or above pH 9.0 as rapid product degradation can occur. Spray solutions of pH 6.0 to 8.0 allow for optimum stability of this product.

1. Fill the tank 1/4 to 1/3 full of water.
2. While agitating, add the required amount of this product
3. Continue agitation until this product is fully dissolved, at least 5 minutes.
4. Once this product is fully dissolved, maintain agitation and continue filling tank with water.
5. As the tank is filling, add tank mix partners and then add the required volume of spray adjuvant. Always add spray adjuvant last.  
Antifoaming agents may be used. Do not use with spray additives that alter the pH of the spray solution below pH 6.0 as rapid product degradation can occur. Spray solutions of pH 7.0 and higher allow for optimum stability of this product.
6. Dispersed tank mix partners can settle if the tank mixture is not continually agitated. If settling occurs, thoroughly re-agitate before using.
7. Apply this product spray mixture within 24 hours of mixing to avoid product degradation.
8. If this product and a tank mix partner are to be applied in multiple loads, fully dissolve this product in clean water prior to adding to the tank.

## GRAZING

Allow at least 7 days between application and grazing of treated forage and 7 days between application and feeding of treated forage to livestock. Allow at least 30 days between application and feeding of hay from treated areas to livestock. Harvested straw may be used for bedding or feed. Allow at least 45 days between application and harvesting grain.

## SPRAY EQUIPMENT

For specific application equipment, refer to the manufacturer's recommendations for additional information on GPA, pressure, speed, nozzle types and arrangements, nozzle heights above the target canopy, etc.

Be sure to calibrate air or ground equipment properly before application. Select a spray volume and delivery system that will ensure thorough coverage and a uniform spray pattern with minimum drift. Use higher spray volumes to obtain better coverage when crop canopy is dense. Avoid swath overlapping, and shut off spray booms while starting, turning, slowing, or stopping, to avoid injury to the crop. Do not make applications using equipment and/or spray volumes or during weather conditions that might cause spray to drift onto nontarget sites. For additional information on spray drift refer to Spray Drift Management section of label.

Continuous agitation may be required to keep this product and tank-mix partners in solution or suspension. Refer to tank-mix partner labels for additional information.

## SPRAYER CLEANUP

The spray equipment must be cleaned before this product is sprayed. Follow the cleanup procedures specified on the labels of the previously applied products. If no directions are provided, follow the six steps outlined in "After Spraying" in this label.

### SPRAYER CLEANUP FOR MULTIPLE LOAD SPRAYING

It is recommended that during periods when multiple loads of this product are applied, at the end of each day of spraying, the interior of the tank be rinsed with fresh water and then partially filled, and the boom and hoses flushed. This will prevent the buildup of dried pesticide deposits which can accumulate in the application equipment.

### AFTER SPRAYING AND BEFORE SPRAYING CROPS OTHER THAN WHEAT AND BARLEY

To avoid subsequent injury to desirable crops, thoroughly clean all mixing and spray equipment immediately following applications of this product as follows:

1. Drain tank; thoroughly rinse spray tanks, boom, and hoses with clean water. Loosen and physically remove any visible deposits.
2. Fill the tank with clean water and 1 gallon of household ammonia\* (contains 3% active) for every 100 gal of water. Flush the hoses, boom, and nozzles with the cleaning solution. Then add more water to completely fill the tank. Circulate the cleaning solution through the tank and hoses for at least 15 minutes. Flush the hoses, boom, and nozzles again with the cleaning solution, and then drain the tank.
3. Remove the nozzles and screens and clean separately in a bucket containing cleaning agent and water.
4. Repeat Step 2.
5. Rinse the tank, boom, and hoses with clean water.
6. If only Ammonia is used as a cleaner, the rinsate solution may be applied back to the crop(s) recommended on this label. Do not exceed the maximum labeled use rate. If other cleaners are used, consult the cleaner label for rinsate disposal instructions. If no instructions are given, dispose of the rinsate on site or at an approved waste disposal facility.

\* Equivalent amounts of an alternate-strength ammonia solution can be used in the cleanout procedure. Carefully read and follow the individual cleaner instructions. Consult your Ag dealer for a listing of approved cleaners.

## Notes:

1. **CAUTION:** Do not use chlorine bleach with ammonia as dangerous gases will form. Do not clean equipment in an enclosed area.
2. Steam-cleaning aerial spray tanks is recommended prior to performing the above cleanout procedure to facilitate the removal of any caked deposits.
3. When this product is tank mixed with other pesticides, all cleanout procedures should be examined and the most rigorous procedure should be followed.
4. In addition to this cleanout procedure, all precleanout guidelines on subsequently applied products should be followed as per the individual labels.
5. Where routine spraying practices include shared equipment frequently being switched between applications of this product and applications of other pesticides to sensitive crops during the same spray season, it is recommended that a sprayer be dedicated to this product to further reduce the chance of crop injury.

## SPRAY DRIFT MANAGEMENT

The interaction of many equipment and weather-related factors determines the potential for spray drift. The applicator is responsible for considering all these factors when making application decisions.

## IMPORTANCE OF DROPLET SIZE

AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. The presence of sensitive species nearby, the environmental conditions, and pest pressure may affect how an applicator balances drift control and coverage. **APPLYING LARGER DROPLETS REDUCES DRIFT POTENTIAL, BUT WILL NOT PREVENT DRIFT IF APPLICATIONS ARE MADE IMPROPERLY OR UNDER UNFAVORABLE ENVIRONMENTAL CONDITIONS!** See **Wind, Temperature and Humidity**, and **Temperature Inversions** sections of this label.

## CONTROLLING DROPLET SIZE - GENERAL TECHNIQUES

- **Volume** - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- **Pressure** - Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. **WHEN HIGHER FLOW RATES ARE NEEDED, USE A HIGHER-CAPACITY NOZZLE INSTEAD OF INCREASING PRESSURE.**
- **Nozzle Type** - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles.

## CONTROLLING DROPLET SIZE - AIRCRAFT

- **Number of Nozzles** - Use the minimum number of nozzles with the highest flow rate that provide uniform coverage.
- **Nozzle Orientation** - Orienting nozzles so that the spray is emitted backwards, parallel to the airstream will produce larger droplets than other orientations.
- **Nozzle Type** - Solid stream nozzles (such as disc and core with swirl plate removed) oriented straight back produce larger droplets than other nozzle types.
- **Boom Length** - The boom length should not exceed 3/4 of the wing or rotor length - longer booms increase drift potential.
- **Application Height** - Application more than 10 feet above the canopy increases the potential for spray drift.

## BOOM HEIGHT

Setting the boom at the lowest referenced height (if specified) which provides uniform coverage reduces the exposure of droplets to evaporation and wind. For ground equipment, the boom should remain level with the crop and have minimal bounce.

## WIND

Drift potential increases at wind speeds of less than 3 mph (due to inversion potential) or more than 10 mph. However, many factors, including droplet size and equipment type, determine drift potential at any given wind speed. **AVOID GUSTY AND WINDLESS CONDITIONS.**

**Note:** Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

## TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, set up equipment to produce larger droplets to reduce effects of evaporation.

## TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog. However, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

## SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce the effects of wind. However, it is the responsibility of the applicator to verify that the shields are preventing drift and not interfering with uniform deposition of the product.

## AIR ASSISTED (AIR BLAST) FIELD CROP SPRAYERS

Air assisted field crop sprayers carry droplets to the target via a downward directed airstream. Some may reduce the potential for drift, but if a sprayer is unsuitable for the application and/or set up improperly, high drift potential can result. It is the responsibility of the applicator to determine that a sprayer is suitable for the intended application, is configured properly, and that drift is not occurring.

**Note:** Air assisted field sprayers can affect product performance by affecting spray coverage and canopy penetration. Consult the spray equipment section of this label to determine if use of an air assist sprayer is recommended.

## RESISTANCE

This product contains the active ingredients thifensulfuron and tribenuron is a Group 2 herbicide based on the mode of action classification system of the Weed Science Society of America. When herbicides with mode of action classifications that affect the same biological sites of action are used repeatedly over several years to control the same weed species in the same treatment area, naturally-occurring resistant biotypes may survive a correctly applied herbicide treatment, propagate, and become dominant in that area. Adequate control of these resistant weed biotypes cannot be expected. If weed control is unsatisfactory, it may be necessary to retreat the problem area using a product affecting a different biological site of action.

To better manage herbicide resistance through delaying the proliferation and possible dominance of herbicide resistant weed biotypes, it may be necessary to change cultural practices within and between crop seasons such as using a combination of tillage, retreatment, tank-mix partners and/or sequential herbicide applications that affect a different site of action. Weed escapes that are allowed to go to seed, and movement of plant material between treatment areas on equipment will promote the spread of resistant biotypes.

It is advisable to keep accurate records of pesticides applied to individual fields to help obtain information on the spread and dispersal of resistant biotypes. Consult your agricultural dealer, consultant, applicator, and/or appropriate state agricultural extension service representative to determine appropriate actions for treating specific resistant weed biotypes in your area.

When herbicides that affect the same biological site of action are used repeatedly over several years to control the same weed species in the same field, naturally-occurring resistant biotypes may survive a correctly applied herbicide treatment, propagate, and become dominant in that field. Adequate control of these resistant weed biotypes cannot be expected. If weed control is unsatisfactory, it may be necessary to retreat the problem area using a product affecting a different site of action.

## INTEGRATED PEST MANAGEMENT

**Nufarm recommends the use of Integrated Pest Management (IPM) programs to control pests. This product may be used as part of an Integrated Pest Management (IPM) program that can include biological, cultural, and genetic practices aimed at preventing economic pest damage. Application of this product should be based on IPM principles and practices include field scouting or other detection methods, correct target pest identification, population monitoring, and treating when target pest populations reach locally determined action thresholds.**

**Consult your state cooperative extension service, professional consultants or other qualified authorities to determine appropriate action treatment threshold levels for treating specific pest/crop systems in your area.**

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Store product in original container only. Store in a cool, dry place.

**PESTICIDE DISPOSAL:** Waste resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

### CONTAINER HANDLING:

**Nonrefillable Plastic and Metal Containers (Capacity Equal to or Less Than 50 Pounds):** Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes. In the event of a major spill, fire or other emergency contact CHEMTREC 1-800-424-9300.

## WARRANTY DISCLAIMER

**The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.**

## LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

**If you do not agree with or do not accept any of the directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.**

RV121514 [1]

Rapport, Diabolo, Clash, Maestro, Comet, TruSlate and Assert are a registered trademark of Nufarm Americas Inc.

Everest and Raze are registered trademarks of Arysta LifeScience Corporation.

Banvel and Clarity are trademarks or registered trademarks of BASF Corporation.

Bronate, Buctril, Hoelon, Huskie, Rimfire and Wolverine are registered trademarks of Bayer CropScience.

CleanWave, GoldSky, Starane and Widematch are registered trademarks of Dow AgroSciences, LLC.

Colt, Saber, Salvo and Sword are trademarks or registered trademarks of UAP Loveland Products Inc.

Raindrop RA is a registered trademark of Delavan.