



# Tide Paclo 2SC

Plant Growth Regulator for Turfgrass, Trees, Shrubs & Vines

**ACTIVE INGREDIENT:**

Pacllobutrazol: ((±)-(R\*,R\*)-β-[(4-chlorophenyl)methyl]-α-(1,1-dimethylethyl)

-1H-1,2,4-triazole-1-ethanol)..... 22.3%

**OTHER INGREDIENTS:**..... 77.7%

**TOTAL:**..... 100.0%

Tide Paclo 2SC contains 2 lbs. active ingredient per gallon.

**KEEP OUT OF REACH OF CHILDREN**

## CAUTION

See inside label booklet for First Aid, additional Precautionary Statements and Directions for Use including Storage and Disposal instructions.

**Net Contents: 1 Gallon**

EPA Reg. No.: 80697-4

EPA Est. No.: □ 69845-CHN-002 □ 44616-MO-001

□ 70253-CA-002 □ 71764-NC-001

□ 39578-TX-001 □ 37429-GA-002

□ 70815-GA-001 □ 83411-MN-001

□ 69016-MS-001 □ 5905-IA-001

Manufactured for:  
Zhejiang Tide CropScience Co., Ltd.  
21 Hubble, Irvine, CA 92618

**SHAKE WELL BEFORE USE**

### PROOF

Proof date: 01/09/2023

Customer: Tide USA

Job number: ZTC-PGR806974

Label size: 149 x 165mm

Leaflet flat size: 286x 165mm

Leaflet folded size: 151 x 165mm

Label colors: PANTONE Process Black C,  
PANTONE 3282 C,

Leaflet "in" colors: Black

Leaflet "out" colors: PANTONE Process Black C,  
PANTONE 3282 C,

PEEL BACK BOOK HERE AND RESEAL AFTER OPENING



# Tide Paclo 2SC

Plant Growth Regulator for Turfgrass, Trees, Shrubs & Vines

**ACTIVE INGREDIENT:**

Paclobutrazol: ((±)-(R\*,R\*)-β-[(4-chlorophenyl)methyl]-α-(1,1-dimethylethyl)-1H-1,2,4-triazole-1-ethanol).....22.3%

**OTHER INGREDIENTS:**.....77.7%

**TOTAL:**.....100.0%

Tide Paclo 2SC contains 2 lbs. active ingredient per gallon.

**KEEP OUT OF REACH OF CHILDREN**

## CAUTION

See inside label booklet for First Aid, additional Precautionary Statements and Directions for Use including Storage and Disposal instructions.

**Net Contents: 1 Gallon**

EPA Reg. No.: 80697-4

- EPA Est. No.:  69845-CHN-002     44616-MO-001
- 70253-CA-002             71764-NC-001
- 39578-TX-001             37429-GA-002
- 70815-GA-001             83411-MN-001
- 69016-MS-001             5905-IA-001

Manufactured for:  
Zhejiang Tide CropScience Co., Ltd.  
21 Hubble, Irvine, CA 92618

**SHAKE WELL BEFORE USE**

PEEL BACK BOOK HERE AND RESEAL AFTER OPENING

## FIRST AID

### If swallowed:

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to by a poison control center or doctor.
- Do not give anything to an unconscious person.

### If on skin or clothing:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

### If in eyes:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

## HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact CHEMTREC at 1-800-424-9300 for emergency medical treatment information.

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

## CAUTION

Harmful if swallowed or absorbed through the skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Wear long-sleeved shirt and long pants, socks, shoes, and waterproof gloves. Remove and wash contaminated clothing before reuse.

## PERSONAL PROTECTIVE EQUIPMENT (PPE)

### Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, polyvinyl chloride, or viton
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Wash PPE after each day's use.

## USER SAFETY RECOMMENDATIONS

### Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

## ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters.

## PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

---

## DIRECTIONS FOR USE

---

### **Failure to follow directions and precautions on this label may result in plant injury or poor disease control.**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. Read all label directions carefully before use.

**Do not** apply this product when wind speed favors drift beyond the area intended for treatment.

**Do not** use on residential lawns.

### **AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, polyvinyl chloride, or viton
- Shoes plus socks

---

## GENERAL INFORMATION

---

Tide Paclo 2SC is a systemic plant growth regulator that slows vegetative growth by inhibiting gibberellin biosynthesis for use on turfgrass, trees, shrubs, vines, and landscaping plants.

For turf, when Tide Paclo 2SC is applied as directed, vertical growth of turf will be slowed within 3-10 days, resulting in reduced mowing frequency for a 6 to 8 week period on St. Augustine grass and Kentucky bluegrass/perennial ryegrass turf areas, as well as established bent grass, hybrid bermuda grass and perennial ryegrass tees, fairways and roughs.

Following Tide Paclo 2SC applications, turf will gradually undergo increased greening and density, which may persist up to 12 weeks.

Tide Paclo 2SC can be used on turfgrass in the following areas: golf courses, (tees, fairways, greens and rough), athletic fields, commercial and industrial areas, parks, and boulevards.

For trees, Tide Paclo 2SC is a xylem mobile plant regulator. It reduces the above ground vegetative growth and changes specific morphological characteristics of the treated plant. It slows vegetative growth as well as creating other physiological effects by inhibiting gibberellin biosynthesis. Slowed vegetative growth reduces the volume of woody growth that must be removed when trimming occurs and may extend the trimming cycle.

Tide Paclo 2SC is most effective when applied to the soil near the base of the tree, shrub, or vine either by soil injection or with basal soil dress.

Tide Paclo 2SC may be used on listed tree species in residential areas, utility rights-of-way (e.g. trees below transmission lines, distribution lines and feeder lines; and railroad and highway rights-of-way), other urban settings (e.g. parks, parking lot areas, storage areas, street medians and easements, commercial/municipal buildings or office sites, plazas, around walkways and sidewalks, above ground planters, and around schools, hospitals and other buildings), and other non-crop

areas (e.g. industrial sites, athletic fields and other recreational areas, naturalized areas and other uncultivated non-agricultural areas, fence and property lines, and airports).

## NON-RESIDENTIAL TURF

Tide Paclo 2SC, a specialty plant growth regulator that slows vertical grass growth and reduces mowing for up to two (2) months after application. The frequency of mowing can be reduced by up to 50% during the period of effective retardation. Use of Tide Paclo 2SC on fine turf should be accompanied by moderate-to-high fertility to maintain turfgrass appearance and reduce discoloration.

Tide Paclo 2SC can be used on turfgrass in the following areas: golf courses, (tees, fairways, greens and rough), athletic fields, commercial and industrial areas, parks, and boulevards.

### USE DIRECTIONS

| Turf   | Application Rate<br>(Product amount/a.i.-lbs per acre)   | Potential Effects  |
|--|--|--|
| St Augustine, Bermuda & similar warm season grasses                                | 0.5 lbs. a.i.<br>32 fl. oz prod/A (Sandy soil)<br>0.75 lbs a.i.<br>48 fl. oz prod/A (Clay soil)                        | Reduced mowing for 6-8 weeks on established hybrid bermudagrass, bent grass and perennial ryegrass fairways, tees and roughs and on St. Augustinegrass and Kentucky bluegrass/perennial ryegrass turf. |
| Bluegrass, Rye, Fescue, Bent & similar cool season grasses (except putting greens) | 0.25 to 0.50 lbs a.i.<br>16 -32 fl. oz prod/A<br><br>Reduce by 50% if <i>Poa annua</i> is a major turfgrass population | Following application, turf will gradually increase in density with tighter knit areas of turfgrass. This will result in a reduced potential for scalping of all turfgrass areas.                      |
| Bentgrass Putting Greens & Overseeded Bermuda                                      | 0.10 to 0.25 lbs a.i.<br>6.4 -16 fl. oz prod/A   | Suppression of <i>Poa annua</i> by reducing its growth and its compatibility, leading to selective control after prolonged use.  |

Apply Tide Paclo 2SC with standard pressurized application equipment with by-pass or mechanical agitation using 50 mesh strainer to turf in sufficient amount of water (minimum 1 gal water /1,000 sq ft. = 43.5 gals/Ac) to ensure spray coverage uniformly to turf and to avoid skipping and/or overlapping.

For best results, use 2 to 5 gals water /1,000 sq ft.

After application of Tide Paclo 2SC, water-in within 24 hours to limit surface movement, but not to the point of runoff. To prevent product runoff, time applications to allow for watering-in and maximum absorption into treated turf prior to a rain event.

Tide Paclo 2SC can be used any time when established hybrid bermudagrass and St. Augustinegrass are green, actively growing and have recovered from dormancy after winter.

Tide Paclo 2SC can be applied in spring after green-up and after turf has been mowed once or twice. Apply at least 1 month before onset of high air temperatures. In late summer/early fall, apply at least 1 month before anticipated first killing frost.

Apply any time after overseeded turf has established itself successfully. Do not apply after March 15<sup>th</sup> to avoid delay in Bermudagrass green-up.

Maintain moderate soil moisture conditions before and after each application to achieve best plant growth retardation effect.

A repeat application of Tide Paclo 2SC within the same growing season may be made 8 weeks apart for growth regulation & 4 to 6 weeks apart for color, quality enhancement after the initial application. Do not apply more than 4 quarts per acre per year (2 lbs. a.i./A).

Tide Paclo 2SC will not prevent seedhead production if used alone. A tank mixture for a spray program using a seed head control agent is required when seedhead visibility is a problem. If a seed head control agent is used for *Poa annua*, apply Tide Paclo 2SC at least 14 days after application of the seed head control agent.

### Weeds, Insects and Diseases:

If crabgrass or other annual grassy weeds have been a problem in the past, an application of an appropriate preemergence weed control product should be made before the use of Tide Paclo 2SC. Space applications of Tide Paclo 2SC and preemergence product (Goosegrass/Crabgrass Preventer) at least 4 weeks apart.

Broadleaf weeds are not significantly affected by Tide Paclo 2SC. To control the growth of weeds, treat with an appropriate herbicide when weeds are actively growing. Carefully follow label directions.

If a weed, disease, or insect problem occurs after application of Tide Paclo 2SC, apply control product at the recommended rate. Tide Paclo 2SC is compatible with most existing control products. A compatibility test should be conducted with product(s) with which you do not have prior experience.

### Regulator response and Turf cultivar

Excessive irrigation and/or nitrogen fertilization may shorten period of growth regulation. Regulator response will vary somewhat according to turf variety. For St. Augustinegrass, Bitter Blue will be the most responsive and Floratam the least responsive.

### HYBRID BERMUDAGRASS

| Sensitivity/Activity | Cultivar         | Period of Growth Regulation | Period of Color Response |
|----------------------|------------------|-----------------------------|--------------------------|
| Medium/Good          | Tifway I, II     | 5 - 6 weeks                 | 6 - 8 weeks              |
| High/Excellent       | Tifgreen, Ormond | 6 - 8 weeks                 | 8 - 10 weeks             |

### ST. AUGUSTINEGRASS

| Sensitivity/Activity | Cultivar                          | Period of Growth Regulation | Period of Color Response |
|----------------------|-----------------------------------|-----------------------------|--------------------------|
| Low/Moderate         | Floratam, Floralawn               | 4 - 6 weeks                 | 5 - 8 weeks              |
| Medium/Good          | Floratine, Raleigh, Texas, Common | 6 - 7 weeks                 | 7 - 8 weeks              |
| High/Excellent       | Bitter Blue, Seville              | 7 - 8 weeks                 | 8 - 10 weeks             |

### USE AND RESTRICTIONS

**Do not** use on residential lawns.

**Do not** apply Tide Paclo 2SC to turf within right-of-ways.

**Do not** use on bermudagrass putting greens except for winter overseeding enhancement use.

**Do not** use on athletic fields under heavy traffic where maximum growth potential of turf is desired.

Applications to turf areas under trees will not affect/harm trees.

**Do not** use during periods of extreme dry or cold weather conditions, or during heavy insect or disease activity.

**Do not** apply product when soil is already saturated. Heavy rainfall or irrigation in the treated areas may cause active ingredient to move laterally on slopes and collect in low areas. These areas may undergo more severe growth control for a longer period of time.

**Do not** use on areas containing greater than 70% *Poa annua*, since discoloration of *Poa* may be unacceptable.

Delay treatment of newly-sodded or sprigged turf until grass has knitted down and rooted firmly.

Delay sprigging for at least 4 weeks and sodding at least 2 weeks after application is made.

Withhold application on large turf areas that have been thinned from winter drainage, disease, or insects until desired fill-in is achieved.

Assure that dosage rates are measured accurately since rates greater than those recommended may cause undesirable turf growth control and may discolor areas temporarily.

**Do not** use on areas to be cultivated for food or food crops or to be resown with grasses within two years of treatment.

- Do not** apply more than 4 qts. per acre per year (2 lbs. a.i./A).
- Do not** graze treated areas or harvest for forage or hay.
- Do not** apply this product through any type of irrigation equipment.

---

## TREES

---

Tide Paclo 2SC as tree growth regulator is a xylem mobile plant growth regulator that slows vegetative growth by inhibiting gibberellin biosynthesis. Tide Paclo 2SC reduces vegetative growth thereby reducing the volume of woody growth that must be removed from trees when trimming occurs and may extend the trimming cycle. Tide Paclo 2SC is equally effective when applied as a basal soil drench or by soil injection.

Use Tide Paclo 2SC on listed tree species in residential areas, utility rights-of-way (e.g. trees below transmission lines, distribution lines and feeder lines; and railroad and highway rights-of-way), other urban settings (e.g. parks, parking lot areas, storage areas, street medians and easements, commercial/municipal building or office sites, plazas, around walkways and sidewalks, above ground planters, and around schools, hospitals and other buildings), and other non-crop areas (e.g. industrial sites, athletic fields and other recreational areas, naturalized areas and other uncultivated non-agricultural areas, fence and property lines, and airports).

### Symptoms of Growth Regulation

Tide Paclo 2SC demonstrates little or no phloem mobility. Activity occurs following root uptake and xylem translocation throughout the tree canopy. Symptoms of growth regulation may not be visible for up to 18 months following application of Tide Paclo 2SC. Initial effects of Tide Paclo 2SC may be observed as intense greening of foliage with no resulting phytotoxicity. At the time when normal vegetative growth extension should occur, trees treated with Tide Paclo 2SC will exhibit shortened internodes which may be somewhat thickened. Smaller leaf size and enhanced flowering may also be observed in some species.

## USE DIRECTIONS

### Application Timing

Applications can be made throughout the year, weather permitting, except when soil is frozen or saturated with water.

**Note:** Tide Paclo 2SC is absorbed by plant roots and translocated to the growing tissues in response to evaporative water loss (transpiration). If applications are made after fall leaf drop, uptake of Tide Paclo 2SC will not occur until development of new leaves in the spring and resumption of transpiration.

### Mixing Directions

Applications are made with a diluted mixture composed of 1 part Tide Paclo 2SC to 11 parts water. Mix 317 ml (10.7 fl oz, 0.167 lbs. a.i.) of Tide Paclo 2SC with water to make one gallon of diluted mixture. Mix one gallon (2 lbs. a.i.) of Tide Paclo 2SC with water to make 12 gallons of diluted mixture. To improve the handling characteristics of the diluted mixture, the addition of a suspension aid is recommended such as, nonionic and organosilicone surfactants.

Follow all label directions and precautions on the product label of the suspension aid.

### Application Methods

Tide Paclo 2SC may be applied as a basal drench or by soil injection. Treat only trees that are well established in their final location. Treatment of trees less than 4 inches in diameter is not recommended.


### Basal Drench

Prior to application, make a 2" deep furrow around the base of the tree near the point of contact between the soil and the tree trunk. Apply the required dose as listed in the table "**Amount of Tide Paclo 2SC Diluted Mixture Required for Specific Application Rates and Tree Sizes**" below. Refer to Figure 1 for placement as a basal drench. Carefully pour the diluted mixture of Tide Paclo 2SC into the furrow with a graduated container/jug or with a handheld hose connected to a truck-mounted tank/hydraulic sprayer. To avoid possible runoff after applying, refill the furrow with untreated soil.


### Soil Injection

The diluted mixture of Tide Paclo 2SC should be injected approximately 6 inches deep. Use soil injection equipment capable of delivery at 100 to 200 psi. Injection orifices should be oriented to release the diluted product horizontally at the point of injection. The required dose should be divided evenly among injection sites spaced as uniformly as possible around the tree

trunk. The injection sites should be positioned to release Tide Paclo 2SC diluted mixture as close as possible to the point of contact between soil and unthickened bark beneath the soil so that the active ingredient may be readily absorbed by the tree (Figure 2). Injection sites should also be located next to buttress roots (Figure 2). For trees less than 6 inches DHB, use at least 4 evenly spaced injection sites per tree.



**Figure 1.** Placement Tide Paclo 2SC as basal drench



**Figure 2.** Placement of Tide Paclo 2SC as a soil injected treatment.



## Application Rate

| Application Rate Ranges for Diluted Mixture of Tide Paclo 2SC<br>for Treatment of Various Tree Species <sup>1</sup>   |  |  |
|---|--|--|
| 1.7 to 3.4 fl oz<br>(50 to 100 ml,<br>0.0022 to 0.0044 lbs a.i.,<br>1 to 2 grams a.i.)<br>per inch DBH  | 3.4 to 5.1 fl oz<br>(100 to 150 ml,<br>0.0044 to 0.0066 lbs a.i.,<br>2 to 3 grams a.i.)<br>per inch DBH  | 5.1 to 6.8 fl oz<br>(150 to 200 ml,<br>0.0066 to 0.0088 lbs. a.i.<br>3 to 4 grams a.i.)<br>per inch DBH  |
| Australian Pine<br>Crape Myrtle<br>Cypress<br>Dogwood<br>Elm (Cedar, Chinese,<br>Siberian)<br>Holly, Yaupon<br>Maple (Amur,<br>Japanese)<br>Redbud<br>Sweetgum<br>Tupelo<br>Zelkova | Australian Bottle<br>Basswood<br>Beech<br>Boxelder<br>Buckeye<br>California Pepper<br>Camphor<br>Catalpa<br>Cherry Laurel<br>Elm, American<br>Hawthorne<br>Hickory<br>Holly, American<br>Horse Chestnut<br>Linden<br>Locust, Honey<br>Maple (Big Leaf,<br>Norway, Red,<br>Silver, Sugar)<br>Mimosa<br>Mountain Ash | Oak (Black, Bur,<br>Live, Pin, Post,<br>Red, Sand<br>Shinnery,<br>Shumard,<br>Water, White,<br>Willow)<br>Oleander<br>Olive, Russian<br>Pecan<br>Persimmon<br>Plum - Ornamental<br>Rosewood<br>Sassafras<br>Sumac - African<br>Walnut<br>Acacia<br>Ailanthus<br>Alder<br>Anaqua<br>Arborvitae<br>Ash<br>Aspen<br>Banyan<br>Birch<br>Bottlebrush<br>Bradford Pear<br>Cedar (Deodora,<br>Eastern Red,<br>Western Red,<br>all others)<br>Cherry (Black,<br>Ornamental)<br>Chinaberry<br>Cottonwood<br>Crabapple<br>Cypress, Leyland<br>Ebony<br>Eucalyptus<br>Fir<br>Ginkgo<br>Gumbo Limbo<br>Hackberry<br>Hemlock<br>Huisache<br>Ironwood<br>Jacaranda<br>Juniper<br>Laurel<br>Locust, Black<br>Magnolia<br>Mahogany<br>Melaleuca<br>Mesquite<br>Mulberry<br>Oak (Blackjack,<br>Laurel, Valley)<br>Olive, Black<br>Osage Orange<br>Palms<br>Palo Verde<br>Paulownia<br>Pines<br>Poinciana<br>Poplar<br>Raintree - Golden<br>Redwood<br>Saltcedar<br>Sea Grape<br>Serviceberry<br>Silk Oak<br>Soapberry<br>Spruce<br>Sugarberry<br>Sycamore<br>Tallow, Chinese<br>Tamarisk<br>Tuliptree<br>Willow<br>Yellow Poplar<br>Yew |

<sup>1</sup>For acidic soils (low pH), soils with high organic matter content or soils with high clay content, use the higher rate, not to exceed 0.0088 lbs. a.i. 4 grams a.i. per inch DBH.

| Amount of Tide Paclo 2SC Diluted Mixture Required for Specific Application Rates and Tree Sizes |  |                 |   |                 |   |                 |   |                 |
|---|--|-----------------|---|-----------------|---|-----------------|---|-----------------|
| Dia. Of Tree at Breast Height (DBH) (inches)  | 1.7 fl oz (50 ml, 0.0022 lbs. a.i.) per inch DBH |                 | 3.4 fl oz (100 ml, 0.0044 lbs. a.i.) per inch DBH |                 | 5.1 fl oz (150 ml, 0.0066 lbs. a.i.) per inch DBH |                 | 6.8 fl oz (200 ml, 0.0088 lbs. a.i.) per inch DBH |                 |
|   | Total fl oz (ml) required                        | Total lbs. a.i. | Total fl oz (ml) required                         | Total lbs. a.i. | Total fl oz (ml) required                         | Total lbs. a.i. | Total fl oz (ml) required                         | Total lbs. a.i. |
| 4   | 6.8 (200)  | 0.0088          | 13.6 (400)  | 0.0176          | 20.4 (600)  | 0.0264          | 27.2 (800)  | 0.0352          |
| 5   | 8.5 (250)  | 0.0110          | 17.0 (500)  | 0.0220          | 25.5 (750)  | 0.0330          | 34.0 (1000)                                       | 0.0440          |
| 6   | 10.2 (300)                                       | 0.0132          | 20.4 (600)  | 0.0264          | 30.6 (900)  | 0.0396          | 40.8 (1200)                                       | 0.0528          |
| 7   | 11.9 (350)                                       | 0.0154          | 23.8 (700)  | 0.0308          | 35.7 (1050)                                       | 0.0462          | 47.6 (1400)                                       | 0.0616          |
| 8   | 13.6 (400)                                       | 0.0176          | 27.2 (800)  | 0.0352          | 40.8 (1200)                                       | 0.0528          | 54.4 (1600)                                       | 0.0704          |
| 9   | 15.3 (450)                                       | 0.0198          | 30.6 (900)  | 0.0396          | 45.9 (1350)                                       | 0.0594          | 61.2 (1800)                                       | 0.0792          |
| 10  | 17.0 (500)                                       | 0.0220          | 34.0 (1000)                                       | 0.0440          | 51.0 (1500)                                       | 0.0660          | 68.0 (2000)                                       | 0.0880          |
| 12  | 20.4 (600)                                       | 0.0264          | 40.8 (1200)                                       | 0.0528          | 61.2 (1800)                                       | 0.0792          | 81.6 (2400)                                       | 0.1056          |
| 14  | 23.8 (700)                                       | 0.0308          | 47.6 (1400)                                       | 0.0616          | 71.4 (2100)                                       | 0.0924          | 95.2 (2800)                                       | 0.1232          |
| 16  | 27.2 (800)                                       | 0.0352          | 54.4 (1600)                                       | 0.0704          | 81.6 (2400)                                       | 0.1056          | 108.8 (3200)                                      | 0.1408          |
| 18  | 30.6 (900)                                       | 0.0396          | 61.2 (1800)                                       | 0.0792          | 91.8 (2700)                                       | 0.1188          | 122.4 (3600)                                      | 0.1584          |
| 20  | 34.0 (1000)                                      | 0.0440          | 68.0 (2000)                                       | 0.0880          | 102.0 (3000)                                      | 0.1320          | 136.0 (4000)                                      | 0.1760          |
| 22  | 37.4 (1100)                                      | 0.0484          | 74.8 (2200)                                       | 0.0968          | 112.2 (3300)                                      | 0.1452          | 149.6 (4400)                                      | 0.1936          |
| 24  | 40.8 (1200)                                      | 0.0528          | 81.6 (2400)                                       | 0.1056          | 122.4 (3600)                                      | 0.1584          | 163.2 (4800)                                      | 0.2112          |
| 26  | 44.2 (1300)                                      | 0.0572          | 88.4 (2600)                                       | 0.1144          | 132.6 (3900)                                      | 0.1716          | 176.8 (5200)                                      | 0.2288          |
| 28  | 47.6 (1400)                                      | 0.0616          | 95.2 (2800)                                       | 0.1232          | 142.8 (4200)                                      | 0.1848          | 190.4 (5600)                                      | 0.2464          |
| 30  | 51.0 (1500)                                      | 0.0660          | 102.0 (3000)                                      | 0.1320          | 153.0 (4500)                                      | 0.1980          | 204.0 (6000)                                      | 0.2640          |
| 32  | 54.4 (1600)                                      | 0.0704          | 108.8 (3200)                                      | 0.1408          | 163.2 (4800)                                      | 0.2112          | 217.6 (6400)                                      | 0.2816          |
| 34  | 57.8 (1700)                                      | 0.0748          | 115.6 (3400)                                      | 0.1496          | 173.4 (5100)                                      | 0.2244          | 231.2 (6800)                                      | 0.2992          |
| 36  | 61.2 (1800)                                      | 0.0792          | 122.4 (3600)                                      | 0.1584          | 183.6 (5400)                                      | 0.2376          | 244.8 (7200)                                      | 0.3168          |

## USE AND RESTRICTIONS

- Apply at prescribed rates and follow safety procedures.
- Trees not used for food production that are not specifically listed on this label may be treated if all other label directions are followed.
- Trees which are severely stressed due to moisture, temperature, low soil fertility, or exhibit mechanical or chemical injury should not be treated.
- Do not reapply Tide Paclo 2SC until symptoms from previous applications begin to subside.
- If soil is heavily compacted or a high water table exists, use of a registered tree growth regulator such as flurprimidol may be more satisfactory.
- Basal drench and soil injection application of Tide Paclo 2SC may result in localized, temporary discoloration of turfgrass immediately adjacent to the treatment site.
- Avoid basal drench applications on slopes or other areas where Tide Paclo 2SC or treated soil may be washed away from the base of the tree by rainfall or irrigation.
- Treatment of trees bordered by shrubs and/or herbaceous ornamentals may cause these plants to be affected if their roots extend into the treatment zone.
- Do not treat sugar maple trees or any other trees that are or could be tapped for sugar within one year of application.
- Do not treat nut or fruit trees that will be harvested within one year.

**Chemigation:** Do not apply this product through any type of irrigation system.

## SHRUBS AND VINES

On woody shrubs, Tide Paclo 2SC reduces vegetative growth and creates beneficial changes including increased root hair growth and darker, green foliage. Soil applications on woody shrubs show 90-95% growth control for 1-2 years, depending on the length of the growing season, sunlight exposure, soil texture, rainfall and temperature conditions. Tide Paclo 2SC is equally effective when applied as a basal drench or by soil injection.

Foliar sprays of Tide Paclo 2SC on shrubs, vines, and specified landscape plants typically provide up to 3-6 months of activity. Length of control with foliar sprays is influenced by application timing, amount of pruning performed, soil type, growing conditions and plant species.

### USE DIRECTIONS


#### Soil Injection/Basal Drench

Soil injections or basal drenches of Tide Paclo 2SC are an effective tool for controlling the growth of woody shrubs in landscape settings. The growth control is approximately 90% and lasts for 1-2 years, depending on the length of the growing season.

Applications are made with a diluted mixture composed of 1 part Tide Paclo 2SC to 11 parts water. Mix 10.7 fl oz (317 mL, 0.167 lbs. a.i.) of Tide Paclo 2SC with water to make one gallon of diluted mixture. Mix one gallon (2 lbs. a.i.) of Tide Paclo 2SC with water to make 12 gallons of diluted mixture.

1. Identify the shrub species and calculate the size of the "shrub crown".
  - a) The size (or volume) of the "shrub crown" is calculated by using the volume formula  $L \times W \times H$  (in feet).
  - b) When measuring crown volume, include only significant branches. Do not include stems or the trunk in the volume calculations if it is not within the crown itself.

$$\text{Volume of Shrub} = \text{Height} \times \text{Length} \times \text{Depth} \\ 3 \text{ ft} \times 4 \text{ ft} \times 2 \text{ ft} = 24 \text{ ft}^3$$



- c) Determine how much Tide Paclo 2SC diluted mixture should be used on the selected woody shrubs. Match the total shrub volume in cubic feet (ft<sup>3</sup>) with the volume amounts in the table "**Drench Rates for Individual Woody Shrubs Using Diluted Mixture of Tide Paclo 2SC Tree Growth Regulator Total Volume of Diluted Mixture In Milliliters & Ounces**" to determine the total volume. If the amount needed is found to be between two volumes, select either the average of the two volumes or the higher of the two volumes.

#### Basal Drench

Prior to application, dig a shallow furrow 2 – 6 inches deep around the base of the shrub near the point of contact between the soil and the shrub trunk (Figure 1). Carefully pour the diluted mixture of Tide Paclo 2SC evenly around the shrub into the furrow with a graduated container/jug or with a hand-held hose connected to a trunk-mounted tank/hydraulic sprayer. To avoid possible product runoff after applying, refill the furrow with untreated soil.

## Restriction

Do not apply product to soil when soil is already saturated. Heavy rainfall or irrigation in treated areas may cause active ingredient to move laterally on slopes and collect in low areas. These areas may undergo more severe growth control for a longer period of time.

## Soil Injection

Inject the Ready to Use solution approximately 2-6 inches deep at 50-200 psi using the volumes in the table "Drench Rates for Individual Woody Shrubs using Diluted Mixture of Tide Paclo 2SC Tree Growth Regulator Total Volume of Diluted Mixture In Milliliters & Ounces." Orient injection orifices to release the diluted product horizontally at the point of injection. Divide the required dose evenly among injection sites spaced as uniformly as possible around the base of the tree.

Position the injection sites to release the diluted Tide Paclo 2SC as close as possible to the point of contact between the soil and the plant beneath the soil so that the solution is readily absorbed by the roots (Figure 2). Use at least 4 injection sites evenly spaced around the plant.

## Notes for Woody Shrubs Soil Applications

- Newly planted or containerized shrubs need only 60% of the prescribed volume.
- For single stem applications, place product closely around the shrub base.
- For multiple stem applications, place product close around the base and within stem groupings.
- Intermittent trimming to keep tight formal appearance will not remove enough product to lose growth control in most shrubs.

| Drench Rates for Individual Woody Shrubs Using Diluted Mixture of Tide Paclo 2SC Tree Growth Regulator Total Volume of Diluted Mixture In Milliliters & Ounces |              |      |           |  |      |           |   |      |           |                              |      |           |
|--|--------------|------|-----------|--|------|-----------|---|------|-----------|------------------------------|------|-----------|
| Crown Volume (ft <sup>3</sup> )  | Spirea       |      |           | Boxwood, Privet, Ligustrum, Rhododendron |      |           | Abelia, Elaeagnus, Forsythia japonicum, Korean viburnum |      |           | Barberry, Camellia, Photinia |      |           |
|  | mLs          | oz   | lbs. a.i. | mLs                                      | oz   | lbs. a.i. | mLs   | oz   | lbs. a.i. | mLs                          | oz   | lbs. a.i. |
| 4  | 100          | 3.4  | 0.0044    | 155                                      | 5.2  | 0.0068    | 190   | 6.4  | 0.0084    | 200                          | 6.8  | 0.0089    |
| 8  | 115          | 3.9  | 0.0051    | 175                                      | 5.9  | 0.0077    | 200   | 6.8  | 0.0089    | 230                          | 7.8  | 0.0102    |
| 12   | 130          | 4.4  | 0.0057    | 200                                      | 6.8  | 0.0089    | 210   | 7.1  | 0.0093    | 270                          | 9.1  | 0.0119    |
| 18   | 150          | 5.1  | 0.0067    | 220                                      | 7.4  | 0.0097    | 220   | 7.4  | 0.0097    | 305                          | 10.3 | 0.0135    |
| 27   | 165          | 5.6  | 0.0073    | 240                                      | 8.1  | 0.0106    | 235   | 7.9  | 0.0103    | 340                          | 11.5 | 0.0150    |
| 36   | 180          | 6.1  | 0.0080    | 260                                      | 8.8  | 0.0115    | 250   | 8.4  | 0.0110    | 380                          | 12.8 | 0.0167    |
| 48   | 200          | 6.8  | 0.0089    | 275                                      | 9.3  | 0.0121    | 265   | 9    | 0.0118    | 405                          | 13.7 | 0.0179    |
| 64   | 215          | 7.3  | 0.0095    | 290                                      | 9.8  | 0.0128    | 300   | 10.1 | 0.0132    | 420                          | 14.2 | 0.0185    |
| 80   | 230          | 7.8  | 0.0102    | 305                                      | 10.3 | 0.0135    | 325   | 11   | 0.0144    | 435                          | 14.7 | 0.0192    |
| 100  | 250          | 8.4  | 0.0110    | 315                                      | 10.6 | 0.0138    | 340   | 11.5 | 0.0150    | 455                          | 15.4 | 0.0201    |
| 130  | 260          | 8.8  | 0.0115    | 345                                      | 11.7 | 0.0153    | 360   | 12.2 | 0.0159    | 500                          | 16.9 | 0.0221    |
| 150  | 270          | 9.1  | 0.0119    | 365                                      | 12.3 | 0.0161    | 370   | 12.5 | 0.0163    | 520                          | 17.6 | 0.0230    |
| 180  | 280          | 9.5  | 0.0124    | 390                                      | 13.2 | 0.0172    | 385   | 13   | 0.0170    | 545                          | 18.4 | 0.0240    |
| 220  | 290          | 9.8  | 0.0128    | 420                                      | 14.2 | 0.0185    | 405   | 13.7 | 0.0179    | 565                          | 19.1 | 0.0249    |
| 250  | 300          | 10.1 | 0.0132    | 435                                      | 14.7 | 0.0192    | 415   | 14   | 0.0183    | 575                          | 19.4 | 0.0253    |
| 300  | 310          | 10.5 | 0.0137    | 460                                      | 15.5 | 0.0202    | 430   | 14.5 | 0.0189    | 590                          | 19.9 | 0.0260    |
| Crown Volume (ft <sup>3</sup> )  | Burning Bush |      |           | Viburnum, Rose of Sharon, Holly          |      |           | Arborvitae, Cherry, Laurel, Juniper, Yew                |      |           |                              |      |           |
|  | mLs          | oz   | lbs. a.i. | mLs                                      | oz   | lbs. a.i. | mLs   | oz   | lbs. a.i. |                              |      |           |
| 4  | 250          | 8.4  | 0.0110    | 340                                      | 11.5 | 0.0150    | 400   | 13.5 | 0.0176    |                              |      |           |
| 8  | 270          | 9.1  | 0.0119    | 345                                      | 11.7 | 0.0153    | 450   | 15.2 | 0.0199    |                              |      |           |
| 12   | 300          | 10.1 | 0.0132    | 350                                      | 11.8 | 0.0154    | 510   | 17.2 | 0.0225    |                              |      |           |
| 18   | 350          | 11.8 | 0.0154    | 370                                      | 12.5 | 0.0163    | 600   | 20.3 | 0.0265    |                              |      |           |

|     |     |      |        |     |      |        |      |      |        |
|-----|-----|------|--------|-----|------|--------|------|------|--------|
| 27  | 370 | 12.5 | 0.0163 | 380 | 12.8 | 0.0167 | 650  | 22   | 0.0287 |
| 36  | 410 | 13.9 | 0.0182 | 400 | 13.5 | 0.0176 | 700  | 23.6 | 0.0308 |
| 48  | 450 | 15.2 | 0.0199 | 430 | 14.5 | 0.0189 | 800  | 27   | 0.0353 |
| 64  | 480 | 16.2 | 0.0212 | 450 | 15.2 | 0.0199 | 900  | 30.4 | 0.0397 |
| 80  | 510 | 17.2 | 0.0225 | 480 | 16.2 | 0.0212 | 1000 | 33.8 | 0.0441 |
| 100 | 550 | 18.6 | 0.0243 | 510 | 17.2 | 0.0225 | 1200 | 40.5 | 0.0529 |
| 130 | 580 | 19.6 | 0.0256 | 550 | 18.6 | 0.0243 | 1300 | 43.9 | 0.0573 |
| 150 | 610 | 20.6 | 0.0269 | 580 | 19.6 | 0.0256 | 1400 | 47.3 | 0.0618 |
| 180 | 660 | 22.3 | 0.0291 | 650 | 22   | 0.0287 | 1550 | 52.4 | 0.0684 |
| 220 | 680 | 23   | 0.0300 | 710 | 24   | 0.0313 | 1650 | 55.7 | 0.0728 |
| 250 | 710 | 24   | 0.0313 | 750 | 25.3 | 0.0330 | 1850 | 62.5 | 0.0816 |
| 300 | 750 | 25.3 | 0.0330 | 770 | 26   | 0.0340 | 2000 | 67.6 | 0.0883 |

### Soil Treatment for Vines

Tide Paclo 2SC can also be used as a basal drench or soil injection to reduce the growth of vines. Refer to sections above for application instructions. Mix 1 part Tide Paclo 2SC to 11 parts water. Mix 10.7 fl oz (317 mL, 0.167 lbs. a.i.) of Tide Paclo 2SC with water to make one gallon of diluted mixture. Mix one gallon (2.0 lbs. a.i.) of Tide Paclo 2SC with water to make 12 gallons of diluted mixture. The amount diluted mixture is based upon the accumulated surface area of the vines to be regulated.

### Soil Treatment for Vine Species and Rates for Vegetative Growth Control

| Plant Name           | Scientific Name                           | Rate   |
|----------------------|---|--|
| Boston Ivy           | <i>Parthenocissus tricuspidata</i>        | 0.33 gallons of diluted mixture<br>(0.055 lbs. a.i.) per 100 ft <sup>2</sup> of vine |
| Creeping Fig         | <i>Ficus pumila</i> , <i>Ficus repens</i> |  |
| English Ivy          | <i>Hedera</i> spp.                        |  |
| Japanese Honeysuckle | <i>Lonicera japonica</i>                  |  |
| Trumpet Creeper      | <i>Campsis radicans</i>                   |  |
| Wisteria             | <i>Wisteria</i> spp.                      |  |

### Foliar Sprays

Tide Paclo 2SC as a foliar spray can provide growth suppression of a wide range of shrubs and vines growing in outdoor, non-crop areas, nurseries, parks, commercial buildings, right-of-way areas, street medium, roadsides, fence rows and non-irrigation ditch banks. Tide Paclo 2SC is absorbed through leaves, buds, new shoots and roots and applications can be made throughout the growing season. See the table "Foliar Spray Application Rate Ranges for Diluted Mixture of Tide Paclo 2SC for Treatment of Various Shrub and Vine Species" for foliar spray rates and associated application notes. To limit unwanted surface runoff in outdoor ornamental uses, do not apply when growth media is saturated.

### Foliar Spray Application Rate Ranges for Diluted Mixture of Tide Paclo 2SC for Treatment of Various Shrub and Vine Species

| Plant Name     | Scientific Name             | Rate per Gallon of Spray Solution |          |                 |
|----------------|-----------------------------|-----------------------------------|----------|-----------------|
|                |                             | fl. oz.                           | mL       | lbs. a.i.       |
| Abelia         | <i>Abella x grandiflora</i> | 1.0 - 3.5                         | 30 - 105 | 0.0156 - 0.0547 |
| Alpine Current | <i>Ribes</i> spp.           | 1.0 - 2.5                         | 30 - 75  | 0.0156 - 0.0391 |
| Arboricola     | <i>Shefflera arboricola</i> | 2.5 - 5.0                         | 75 - 150 | 0.0391 - 0.0781 |
| Arborvitae     | <i>Thuja</i> spp.           | 2.5 - 5.0                         | 75 - 150 | 0.0391 - 0.0781 |
| Azalea         | <i>Rhododendron</i> spp.    | 0.5 - 2.5                         | 15 - 75  | 0.0078 - 0.0391 |
| Barberry       | <i>Berberis</i> spp.        | 1.0 - 2.5                         | 30 - 75  | 0.0156 - 0.0391 |
| Bottlebrush    | <i>Callistemon</i> spp.     | 1.0 - 2.5                         | 30 - 75  | 0.0156 - 0.0391 |

**Foliar Spray Application Rate Ranges for Diluted Mixture of Tide Paclo 2SC for  
Treatment of Various Shrub and Vine Species**

| Plant Name                     | Scientific Name                               | Rate per Gallon of Spray Solution |          |                 |
|--------------------------------|---|-----------------------------------|----------|-----------------|
|                                |   | fl. oz.                           | mL       | lbs. a.i.       |
| Boston Ivy                     | <i>Parthenocissus tricuspidata</i>            | 0.5 - 2.5                         | 15 - 75  | 0.0078 - 0.0391 |
| Bougainvillea                  | <i>Bougainvillea</i> spp.                     | 2.5 - 5.0                         | 75 - 150 | 0.0391 - 0.0781 |
| Boxwood                        | <i>Buxus</i> spp.                             | 1.0 - 3.5                         | 30 - 105 | 0.0156 - 0.0547 |
| Butterfly Bush                 | <i>Buddleia</i> spp.                          | 0.5 - 1.0                         | 15 - 30  | 0.0078 - 0.0156 |
| Cherry Laurel & English Laurel | <i>Prunus</i> spp.                            | 0.5 - 1.0                         | 15 - 30  | 0.0078 - 0.0156 |
| Cocoplum                       | <i>Chrysobalanus icaco</i>                    | 1.0 - 5.0                         | 30 - 150 | 0.0156 - 0.0781 |
| Copperleaf                     | <i>Acalypha wilkesiana</i>                    | 1.0 - 5.0                         | 30 - 150 | 0.0156 - 0.0781 |
| Cotoneaster                    | <i>Cotoneaster</i> spp.                       | 1.0 - 2.5                         | 30 - 75  | 0.0156 - 0.0391 |
| Creeping Fig                   | <i>Ficus pumila, Ficus repens</i>             | 2.5 - 5.0                         | 75 - 150 | 0.0391 - 0.0781 |
| Elaeagnus                      | <i>Elaeagnus pungens</i>                      | 2.5 - 5.0                         | 75 - 150 | 0.0391 - 0.0781 |
| English Ivy                    | <i>Hedera</i> spp.                            | 0.5 - 2.5                         | 15 - 75  | 0.0078 - 0.0391 |
| Escallonia                     | <i>Escallonia</i> spp.                        | 2.5 - 5.0                         | 75 - 150 | 0.0391 - 0.0781 |
| Eugenia                        | <i>Eugenia myrtifolia</i>                     | 1.0 - 2.5                         | 30 - 75  | 0.0156 - 0.0391 |
| Euonymus                       | <i>Euonymus</i> spp.                          | 2.5 - 3.5                         | 75 - 105 | 0.0391 - 0.0547 |
| Ficus                          | <i>Fiscus</i> spp.                            | 2.5 - 5.0                         | 75 - 150 | 0.0391 - 0.0781 |
| Firebush                       | <i>Hamelia patens</i>                         | 2.5 - 5.0                         | 75 - 150 | 0.0391 - 0.0781 |
| Firecracker Plant              | <i>Russelia equisetiformis</i>                | 2.5 - 5.0                         | 75 - 150 | 0.0391 - 0.0781 |
| Forsythia                      | <i>Forsythia</i> spp.                         | 1.0 - 1.7                         | 30 - 50  | 0.0156 - 0.0266 |
| Hibiscus                       | <i>Hibiscus</i> spp.                          | 0.5 - 1.0                         | 15 - 30  | 0.0078 - 0.0156 |
| Holly                          | <i>Ilex</i> spp.                              | 2.5 - 5.0                         | 75 - 150 | 0.0391 - 0.0781 |
| Honeylocust                    | <i>Gleditsia</i>                              | 1.0 - 2.5                         | 30 - 75  | 0.0156 - 0.0391 |
| Honeysuckle                    | <i>Lonicera</i> spp.                          | 0.5 - 1.0                         | 15 - 30  | 0.0078 - 0.0156 |
| Ice Plant                      | <i>Delosperma</i> spp.                        | 0.5 - 1.0                         | 15 - 30  | 0.0078 - 0.0156 |
| Indian Hawthorne               | <i>Raphiolepis indica</i>                     | 2.5 - 5.0                         | 75 - 150 | 0.0391 - 0.0781 |
| Itea                           | <i>Itea virginiana</i>                        | 0.5-0.5                           | 15 - 15  | 0.0078 - 0.0078 |
| Ixora                          | <i>Ixora coccinea</i>                         | 2.5 - 5.0                         | 75 - 150 | 0.0391 - 0.0781 |
| Jasmine (Asiatic)              | <i>Trachelospermum asiaticum</i>              | 1.0 - 3.5                         | 30 - 105 | 0.0156 - 0.0547 |
| Jasmine (Confederate/Star)     | <i>Trachelospermum jasminoides</i>            | 1.0 - 2.5                         | 30 - 75  | 0.0156 - 0.0391 |
| Juniper                        | <i>Juniperus</i> spp.                         | 2.5 - 5.0                         | 75 - 150 | 0.0391 - 0.0781 |
| Lantana                        | <i>Lantana camara</i>                         | 1.0 - 2.5                         | 30 - 75  | 0.0156 - 0.0391 |
| Lilac                          | <i>Syringa</i> spp.                           | 0.5 - 2.5                         | 15 - 75  | 0.0078 - 0.0391 |
| Loropetalum                    | <i>Loropetalum chinensis</i>                  | 0.5 - 2.5                         | 15 - 75  | 0.0078 - 0.0391 |
| Manhattan Euonymus             | <i>Euonymus kiautschovicus</i><br>"Manhattan" | 2.5 - 5.0                         | 75 - 150 | 0.0391 - 0.0781 |
| Ninebark                       | <i>Physocarpus</i> spp.                       | 0.5 - 1.0                         | 15 - 30  | 0.0078 - 0.0156 |
| Oleander                       | <i>Nerium</i> spp.                            | 1.0 - 2.5                         | 30 - 75  | 0.0156 - 0.0391 |
| Orange Jasmine                 | <i>Murraya paniculata</i>                     | 0.5 - 3.5                         | 15 - 105 | 0.0078 - 0.0547 |

**Foliar Spray Application Rate Ranges for Diluted Mixture of Tide Paclo 2SC for  
Treatment of Various Shrub and Vine Species**

| Plant Name          | Scientific Name              | Rate per Gallon of Spray Solution |          |                 |
|---------------------|------------------------------|-----------------------------------|----------|-----------------|
|                     |                              | fl. oz.                           | mL       | lbs. a.i.       |
| Photinia            | <i>Photinia fraseri</i>      | 2.5 - 5.0                         | 75 - 150 | 0.0391 - 0.0781 |
| Pittosporum         | <i>Pittosporum</i> spp.      | 2.5 - 5.0                         | 75 - 150 | 0.0391 - 0.0781 |
| Plumbago            | <i>Plumbago auriculata</i>   | 1.0 - 5.0                         | 30 - 150 | 0.0156 - 0.0781 |
| Podocarpus          | <i>Podocarpus</i> spp.       | 1.0 - 2.5                         | 30 - 75  | 0.0156 - 0.0391 |
| Privet              | <i>Ligustrum</i> spp.        | 2.5 - 5.0                         | 75 - 150 | 0.0391 - 0.0781 |
| Pyracantha          | <i>Pyracantha</i> spp.       | 2.5 - 3.5                         | 75 - 105 | 0.0391 - 0.0547 |
| Rhaphiolepis        | <i>Rhaphiolepis indica</i>   | 1.0 - 5.0                         | 30 - 150 | 0.0156 - 0.0781 |
| Rhododendron        | <i>Rhododendron</i> spp.     | 0.5 - 2.5                         | 15 - 75  | 0.0078 - 0.0391 |
| Rose                | <i>Rosa</i> spp.             | 1.0 - 3.5                         | 30 - 105 | 0.0156 - 0.0547 |
| Rose of Sharon      | <i>Hibiscus syriacus</i>     | 0.5 - 1.0                         | 15 - 30  | 0.0078 - 0.0156 |
| Schefflera          | <i>Schefflera arboricola</i> | 2.5 - 5.0                         | 75 - 150 | 0.0391 - 0.0781 |
| Serviceberry        | <i>Amelanchier</i> spp.      | 0.5 - 1.0                         | 15 - 30  | 0.0078 - 0.0156 |
| Spirea              | <i>Spirea</i> spp.           | 0.5 - 1.0                         | 15 - 30  | 0.0078 - 0.0156 |
| Surinam Cherry      | <i>Eugenia uniflora</i>      | 1.0 - 2.5                         | 30 - 75  | 0.0156 - 0.0391 |
| Trifoliolate Orange | <i>Poncirus trifoliata</i>   | 2.5 - 5.0                         | 75 - 150 | 0.0391 - 0.0781 |
| Viburnum            | <i>Viburnum</i> spp.         | 1.0 - 5.0                         | 30 - 150 | 0.0156 - 0.0781 |
| Weigela             | <i>Weigela florida</i>       | 0.5 - 1.0                         | 15 - 30  | 0.0078 - 0.0156 |
| Winter Jasmine      | <i>Jasminum nudiflorum</i>   | 2.5 - 3.5                         | 75 - 105 | 0.0391 - 0.0547 |
| Yew                 | <i>Taxus</i> spp.            | 2.5 - 5.0                         | 75 - 150 | 0.0391 - 0.0781 |

**Plant Foliar Sprays**

- To minimize regrowth after pruning make applications within two weeks following pruning.
- Pruning after applications have been made can remove the product from the plant and decrease the amount of growth reduction.
- Use a non-ionic surfactant for best results. Always read and follow all cautionary statements and other information appearing on the surfactant label.
- Maintain agitation in the spray tank to ensure uniform distribution within the spray solution.
- Apply the foliar spray using a medium-coarse spray until spray solution begins to drip from all plant surfaces. Ensure that all inner stems are thoroughly covered.
- Wait at least 8 weeks between treatments.
- In outdoor commercial ornamental and nursery uses, follow foliar applications of Tide Paclo 2SC with irrigation within 24 hours to remove product from foliage and limit surface movement. If overhead irrigation is not available, time treatment applications to allow Tide Paclo 2SC to dry on the treated surface prior to rainfall.
- To limit unwanted surface runoff in outdoor ornamental uses, do not apply when growth media is saturated.
- The rates in the tables "**Drench Rates for Individual Woody Shrubs Using Diluted Mixture of Tide Paclo 2SC Tree Growth Regulator Total Volume of Diluted Mixture in Milliliters & Ounces,**" "**Soil Treatment for Vine Species and Rates for Vegetative Growth Control,**" and "**Foliar Spray Application Rate Ranges for Diluted Mixture of Tide Paclo 2SC for Treatment of Various Shrub and Vine Species**" are guidelines as some species may respond more or less to Tide Paclo 2SC. Efficacy may also vary depending on weather conditions, geographic conditions and other biological factors. Treat a small number of plants prior to determining specific application rates for different species under actual use conditions. Use the higher rates when treating subtropical plants growing in

locations with longer growing seasons.

- Foliar sprays at higher dosage rates may leave a white residue on the plant foliage. Take precautions when treating around sidewalks, driveways, buildings, decks, fences, vehicles or other structural surfaces as staining may occur. Wash immediately with water if product comes into contact with these surfaces.
- Take precaution to minimize application to non-target plants that come in contact with Tide Paclo 2SC.
- Avoid pruning following application in order to not remove the terminal shoots of the treated plant. However, certain species may require light pruning to remove unregulated shoots and maintain shape and form.

---

## USE AND RESTRICTIONS

---

- Apply at labeled rates and always follow safety procedures.
- For shrub and vine use, **do not** use on areas to be cultivated for food or food crops within two years of treatment.
- For shrub and vine use, **do not** apply more than 1 gallon per acre per application (2 lbs. ai/A).
- For shrub and vine use, **do not** apply more than 1 gallon per acre per year (2 lbs. ai/A).
- For shrub and vine use, **do not** treat plants more than 4 times per year.
- Basal drench and soil injection application of Tide Paclo 2SC may result in localized, temporary discoloration of turfgrass immediately adjacent to the treatment site.
- Ensure that basal drenches on slopes or other areas subject to erosion by rainfall or irrigation are deep enough and adequately covered with fresh soil. The use of one or more soil dams in the temporary berm can increase application uniformity.

**Chemigation:** Do not apply this product through any type of irrigation system.

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.

**PESTICIDE STORAGE:** Keep container closed when not in use. Do not store near food or feed. Protect from freezing. In case of spill or leak on floor or paved surfaces, soak up with sand, earth or synthetic absorbent. Remove to chemical waste area.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the hazardous waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

## LIMITED WARRANTY AND DISCLAIMER

Zhejiang Tide CropScience Co., Ltd. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below.

Our recommendations for use of this product are based upon tests believed to be reliable. The use of this product being beyond control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such use or the results to be obtained if not used in accordance with printed directions and established safe practice. To the fullest extent



permitted by law, buyer's exclusive remedy and manufacturer's or seller's exclusive liability for any and all claims, losses, damages or injuries resulting from the use or handling of this product, whether or not based in contract, negligence, strict liability in tort or otherwise shall be limited, at the manufacturer's option to replacement of, or the repayment of the purchase price for, the quantity of product with respect to which damages are claimed.

#### **Inherent Risks of Use**

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Zhejiang Tide CropScience Co., Ltd. or the seller. To the fullest extent permitted by law, all such risks shall be assumed by buyer.

#### **Terms and Conditions of Use**

If terms of the Limited Warranty & Disclaimer and Inherent Risks of Use are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, to the fullest extent permitted by law, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer and Inherent Risks of Use.

Label Code No.: ZTC-PGR806974

EPA 20221215

Created Date: 01/09/2023