



0 - 0 - 1

GUARANTEED ANALYSIS

Soluble Potash (K_2O) 1%

Derived from seaweed extract (*Ascophyllum nodosum*) and potassium hydroxide

F000148

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.html>

For your convenience, **TOGGLE**® product information is available on Agrian database (<http://agrian.com/home/>) as well as in Crop Data Management Systems database (<http://www.cdms.net/>)

STORAGE AND HANDLING: This product should be stored away from intense sunlight and heat.

STIR BEFORE USE

CAUTION

Keep out of reach of children.

WARRANTY STATEMENT:

The manufacturer warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions under normal conditions of use. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials or the manner of use or application, all of which are beyond the control of the manufacturer. In no case shall the manufacturer be liable for consequential, special or indirect damages resulting from the use or handling of this product. The manufacturer makes no warranties of merchantability or fitness for a particular purpose nor any other express or implied warranty except as stated above.

Manufactured and guaranteed by:

Acadian Seaplants Limited
30 Brown Avenue, Dartmouth, Nova Scotia
Canada, B3B 1X8
Tel: +1 902 468 2840
Fax: +1 902 468 3474

Acadian Plant Health™ is a division of
Acadian Seaplants Limited



NET CONTENT:
2.5 U.S. Gal.
9.4 Liters

NET WEIGHT:
21.53 lb
9.77 kg

USA_TG1427a-2_0623



ACADIAN™
PLANT HEALTH

GENERAL INFORMATION: Consistent use of **TOGGLE®** supplements a well-balanced crop nutrition program. Use **TOGGLE®** to:

- Maximize crop potential during periods of stress
- Improve drought tolerance and resistance to extreme temperatures (cold and heat)
- Increase and maximize yield
- Optimize nutrient uptake
- Improve root growth and development
- Improve photosynthesis

To achieve the desired results, the levels of major and minor nutrients must be adequate to support the increase in production. The following rates and timings of application are recommended for optimum efficacy.

COMPATIBILITY: **TOGGLE®** is compatible with most insecticides, fungicides and fertilizers. When mixing with calcium products, thoroughly mix **TOGGLE®** with the water in the tank prior to adding the calcium product. If interaction of chemicals is unknown, a “jar” compatibility test is suggested.

DIRECTIONS FOR USE

TOGGLE® fully dissolves in water and is suitable for use in liquid foliar, soil applied, and irrigation water applications. With the versatility to apply in-furrow or by foliar application, **TOGGLE®** works synergistically with a grower's fertilizer program to improve the crop's use of nutrients and make the best use of production inputs. Regular applications are important for maximizing crop potential during unexpected stress.

TOGGLE® can be tank mixed with most commonly used insecticides, fungicides and herbicides.

The product may not be applied near water, storm drains or drainage ditches or to any impervious surface.

The product may not be applied if heavy rain is expected.

The product may only be applied to the intended application site.

| Crop | Application | Rate per Acre | Recommendation and Timings |
|---|-------------|-----------------|--|
| Corn (Field corn, silage, sweet corn, popcorn, seed corn) | 1st | 8 ounces | In-furrow or banded at planting, or with row starter. † |
| | 2nd | 32 to 48 ounces | Foliar broadcast spray at V4 - VT. Note: Always follow fungicide labels regarding surfactant use at/near tassel stages. |
| Soybean | 1st | 8 ounces | In-furrow or banded at planting, or with row starter. † |
| | 2nd | 32 to 48 ounces | Foliar broadcast from V4 - R3. |
| Wheat | 1st | 8 ounces | In-furrow or banded at planting, or with row starter. † |
| | 2nd | 32 to 48 ounces | At tiller initiation. |
| | 3rd | 32 to 48 ounces | At early boot to early panicle emergence. |
| Cotton | 1st | 8 ounces | In-furrow or banded at planting, or with row starter. † |
| | 2nd | 32 to 48 ounces | Foliar broadcast at pinhead square. † |
| | 3rd | 32 to 48 ounces | Foliar broadcast two weeks after pinhead square. |
| Rice | 1st | 8 ounces | In-furrow or banded at planting, or with row starter. † |
| | 2nd | 32 to 48 ounces | Foliar broadcast at tiller initiation. † |
| | 3rd | 32 to 48 ounces | Foliar broadcast at early panicle emergence. † |
| Alfalfa, Mixed Hay | 1st | 8 ounces | In-furrow or banded at planting, or with row starter. † |
| | 2nd | 32 to 48 ounces | Soon after each cutting. † |
| | 3rd | 32 to 48 ounces | Two weeks after cutting. † |
| Potatoes | 1st | 8 ounces | In-furrow or banded at planting, or with row starter. † |
| | 2nd | 32 to 48 ounces | At tuber initiation. † |
| | 3rd | 32 to 48 ounces | Through late tuber set (if more potatoes are desired). † |
| Peanuts | 1st | 8 ounces | In-furrow or banded at planting, or with row starter. † |
| | 2nd | 32 to 48 ounces | After plants are established through to harvest. † |
| All other row crops | 1st | 8 ounces | In-furrow or banded at planting, or with row starter. † |
| | 2nd | 32 to 48 ounces | After planting through to harvest. † |

Additional applications are recommended prior to major stress events or after severe hail or insect damage or herbicide damage.

† Key timings for best results. For more information see your **TOGGLE®** authorized distributor or **TOGGLE®** representative.



0 - 0 - 1

GUARANTEED ANALYSIS

Soluble Potash (K_2O)

1%

Derived from seaweed extract (*Ascophyllum nodosum*) and potassium hydroxide

Density at 68°F (20°C) 8.67 lb / U.S. Gal (1.04 kg/L)

F000148

STORAGE AND HANDLING: This product should be stored away from intense sunlight and heat.

STIR BEFORE USE

CAUTION

Keep out of reach of children.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.html>

For your convenience, **TOGGLE**® product information is available on Agrian database (<http://agrian.com/home/>) as well as in Crop Data Management Systems database (<http://www.cdms.net/>)

WARRANTY STATEMENT:

The manufacturer warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions under normal conditions of use. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials or the manner of use or application, all of which are beyond the control of the manufacturer. In no case shall the manufacturer be liable for consequential, special or indirect damages resulting from the use or handling of this product. The manufacturer makes no warranties of merchantability or fitness for a particular purpose nor any other express or implied warranty except as stated above.

Manufactured and guaranteed by:

Acadian Seaplants Limited
30 Brown Avenue, Dartmouth, Nova Scotia
Canada, B3B 1X8
Tel: +1 902 468 2840
Fax: +1 902 468 3474

Acadian Plant Health™ is a division of
Acadian Seaplants Limited

PRODUCT OF CANADA



NET CONTENT:
2 x 2.5 U.S. Gal.
2 x 9.4 Liters

NET WEIGHT:
43.06 lb
19.54 kg

GENERAL INFORMATION: Consistent use of **TOGGLE**® supplements a well-balanced crop nutrition program. Use **TOGGLE**® to:

- Maximize crop potential during periods of stress
- Improve drought tolerance and resistance to extreme temperatures (cold and heat)
- Increase and maximize yield
- Optimize nutrient uptake
- Improve root growth and development
- Improve photosynthesis

To achieve the desired results, the levels of major and minor nutrients must be adequate to support the increase in production. The following rates and timings of application are recommended for optimum efficacy.

COMPATIBILITY: **TOGGLE**® is compatible with most insecticides, fungicides and fertilizers. When mixing with calcium products, thoroughly mix **TOGGLE**® with the water in the tank prior to adding the calcium product. If interaction of chemicals is unknown, a "jar" compatibility test is suggested.

DIRECTIONS FOR USE

TOGGLE® fully dissolves in water and is suitable for use in liquid foliar, soil applied, and irrigation water applications. With the versatility to apply in-furrow or by foliar application, **TOGGLE**® works synergistically with a grower's fertilizer program to improve the crop's use of nutrients and make the best use of production inputs. Regular applications are important for maximizing crop potential during unexpected stress. **TOGGLE**® can be tank mixed with most commonly used insecticides, fungicides and herbicides.

The product may not be applied near water, storm drains or drainage ditches or to any impervious surface.

The product may not be applied if heavy rain is expected.

The product may only be applied to the intended application site.

| Crop | Application | Rate per Acre | Recommendation and Timings |
|---|-------------|-----------------|--|
| Corn (Field corn, silage, sweet corn, popcorn, seed corn) | 1st | 8 ounces | In-furrow or banded at planting, or with row starter. † |
| | 2nd | 32 to 48 ounces | Foliar broadcast spray at V4 - VT. Note: Always follow fungicide labels regarding surfactant use at/near tassel stages. |
| Soybean | 1st | 8 ounces | In-furrow or banded at planting, or with row starter. † |
| | 2nd | 32 to 48 ounces | Foliar broadcast from V4 - R3. |
| Wheat | 1st | 8 ounces | In-furrow or banded at planting, or with row starter. † |
| | 2nd | 32 to 48 ounces | At tiller initiation. |
| | 3rd | 32 to 48 ounces | At early boot to early panicle emergence. |
| Cotton | 1st | 8 ounces | In-furrow or banded at planting, or with row starter. † |
| | 2nd | 32 to 48 ounces | Foliar broadcast at pinhead square. † |
| | 3rd | 32 to 48 ounces | Foliar broadcast two weeks after pinhead square. |
| Rice | 1st | 8 ounces | In-furrow or banded at planting, or with row starter. † |
| | 2nd | 32 to 48 ounces | Foliar broadcast at tiller initiation. † |
| | 3rd | 32 to 48 ounces | Foliar broadcast at early panicle emergence. † |
| Alfalfa, Mixed Hay | 1st | 8 ounces | In-furrow or banded at planting, or with row starter. † |
| | 2nd | 32 to 48 ounces | Soon after each cutting. † |
| | 3rd | 32 to 48 ounces | Two weeks after cutting. † |
| Potatoes | 1st | 8 ounces | In-furrow or banded at planting, or with row starter. † |
| | 2nd | 32 to 48 ounces | At tuber initiation. † |
| | 3rd | 32 to 48 ounces | Through late tuber set (if more potatoes are desired). † |
| Peanuts | 1st | 8 ounces | In-furrow or banded at planting, or with row starter. † |
| | 2nd | 32 to 48 ounces | After plants are established through to harvest. † |
| All other row crops | 1st | 8 ounces | In-furrow or banded at planting, or with row starter. † |
| | 2nd | 32 to 48 ounces | After planting through to harvest. † |

Additional applications are recommended prior to major stress events or after severe hail or insect damage or herbicide damage.

† Key timings for best results. For more information see your **TOGGLE**® authorized distributor or **TOGGLE**® representative.