



Tourney® EZ



F U N G I C I D E



**TURFGRASS FUNGICIDE
A SYSTEMIC FUNGICIDE
FOR CONTROL OF THE
LISTED TURFGRASS DISEASES**

Active Ingredient	By Wt
Metconazole**	42.5%
Other Ingredients	57.5%
Total	100.0%

** (1*RS*,5*RS*;1*RS*,5*SR*)-5-(4-chlorobenzyl)-
2,2-dimethyl-1-(1*H*-1,2,4-triazol-1-
ylmethyl)cyclopentanol

Tourney® EZ Fungicide is a suspension concentrate containing 4 lb active ingredient per gallon.

EPA Reg. No. 59639-257

EPA Est. 11773-IA-1® 228-IL-1® 39578-TX-1®

5481-ID-1® 62171-MS-1® 70815-GA-2®

Superscript is first letter of lot number.



**ORNAMENTAL FUNGICIDE*
A SYSTEMIC FUNGICIDE FOR
CONTROL OF THE LISTED
ORNAMENTAL DISEASES**

*Not for use in California

**KEEP OUT OF
REACH OF CHILDREN
CAUTION**

**SEE NEXT PAGE FOR ADDITIONAL
PRECAUTIONARY STATEMENTS
AND FIRST AID.**

NET CONTENTS 1 GALLON

FIRST AID

If swallowed: Call a poison control center or doctor immediately for treatment advice.

Have person sip a glass of water if able to swallow.

DO NOT induce vomiting unless told to do so by the poison control center or doctor.

DO NOT give anything by mouth to an unconscious person.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center, doctor, or going for treatment. You may also contact **800-892-0099** for emergency medical treatment information.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS CAUTION

Harmful if swallowed. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Applicators and other handlers must wear: long-sleeved shirt and long pants, chemical-resistant gloves made of any waterproof material, socks and shoes.

USER SAFETY REQUIREMENTS

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to birds, mammals, fish, and aquatic invertebrates. **DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high-water mark. **DO NOT** contaminate water when disposing of equipment washwater or rinseate.

Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

Groundwater Advisory: This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Surface Water Advisory: This product may contaminate water through drift of spray in wind. Metconazole may impact surface water quality through runoff of rain water. This product has a high potential for runoff for several months or more after application. Poorly-draining soils and soils with shallow water tables are more prone to produce runoff that contains this product. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features including ponds, streams, and springs will reduce the potential for contamination of water from rainfall-runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL, USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, including plants, soil or water, is coveralls, chemical resistant gloves made of any waterproof material, socks and shoes.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to the uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, greenhouses or sodfarms.

Keep all unprotected persons out of operating areas or vicinity where there may be drift.

DO NOT enter or allow others to enter treated areas until sprays have dried.

RISKS OF USING THIS PRODUCT, LIMITED WARRANTY AND DISCLAIMER, AND LIMITATION OF LIABILITY

IMPORTANT: Read the entire Label including this section titled Risks of Using this Product, Limited Warranty and Disclaimer, and Limitation of Liability before using this product. If the terms are not acceptable **THEN DO NOT USE THE PRODUCT**; rather, return the unopened product within 15 days of purchase for a refund of the purchase price.

RISKS OF USING THIS PRODUCT

The buyer and user (referred to collectively herein as "Buyer") of this product should be aware that there are inherent unintended risks associated with the use of this product which are impossible to eliminate. Such risks of crop injury, non-performance, resistance or other unintended consequences are unavoidable and may result because of such factors as weather, soil conditions, disease, moisture conditions, irrigation practices, condition of the crop at the time of application, presence of other materials either applied in the tank mix with this product or prior to application of this product, cultural practices or the manner of use or application, (or a combination of such factors) all of which are factors beyond the control of Valent. If the Buyer chooses not to accept these risks, **THEN THIS PRODUCT SHOULD NOT BE APPLIED.** By applying this product Buyer acknowledges and accepts these inherent unintended risks **AND TO THE FULLEST EXTENT ALLOWED BY LAW, AGREES THAT ALL SUCH RISKS ASSOCIATED WITH THE APPLICATION AND USE ARE ASSUMED BY THE BUYER.**

The Directions for Use of this product must be followed carefully. Valent shall not be responsible for losses or damages (including, but not limited to, loss of yield, increased expenses of farming the crop or such incidental, consequential or special damages that may be claimed) resulting from use of this product in any manner not set forth on the label. To the extent consistent with applicable law, Buyer assumes all risks associated with the use of this product in any manner or under conditions not specifically directed or approved on the label.

(continued)

(continued)

LIMITED WARRANTY AND DISCLAIMER

Valent warrants only that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the label, under average use conditions, when used strictly in accordance with the label and **subject to the Risks of Using This Product as described above. To the extent consistent with applicable law, VALENT MAKES NO OTHER WARRANTIES, EITHER EXPRESSED OR IMPLIED.** No agent or representative of Valent or Seller is authorized to make or create any other express or implied warranty.

LIMITATION OF LIABILITY

To the fullest extent allowed by law, Valent or Seller is not liable for any incidental, consequential, indirect or special damages resulting from the use or handling of this product. TO THE FULLEST EXTENT ALLOWED BY LAW, THE EXCLUSIVE REMEDY OF THE BUYER, AND THE EXCLUSIVE MAXIMUM LIABILITY OF VALENT OR SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT SHALL BE THE RETURN OF THE PURCHASE PRICE OF THIS PRODUCT OR, AT THE ELECTION OF VALENT OR SELLER, THE REPLACEMENT OF THE PRODUCT.

PROMPT NOTICE OF CLAIM

To the extent consistent with applicable law allowing such requirements, Valent must be provided notice as soon as Buyer has reason to believe it may have a claim, but in no event later than twenty-one days from date of planting, or twenty-one days from the date of application, whichever is later, so that an immediate inspection of the affected property and growing crops can be made. To the extent consistent with applicable law, if Buyer does not notify Valent of any claims, in such period, it shall be barred from obtaining any remedy.

NO AMENDMENTS

Valent and Seller offer this product, and Buyer accepts it, subject to the foregoing **Risks of Using This Product, Limited Warranty and Disclaimer, and Limitation of Liability**, which may not be modified by any oral or written agreement.

PRODUCT INFORMATION

Tourney EZ Fungicide is formulated as a suspension concentrate. The active ingredient in *Tourney EZ* Fungicide is metconazole, a broad-spectrum triazole fungicide that works by inhibiting demethylation and other processes in sterol biosynthesis. *Tourney EZ* Fungicide is systemic and is quickly absorbed into plant tissue and can move up, but not down in the plant. Metconazole has no effect on fungal spore germination but interferes with other early developmental processes in the life cycle of certain fungi. Although *Tourney EZ* Fungicide cannot prevent spore germination, it prevents spore formation and inhibits mycelial growth.

Tourney EZ Fungicide may be used for disease control in turfgrass on golf courses, lawns and landscape areas around residential, institutional, public, commercial and industrial buildings, parks, recreational areas, athletic fields, and sod farms.

Tourney EZ Fungicide may be used for disease control on ornamental plants in commercial indoor and outdoor nurseries, greenhouses and on ornamental plants found in landscaped areas around residential, institutional, public, commercial and industrial buildings, parks, recreational areas, and athletic fields.

Tourney EZ Fungicide can be applied pre- or post-infection but is most effective when applied prior to infection. Optimal disease control is achieved when *Tourney EZ* Fungicide is applied in a regularly scheduled spray program used in combination and/or in rotation with other effective fungicides that have different modes of action (i.e., non-Group 3 fungicides).

MODE OF ACTION

The active ingredient in *Tourney EZ* Fungicide, metconazole, belongs to the sterol biosynthesis inhibitor group of fungicides as classified by the U.S. EPA and Canada PMRA as a target site of action Group 3 fungicide.

Resistance Management

For resistance management, *TourneyEZ* Fungicide contains a Group 3 fungicide. Any fungal population may contain individuals naturally resistant to *TourneyEZ* Fungicide and other Group 3 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Appropriate resistance-management strategies must be followed.

To delay fungicide resistance, take one or more of the following steps:

- Avoid application of more than maximum number of applications per year as listed in the Crop Specific Directions and maximum number of consecutive sprays of *TourneyEZ* Fungicide or other fungicides in the same group in a season.
- Use tank mixtures with fungicides from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host-plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological, and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.
- Contact your local extension specialist or certified crop advisor for additional pesticide resistance-management and/or IPM guidelines for specific crops and pathogens.
- For further information or to report suspected resistance, contact Valent U.S.A. LLC at 800-682-5368. You can also contact your pesticide distributor or university extension specialist to report resistance.

JAR TEST TO DETERMINE COMPATIBILITY OF ADJUVANTS AND *TOURNEYEZ* FUNGICIDE

Perform a jar test before mixing commercial quantities of *TourneyEZ* Fungicide, when using this product for the first time, when using new adjuvants, when using new tank mixes, or when using a new water source.

1. Add 1 pint of the water to a quart jar. Use water from the same source and temperature as that to be used in the spray tank.
2. Add 2.5 mL (1/2 teaspoon) of *TourneyEZ* Fungicide to the quart jar, gently mix until product goes into suspension.
3. Add 1 mL of new adjuvant and/or appropriate amount of new tank mix partner and gently mix.
4. Place cap on jar, invert 10 times, let stand for 15 minutes, evaluate.
5. An acceptable tank mix combination will have a smooth, uniform appearance. If any of the following conditions are observed, question the choice of spray mix components:
 - a) Layer of oil or globules on the mixture's surface.
 - b) Flocculation: Formation of fluffy, cloudlike aggregates or masses in suspension or as a layer on the bottom of the jar.
 - c) Clabbering: Thickening texture (coagulated) like gelatin or cottage cheese.

TANK MIX PARTNERS

TourneyEZ Fungicide is compatible with most herbicides, insecticides, fungicides and other types of tank mix components. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. **DO NOT** tank mix with any product which has a prohibition on tank mixing. When evaluating potential tank mix partners, always apply to a small area of target plants and then monitor for adverse effects for at least two weeks before making large-scale applications.

SPRAYER PREPARATION

Before applying *TourneyEZ* Fungicide, start with clean, well-maintained application equipment. The spray tank hoses and

booms must be cleaned to ensure no residue from the previous spraying operations remain in the sprayer. The spray equipment must be cleaned according to the manufacturer's directions for the last product used before the equipment is used to apply *Tourney EZ* Fungicide. If two or more products were tank mixed prior to *Tourney EZ* Fungicide application, the most restrictive cleanup procedure must be followed.

APPLICATION EQUIPMENT

Application equipment must be clean and in good repair. Check nozzles frequently for accuracy.

SPRAYER CLEANUP

Clean sprayer equipment each day following *Tourney EZ* Fungicide application. After application is complete, use the following steps to clean spray equipment:

1. Completely drain the spray tank, rinse the sprayer thoroughly, including the inside and outside of the tank and all in-line screens.
2. Fill the spray tank with clean water and flush all hoses, booms, screens, and nozzles.
3. Drain tank completely.
4. Remove all nozzles and screens and rinse them in clean water.

MIXING INSTRUCTIONS

1. Fill clean spray tank 1/2 to 2/3 of desired level with clean water.
2. While agitating, slowly add the *Tourney EZ* Fungicide to the spray tank. Agitate to create a rippling or rolling action on the water surface.
3. If tank mixing *Tourney EZ* Fungicide with other labeled pesticides, add water soluble bags first, followed by dry formulations, flowables, emulsifiable concentrates and then solutions.
4. If tank mixing *Tourney EZ* Fungicide with other labeled pesticides, follow more restrictive limitations or cautions on labels of all products. **DO NOT** tank mix with any products which contain a prohibition on tank mixing.
5. Add any required adjuvants.
6. Fill spray tank to desired level with water. Continue agitation until all spray solution has been applied.
7. Mix only the amount of spray solution that can be applied the day of mixing. Apply *Tourney EZ* Fungicide within 24 hours of mixing.

MANDATORY SPRAY DRIFT MANAGEMENT

Ground Applications:

- Apply with the nozzle height recommended by the manufacturer, but no more than 3 feet above the ground or crop canopy.
- For all applications, applicators are required to use a medium or coarser spray droplet size (ASABE S572.1).
- **DO NOT** apply when wind speeds exceed 10 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

Boomless Ground Applications:

- Applicators are required to use a medium or coarser droplet size (ASABE S572.1) for all applications.
- **DO NOT** apply when wind speeds exceed 10 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.

BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size – Ground Boom

- Volume – Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure – Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle – Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

BOOM HEIGHT – Ground Boom

For ground equipment, the boom should remain level with the crop and have minimal bounce.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.

Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

Boomless Ground Applications:

Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

Handheld Technology Applications:

Take precautions to minimize spray drift.

TURFGRASS

- Use *Tourney EZ* Fungicide for disease control on golf courses, lawns and landscape areas around residential, institutional, public, commercial and industrial buildings, parks, recreational areas, athletic fields, and sod farms.
- Use *Tourney EZ* Fungicide in conjunction with turf management practices that promote good plant health and

optimum disease control. Proper identification of disease infestations is essential for providing control. Consult diagnostic guides, kits, or other means of identification to determine the best control measures.

- *Tourney EZ* Fungicide is more effective when used preventively or early in disease development. Length of disease control will vary according to local environmental conditions, disease pressure and turfgrass management practices.
- Apply to dry turf or after mowing. For control of foliar diseases, allow turf to dry before irrigation. For control of soilborne diseases, *Tourney EZ* Fungicide can be watered-in after application.
- *Tourney EZ* Fungicide may be tank mixed with other registered turfgrass fungicides. Perform a jar test as described above to determine compatibility before making applications. Directions for use of *Tourney EZ* Fungicide have been developed using no adjuvant in the spray solution.
- *Tourney EZ* Fungicide can injure golf course greens that are under high heat (temperatures exceeding 90°F) and/or drought stress. Under high heat and/or drought stress conditions (mid to late summer), **DO NOT** apply more than 16 fl oz per acre (0.50 lb ai/acre) (0.37 fl oz per 1,000 sq ft) (0.12 lb ai/1,000 sq ft) every 30 days.
- Important: Under conditions of optimal disease development (including high humidity and temperature), use higher labeled rates and shorter application intervals. When applications are made curatively, use higher specified rates of *Tourney EZ* Fungicide and shorter specified application intervals. Use the higher specified application rate in areas with a history of severe disease pressure.
- Failure to follow label directions may result in turfgrass injury and/or inferior disease control.

APPLICATION DIRECTIONS

Apply *Tourney EZ* Fungicide using standard, low-pressure spray equipment designed to deliver coarse- to medium-size spray droplets (**DO NOT** use nozzles designed to deliver fine droplets). Apply in a sufficient volume of water to provide thorough spray coverage and a uniform spray pattern. **DO NOT** position the spray boom more than 20 inches above the turf. For foliar disease, apply 2 gallons of spray solution per 1,000 sq ft. For crown and root diseases,

apply 2 to 4 gallons of spray solution per 1,000 sq ft. For spot treatment, mix 0.1 to 0.2 fl oz (3-6 mL) (0.003-0.006 lb ai) of *Tourney EZ Fungicide* per gallon of water.

TURFGRASS TOLERANCE

Due to the large number of turfgrass types, possible tank mixes and application techniques, it has not been determined if *Tourney EZ Fungicide* can be safely used on all turfgrasses under all conditions. Before making a large-scale application, determine safety in a small test area. Monitor the test area for at least 2 weeks to determine safety under local growing conditions.

TURFGRASS SPECIES

Annual Bluegrass
Bentgrass
Bermudagrass
Centipedegrass
Fescue species
Kentucky Bluegrass
Perennial Ryegrass
St. Augustinegrass
Seashore Paspalum
Zoysiagrass

TURFGRASS RESTRICTIONS

- **DO NOT** apply more than 1/2 gallon (2 lb ai) of *TourneyEZ Fungicide* per acre per year.
- **DO NOT** apply directly to bermudagrass greens.
- **DO NOT** apply through any type of irrigation system.
- **DO NOT** apply by air.
- **DO NOT** graze animals on treated areas for one year after application.
- **DO NOT** feed clippings from treated areas to livestock or poultry.
- **DO NOT** exceed 0.2 fl oz (6 mL) (0.006 lb ai) product per gallon of water for spot treatment.
- **DO NOT** apply more than 19.2 fl oz/A (0.6 lb ai/A) per application.
- **DO NOT** make sequential applications sooner than 14 days after previous application of *Tourney EZ Fungicide*: See Crop Specific Directions, Restrictions and Limitations for details.
- **DO NOT** exceed 8 applications per year.

Tourney EZ Fungicide Rate Conversion Chart for Turfgrass

fl oz product/1,000 sq ft	lb ai/1,000 sq ft	fl oz product/acre	pints product/acre	lb ai/acre
0.18	0.006	8	0.5	0.25
0.28	0.009	12	0.75	0.38
0.37	0.012	16	1.0	0.50
0.44	0.014	19.2	1.2	0.60

CROP SPECIFIC DIRECTIONS, RESTRICTIONS AND LIMITATIONS

TURFGRASS				
Diseases (Pathogens)	Application Rates		Application Interval/ Timing	Special Instructions
	fl oz/1,000 sq ft	fl oz/A		
Dollar Spot (<i>Sclerotinia homoeocarpa</i>)	0.18 to 0.37	8 to 16	14 to 21 days	Use preventively. Begin application when conditions favor disease development.
Brown Patch (<i>Rhizoctonia solani</i>) Anthracnose (<i>Colletotrichum graminicola</i>)	0.28 to 0.37	12 to 16	14 to 21 days	Begin applications when conditions favor disease development.
Gray Leaf Spot (<i>Pyricularia grisea</i>) Necrotic Ring Spot (<i>Ophiostoma korrae</i>) Red Thread (<i>Laetisaria fuciformis</i>) Rhizoctonia Large Patch (<i>Rhizoctonia solani</i>) Rust Diseases (<i>Puccinia</i> spp.) Summer Patch (<i>Magnaporthe poae</i>) Take-all Patch (<i>Gaeumannomyces graminis</i>) Waitea Patch/ Brown Ring Patch (<i>Waitea circinata</i>) Zoysia Patch (<i>Rhizoctonia solani</i>)	0.37	16	14 days	Begin applications when conditions favor disease development or when the disease first appears.

(continued)

CROP SPECIFIC DIRECTIONS, RESTRICTIONS AND LIMITATIONS (continued)

TURFGRASS				
Diseases (Pathogens)	Application Rates		Application Interval/ Timing	Special Instructions
	fl oz/ 1,000 sq ft	fl oz/A		
Fairy Ring (Various <i>Basidiomycetes</i>)	0.37	16	21 days	Apply as soon as possible after first symptoms appear. Apply in a volume of 4 gal per 1,000 sq ft. For optimal control, reapply after 21 days. Symptoms may take several weeks to disappear following application.
Pink Snow Mold/ Fusarium Patch (<i>Microdochium nivale</i>) Gray Snow Mold (<i>Typhula</i> spp.) Yellow Patch (<i>Rhizoctonia cerealis</i>)	0.37 to 0.44	16 to 19.2	Late fall	Make 1 to 2 applications in combination with PCNB, chlorothalonil or other fungicides active on snow mold in late fall before snow cover and if possible, one application during a mid-winter thaw.

ORNAMENTAL PLANTS

Do not use in California

- *Tourney* EZ Fungicide provides disease control on ornamental plants in commercial outdoor nurseries and ornamental plants found in landscaped areas around residential, institutional, public, commercial and industrial buildings, parks, recreational areas and athletic fields.
- *Tourney* EZ Fungicide provides disease control on ornamental plants in indoor greenhouses and nurseries.
- Use *Tourney* EZ Fungicide as part of a preventive disease control program.
- *Tourney* EZ Fungicide is more effective when used preventively or early in disease development. Length of disease control will vary according to local environmental conditions, disease pressure and management practices.
- Follow all use directions on this label. Failure to follow label directions may result in injury and/or inferior disease control.

ORNAMENTAL PLANT TOLERANCE

IMPORTANT: *Tourney* EZ Fungicide may cause growth regulation to some ornamental plant species. The large number of existing ornamental species and their varieties and cultivars coupled with the constant introduction of new varieties makes it impossible to field test *Tourney* EZ Fungicide in every locale where it is sold or in all of the combinations of location and plant variety. Further differences include the soil or media type, pH, moisture or fertility, environmental conditions including temperature, lighting or degree-days and horticultural practice and the manner of use and application of this product. To ensure that *Tourney* EZ Fungicide is compatible with the ornamental plant variety or cultivar under your specific conditions, test this product on a small test area and observe for phytotoxicity or other unintended effects for at least two weeks before making large-scale applications.

APPLICATION DIRECTIONS

Foliar applications

- Apply *TourneyEZ* Fungicide at 1 to 4 fl oz (0.03-0.125 lb ai) of product per 100 gallons of water for disease control in ornamental plants.
- Apply as a full coverage spray to the point of run off.
- Apply as needed on a 14- to 28-day interval.
- Under conditions of optimal disease development (for example high humidity and temperature), use higher specified rates and shorter specified application intervals.
- When applications are made curatively, use higher specified rates of *TourneyEZ* Fungicide and shorter specified application intervals. Use the higher specified application rate in areas with a history of severe disease pressure.
- For disease control in landscapes, apply 1 to 4 fl oz (0.03-0.125 lb ai) of product per 100 gallons of water on a 21- to 28-day interval.

Drench applications

- To control soilborne, seedling and crown diseases, apply *TourneyEZ* Fungicide as a preventive drench treatment prior to disease development.
- Apply at 1 to 4 fl oz (0.03-0.125 lb ai) per 100 gallon of water in a volume of 1 to 2 pints of solution per square foot of surface area on a 14- to 28-day interval.
- Roots must be healthy to ensure systemic uptake, translocation, and disease protection.

COMPATIBILITY

TourneyEZ Fungicide is compatible with most commonly used insecticides, fungicides, and spray adjuvants used in the production of ornamental plants. When using *TourneyEZ* Fungicide in tank mixes with other pesticides, observe all directions for use and precautions on the label of tank mix partner. Before applying a tank mix for the first time, check physical compatibility of the mixture.

DO NOT tank mix with any product which contains a prohibition on tank mixing.

APPLICATION EQUIPMENT

Apply *TourneyEZ* Fungicide using suitable hand- or power-operated application equipment including portable, pump-up, backpack, hydraulic or boom sprayers in a manner to provide complete and uniform coverage.

ORNAMENTAL RESTRICTIONS

- **DO NOT** apply more than 1/2 gallon (2 lb ai) of *TourneyEZ* Fungicide per acre per year.
- **DO NOT** apply through any type of irrigation system.
- **DO NOT** apply by air.
- **DO NOT** graze animals on treated areas of one year after application.
- **DO NOT** feed plants from treated areas to livestock or poultry.
- **DO NOT** apply more than 4 fl oz/100 gallons (0.125 lb ai/100 gallons) per application.
- **DO NOT** make sequential applications sooner than 14 days after previous application of *TourneyEZ* Fungicide: See Crop Specific Directions, Restrictions and Limitations for details.
- **DO NOT** exceed 26 applications per year (at the lowest rate).
- **DO NOT** use in California.

TourneyEZ Ornamental rate summary

fl oz/100 gal of spray volume	lb ai/100 gal of spray volume
1	0.031
2	0.063
3	0.094
4	0.125

DIRECTIONS FOR USE ON ORNAMENTALS Not for use in California

Diseases*	Pathogen	Application Rates	Special Instructions
Anthracnose	<i>Apiognomonia</i> spp. <i>Colletotrichum</i> spp. <i>Cryptocline</i> spp. <i>Discula</i> spp. <i>Gleosporium</i> spp.	1 to 4 fl oz/100 gal on a 14- to 28-day interval	When using a typical high-volume sprayer use 100 gal of spray mix to treat 20,000 sq ft of area. If using a low-volume sprayer, adjust concentration to apply the same amount of product per unit area.
Botrytis	<i>Botrytis</i> spp.		
Blossom Rot/Brown Rot	<i>Monilinia</i> spp.		
Boxwood Blight	<i>Calonectria pseudonaviculatum</i> <i>Cylindrocladium buxicola</i> <i>Cylindrocladium pseudonaviculatum</i>	4 fl oz/100 gal on a 14- to 28-day interval	
Conifer Blight	<i>Diplodia pinea</i> <i>Phomopsis</i> spp. <i>Rhabdocline</i> spp. <i>Rhizosphaera</i> spp. <i>Sphaeropsis</i> spp.	1 to 4 fl oz/100 gal on a 14- to 28-day interval	
Damping-Off/ Stem and Root Rots	<i>Rhizoctonia solani</i>	1 to 4 fl oz/100 gal on a 14- to 28-day interval	
Eastern Filbert Blight	<i>Anisogramma anomala</i>		
Fusarium Root Rot and Damping Off	<i>Fusarium oxysporum</i> <i>Fusarium solani</i>	1 to 4 fl oz/100 gal on a 14- to 28-day interval	

(continued)

*If symptoms are visible of a labeled disease, use a registered contact, protectant fungicide.

DIRECTIONS FOR USE ON ORNAMENTALS Not for use in California (continued)

Diseases*	Pathogen	Application Rates	Special Instructions
Leaf Spots	<i>Alternaria</i> spp. <i>Cercospora</i> spp. <i>Diplocarpon</i> spp. <i>Mycosphaerella</i> spp. <i>Myrothecium</i> spp. <i>Phyllosticta</i> spp. <i>Septoria</i> spp.	1 to 4 fl oz/100 gal on a 14- to 28-day interval	When using a typical high-volume sprayer use 100 gal of spray mix to treat 20,000 sq ft of area. If using a low-volume sprayer, adjust concentration to apply the same amount of product per unit area.
Powdery Mildew	<i>Erysiphe</i> spp. <i>Microsphaera</i> spp. <i>Oidium</i> spp. <i>Phyllactinia</i> spp. <i>Podosphaera</i> spp. <i>Sphaerotheca</i> spp.		
Rusts	<i>Chrysomyxa</i> spp. <i>Coleosporium</i> spp. <i>Gymnosporangium</i> spp. <i>Puccinia</i> spp. <i>Uredo</i> spp. <i>Uromyces</i> spp.		
Scab	<i>Cladosporium</i> spp. <i>Sphaceloma</i> spp. <i>Venturia</i> spp.	2 to 4 fl oz/100 gal on a 14- to 28-day interval	
Shot Hole	<i>Wilsonomyces carpophilus</i> <i>Blumeriella</i> spp.		
White Mold	<i>Sclerotinia sclerotium</i>		

*If symptoms are visible of a labeled disease, use a registered contact, protectant fungicide.

ORNAMENTAL PLANT LIST

Note: *Tourney* EZ Fungicide has been tested on the following plants with no crop injury.

Flowering and Foliage Plants	Herbaceous Ornamental Plants	Woody Ornamental Plants		Non-Bearing Tree, Fruit, Nut and Vine
Geranium Petunia Poinsettia Snapdragon	Carnation Chrysanthemum Daylily Dianthus Hydrangea Iris Lantana Liriope Marigold Phlox Verbena	Amelanchier Ash Azalea Bayberry Beech Boxwood Birch Camellia Conifer Cotoneaster Crabapple Crape Myrtle Dogwood Elm Euonymus Fir Hawthorn Hemlock Holly Juniper	Laurel Lilac Linden London Plane Magnolia Maple Mountain Ash Mountain Laurel Oak Pine Poplar Photinia Pyracantha Rhododendron Rose Sassafras Spirea Spruce Sweetgum Sycamore	Apple Apricot Cherry Citrus Filbert Grape Peach Pear Pecan Pistachio Plum Walnut

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE

Store in a cool dry place.

Keep pesticide in original container.

Keep container closed when not in use.

DO NOT put dilute into food or drink containers.

DO NOT store in or around the home.

PESTICIDE DISPOSAL

Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING

Nonrefillable container. **DO NOT** reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

RECYCLING

Once cleaned, some agricultural plastic pesticide containers can be taken to a container collection site or picked up for recycling. To find the nearest site, contact your chemical dealer or manufacturer, or contact the Ag Container Recycling Council (ACRC) at 877-952-2272 (toll free) or www.acrecycle.org.

©2021 Valent U.S.A. LLC

Tourney and Products That Work, From People Who Care
are registered trademarks of Valent U.S.A. LLC

Manufactured for:

Valent U.S.A. LLC

P.O. Box 5075

San Ramon CA 94583

Made in U.S.A.

Form 2376-A

EPA Reg. No. 59639-257

EPA Est. 11773-IA-1[®], 228-IL-1[®], 39578-TX-1[®],

5481-ID-1[®], 62171-MS-1[®], 70815-GA-2[®]

Superscript is first letter of lot number.

059639-00257.20210816.TourneyEZ.NPR.FINAL

SAL20210819

Information contained in this booklet is accurate at the time of printing. Since product testing is a continuous process, please read and follow the directions on the product label for the most current directions and precautionary statements.

Always check with your state to verify state registration status or call 800-89-VALENT (898-2536).



For state registration and/or supplemental labels, please call or visit us online.

Products That Work, From People Who Care[®] | valentpro.com | 800-89-VALENT (898-2536)

Always read and follow label instructions.

©2021 Valent U.S.A. LLC. All rights reserved. Printed in the U.S.A.

Date 11/24/21