

MEFENOXAM GROUP 4 FUNGICIDE

PYDIFLUMETOFEN GROUP 7 FUNGICIDE

PULL HERE TO OPEN ▶

SEDAXANE GROUP 7 FUNGICIDE

AZOXYSTROBIN GROUP 11 FUNGICIDE

FLUDIOXONIL GROUP 12 FUNGICIDE



# Trebuset®

Peanuts Colorless



## Fungicide

ADEPIDYN® Technology<sup>1</sup>

*Active Ingredients:*

Azoxystrobin <sup>2</sup> .....	11.0%
Pydiflumetofen <sup>3</sup> .....	6.85%
Fludioxonil <sup>4</sup> .....	3.47%
Sedaxane <sup>5</sup> .....	3.34%
Mefenoxam <sup>6</sup> .....	0.64%

*Other Ingredients:* 74.7%

**Total:** 100.00%

<sup>1</sup>Technology denotes the active ingredient Pydiflumetofen.

<sup>2</sup>CAS No. 131860-33-8

<sup>3</sup>CAS No. 1228284-64-7

<sup>4</sup>CAS No. 131341-86-1

<sup>5</sup>CAS No. 874967-67-6

<sup>6</sup>CAS No. 70630-17-0 and 69516-34-3

Trebuset Peanuts Colorless is a flowable concentrate for seed treatment containing 1.03 lb azoxystrobin, 0.64 lb pydiflumetofen, 0.32 lb fludioxonil, 0.31 lb sedaxane, and 0.060 lb mefenoxam per gallon.

**KEEP OUT OF REACH OF CHILDREN.**

## CAUTION

See additional precautionary statements and directions for use inside booklet.

EPA Reg. No. 100-1689 EPA Est. 100-NE-001

Product ID **72355**

SCP 1689A-L1A 0122 4153336

**15 gallons**  
Net Contents



®

## **TABLE OF CONTENTS**

- 1.0 FIRST AID**
- 2.0 PRECAUTIONARY STATEMENTS**
  - 2.1 Hazards to Humans and Domestic Animals**
  - 2.2 Personal Protective Equipment (PPE)**
    - 2.2.1 User Safety Requirements
    - 2.2.2 Engineering Controls
    - 2.2.3 User Safety Recommendations
  - 2.3 Environmental Hazards**
    - 2.3.1 Groundwater Advisory
    - 2.3.2 Surface Water Advisory
  - 2.4 Physical or Chemical Hazards**

## **DIRECTIONS FOR USE**

- 3.0 PRODUCT INFORMATION**
  - 3.1 Resistance Management**
- 4.0 APPLICATION DIRECTIONS**
  - 4.1 Mixing Directions**
    - 4.1.1 Tank-Mix Precautions
    - 4.1.2 Tank-Mix Recommendation
- 5.0 RESTRICTIONS**
- 6.0 SEED CONTAINER LABEL REQUIREMENTS**
- 7.0 Seed Treatment Directions**
  - 7.1 Peanut**
- 8.0 STORAGE AND DISPOSAL**
- 9.0 CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY**

## 1.0 FIRST AID

FIRST AID	
<b>If swallowed</b>	<ul style="list-style-type: none"><li>• Call a poison control center or doctor immediately for treatment advice.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li><li>• Do not give anything by mouth to an unconscious person.</li></ul>
<b>If on skin or clothing</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
<b>HOTLINE NUMBER</b> For 24-Hour Medical Emergency Assistance (Human or Animal) Or Chemical Emergency Assistance (Spill, Leak, Fire or Accident) Call <b>1-800-888-8372</b>	

## 2.0 PRECAUTIONARY STATEMENTS

### 2.1 Hazards to Humans and Domestic Animals

#### CAUTION

Harmful if swallowed or absorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

### 2.2 Personal Protective Equipment (PPE)

**Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Water-resistant gloves
- Shoes plus socks

#### 2.2.1 USER SAFETY REQUIREMENTS

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### 2.2.2 ENGINEERING CONTROLS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240(d)(4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS. IMPORTANT: When reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for "applicators and other handlers" and have such PPE immediately available for use in an emergency, such as a spill or equipment breakdown.

#### 2.2.3 USER SAFETY RECOMMENDATIONS

<b>Users should:</b> <ul style="list-style-type: none"><li>• Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.</li><li>• Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.</li></ul>
---

## 2.3 Environmental Hazards

This product is toxic to fish, aquatic invertebrates, oysters, and shrimp. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. **DO NOT** apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwater or rinsate.

### 2.3.1 GROUNDWATER ADVISORY

Azoxystrobin, a degradate of azoxystrobin, and mefenoxam are known to leach through soil into groundwater under certain conditions as a result of label use. Fludioxonil and pydiflumetofen have properties and characteristics associated with chemicals detected in groundwater. These chemicals may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

### 2.3.2 SURFACE WATER ADVISORY

This product may impact surface water quality due to runoff of rainwater. This is especially true for poorly draining soils and soils with shallow groundwater. This product is classified as having a medium potential for reaching both surface water and aquatic sediment via runoff several months or more after application.

A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of pydiflumetofen from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours. Sound erosion control practices will reduce this product's potential to reach aquatic sediment via runoff.

## 2.4 Physical or Chemical Hazards

**DO NOT** mix or allow to come in contact with oxidizing agents. Hazardous chemical reaction may occur.

## DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. Notify State and/or Federal authorities and Syngenta immediately if you observe any adverse environmental effects due to use of this product.

**Maximum usage when applying metalaxyl- and mefenoxam-containing products to the same crop within the same season: DO NOT** apply more than the maximum yearly total application rate for the active ingredient as stated on the label of the product containing the lowest yearly total on that crop.

**FAILURE TO FOLLOW DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY; POOR DISEASE CONTROL; AND/OR ILLEGAL RESIDUES.**

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard (WPS).

**DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours.**

**Exception:** If the seed is treated with the product and the treated seed is soil-injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Water-resistant gloves
- Shoes plus socks

Treatment of highly mechanically scarred or damaged seed or seed known to be of low vigor and poor quality may result in reduced germination and/or reduction of seed and seedling vigor. Treat a quantity of seed using equipment similar to that planned for treating the total seed lot. Then conduct germination tests with a portion of this treated seed before committing the total seed lot to a selected seed treatment.

Due to seed quality, crop or variety sensitivity, and seed storage conditions beyond the control of Syngenta, no claims are made to guarantee the germination of carry-over seed or propagating material for all crops when treated with Trebuset Peanuts Colorless.

### 3.0 PRODUCT INFORMATION

Trebuset Peanuts Colorless is a seed treatment fungicide effective against seedborne and soilborne fungi causing decay, damping off, seedling blight, and *Cylindrocladium* black rot (CBR). Trebuset Peanuts Colorless also provides suppression of White Mold/Stem Rot (*Sclerotium rolfsii*) and Aspergillus Crown Rot (*Aspergillus niger*)

Read all label directions for use. All applications must be made according to the use directions that follow.

#### 3.1 Resistance Management

MEFENOXAM	GROUP	4	FUNGICIDE
PYDIFLUMETOFEN	GROUP	7	FUNGICIDE
SEDAXANE	GROUP	7	FUNGICIDE
AZOXYSTROBIN	GROUP	11	FUNGICIDE
FLUDIOXONIL	GROUP	12	FUNGICIDE

For resistance management, please note that Trebuset Peanuts Colorless contains Group 4/mefenoxam, Group 7/pydiflumetofen, Group 7/sedaxane, Group 11/azoxystrobin, and Group 12/fludioxonil fungicides. Any fungal population may contain individuals naturally resistant to Trebuset Peanuts Colorless and other Group 4, Group 7, Group 11, or Group 12 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

Mefenoxam belongs to the phenylamide class of chemistry which interferes with fungal RNA synthesis. Pydiflumetofen and sedaxane belong to the carboxamide class of chemistry and are succinate dehydrogenase inhibitors (SDHIs) which disrupt cellular respiration and energy generation. Azoxystrobin belongs to the strobilurin class of chemistry which disrupts cellular respiration and energy generation. Fludioxonil belongs to the phenylpyrrole class of chemistry which interferes with osmotic signal transduction.

To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of Trebuset Peanuts Colorless or other Group 4, Group 7, Group 11, or Group 12 fungicides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicides from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
- For further information or to report suspected resistance contact Syngenta Crop Protection at 1-866-796-4368. You can also contact your university extension specialist to report resistance.

## 4.0 APPLICATION DIRECTIONS

**Important:** Recirculate Trebuset Peanuts Colorless thoroughly before using.

Follow the manufacturer's application instructions for the seed treatment equipment being used. This product must be used only by commercial seed treatment facilities or with commercial seed treatment equipment on farm. **DO NOT** use in hopper-box, planter-box, slurry-box or other seed treatment applications at or immediately before planting.

Apply Trebuset Peanuts Colorless as a water-based slurry through standard liquid-type seed treatment equipment that provides uniform seed coverage. Seed treaters with atomizers or spinning discs are highly recommended for better product coverage on the seed. Uneven or incomplete seed coverage may not give the desired level of disease control. Thoroughly mix the specified amount of Trebuset Peanuts Colorless into the required amount of water for the slurry treater and dilution rate to be used.

The total application volume must be sufficient to provide desired level of coverage. The slurry volume needed to achieve adequate coverage is approximately 11-16 fl oz/100 lb of seed. More or less diluent may be required depending on environmental conditions and peanut seed size in order to obtain optimal coverage.

Fill the mix tank  $\frac{1}{4}$ - $\frac{1}{2}$  full with clean water. Start agitation of the mix tank. Next, open the container and add the appropriate amount of Trebuset Peanuts Colorless into the mix tank, and maintain agitation. Thoroughly blend the mixture while adding the remainder of the required water.

After the mixture appears completely uniform, begin application to the seed. Mix the slurry thoroughly with the seed to provide uniform coverage.

Continuous agitation or mixing of the slurry mixture is necessary to prevent settling out of the solution. Clean out any unused product from the treater after treating or maintain constant agitation if the leftover slurry will be maintained overnight.

Follow planter manufacturer's specifications for use of talc or other hopper additives at planting. Seed must be completely dry before adding to planter.

### 4.1 Mixing Directions

Trebuset Peanuts Colorless mixes easily with water and other water-based seed treatments manufactured by Syngenta and many other companies. When mixing with products from other manufacturers, test the compatibility prior to use by conducting a jar test:

- Mix all intended seed treatments with a proportional amount of water to achieve the desired slurry mixture in a clear glass container.
- Mix well and allow mixture to sit for one hour.
- Remix and observe for incompatibility.

Mixing Trebuset Peanuts Colorless with tank-mix partners:

- Add  $\frac{1}{2}$  of the required water to the mix tank and turn on the agitation. Mechanical agitation is preferred.
- If using wettable powders, add them first to clean water allowing them to completely disperse prior to adding Trebuset Peanuts Colorless or other products.
- Allow each tank-mix partner to completely disperse before adding the next product.
- Add the remaining amount of water and agitate.
- Maintain agitation until the entire slurry mixture has been used.

This product does not contain dye and treated seed is not covered by an appropriate tolerance, tolerance exemption, or other clearance under the Federal Food, Drug and Cosmetic Act. To comply with 40 CFR 153.155, therefore, all seed treated commercially with this product must be colored with an EPA-approved dye or colorant of a suitable color to prevent accidental use as food for man or feed for animals.

**Allow seed to dry before bagging.**

#### 4.1.1 TANK-MIX PRECAUTIONS

Confirm physical compatibility, seed safety, seed safety in stored seed, and biological efficacy before tank mixing with other seed treatment products.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

#### 4.1.2 TANK-MIX RECOMMENDATION

For additional suppression of Aspergillus Crown Rot, Trebuset Peanuts Colorless may be combined with a seed treatment product containing difenoconazole (e.g., Dividend®).

### 5.0 RESTRICTIONS

- Store treated seeds away from feed and foodstuffs.
- **DO NOT** use for feed, food, or oil purposes.
- **DO NOT** allow children, pets, or livestock to have access to treated seeds.
- Wear long-sleeved shirt, long pants, socks, shoes and water-resistant gloves when handling treated seed.
- Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading and planting.
- Dispose of all excess treated seed. Left over treated seed may be double sown around the headland or buried away from water sources in accordance with local requirements.
- **DO NOT** contaminate water bodies when disposing of planting equipment washwater or rinsate.
- Treated seed must be planted into the soil at a depth of at least 1 inch.
- Dispose of seed packaging in accordance with local requirements.
- **DO NOT** use at a rate that will result in more than 0.073 lb (33 g) azoxystrobin per acre per calendar year as a seed treatment application. This seed has been treated with 0.086 mg ai azoxystrobin per seed.
- **DO NOT** apply more than 0.18 lb (82 g) of pydiflumetofen per acre per calendar year as a seed treatment. This seed has been treated with 0.053 mg ai pydiflumetofen per seed.
- **DO NOT** apply more than 0.023 lb (10.5 g) of fludioxonil per acre per calendar year as a seed treatment. This seed has been treated with 0.027 mg ai fludioxonil per seed.
- **DO NOT** apply more than 0.071 lb (32 g) of mefenoxam per acre per calendar year as a seed treatment. This seed has been treated with 0.005 mg ai mefenoxam per seed.
- **DO NOT** apply more than 0.046 lb (21 g) of sedaxane per acre per calendar year as a seed treatment. This seed has been treated with 0.026 mg ai sedaxane per seed.
- The maximum allowed number of treated seeds is 193,800 treated seeds per acre (equivalent to 228 pounds of treated seed per acre).
- This treated seed may be planted on the same acres up to 2 times per year.

- In the event of a crop failure or harvest of a crop grown from seed treated with Trebuset Peanuts Colorless, the field may be replanted according to the following schedule:

Crop	Plant-Back Interval
Cereals, Small Grain (barley, oats, rye, triticale, wheat) Corn (field, pop, seed, sweet) Cottonseed Legume Vegetables, Succulent or Dried, Crop Group 6, Except Cowpea Forage and Cowpea Hay Peanut Potato Rapeseed (including canola), Crop Subgroup 20A Sorghum Soybean, Except Forage, Hay, and Silage Sugarbeet	0 days
Brassica Head and Stem Vegetables, Crop Group 5-16 Brassica Leafy Greens, Crop Subgroup 4-16B Bulb Vegetables, Crop Group 3-07 Cucurbit Vegetables, Crop Group 9 Fruiting Vegetables, Crop Group 8-10 Grasses Grown for Seed Leaf Petiole Vegetables, Crop Subgroup 22B Leafy Greens, Crop Subgroup 4-16A Leaves of Root and Tuber Vegetables, Crop Group 2 Non-grass Animal Feeds, Crop Group 18 Quinoa Rice Root Vegetables, Crop Subgroup 1A Strawberry Sunflower, Crop Subgroup 20B Tobacco Tuberous and Corm Vegetables, Crop Subgroup 1C	30 days
All Other Crops Intended for Food and Feed	365 days



## 6.0 SEED CONTAINER LABEL REQUIREMENTS

The Federal Seed Act requires that bags containing treated seeds shall be labeled with the following statements:

- This seed has been treated with azoxystrobin, fludioxonil, mefenoxam, pydiflumetofen, and sedaxane fungicides.
- **DO NOT** use for feed, food, or oil purposes.

In addition, the U.S. Environmental Protection Agency requires the following statements on bags containing seed treated with Trebuset Peanuts Colorless fungicide:

- Store away from feed and foodstuffs.
- **DO NOT** allow children, pets, or livestock to have access to treated seeds.
- Wear long-sleeved shirt, long pants, shoes, socks, and water-resistant gloves when handling treated seed.
- Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading and planting.
- Dispose of all excess treated seed. Left over treated seed may be double sown around the headland or buried away from water sources in accordance with local requirements.
- **DO NOT** contaminate water bodies when disposing of planting equipment washwater or rinsate.
- **Groundwater Advisory:** Azoxystrobin, a degradate of azoxystrobin and mefenoxam are known to leach through soil into groundwater under certain conditions as a result of label use. Fludioxonil and Pydiflumetofen have properties and characteristics associated with chemicals detected in groundwater. These chemicals may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.
- Treated seed must be planted into the soil at a depth of at least 1 inch.
- Dispose of seed packaging in accordance with local requirements.
- **DO NOT** exceed 0.073 lb ai/A/year of azoxystrobin-containing products as a seed treatment.
- **DO NOT** exceed 0.18 lb ai/A/year of pydiflumetofen-containing products through any combination of seed and foliar applications.
- **DO NOT** exceed 0.023 lb ai/A/year of fludioxonil-containing products as a seed treatment.
- **DO NOT** exceed 0.071 lb ai/A/year of mefenoxam- or metalaxyl-containing products as a seed treatment.
- **DO NOT** exceed 0.046 lb ai/A/year of sedaxane-containing products as a seed treatment.
- This treated seed may be planted on the same acres up to 2 times per year.

- In the event of crop failure or harvest of a crop grown from seed treated with Trebuset Peanuts Colorless, crops may be replanted according to the following schedule:

Crop	Plant-Back Interval
Cereals, Small Grain (barley, oats, rye, triticale, wheat) Corn (field, pop, seed, sweet) Cottonseed Legume Vegetables, Succulent or Dried, Crop Group 6, Except Cowpea Forage and Cowpea Hay Peanut Potato Rapeseed (including canola), Crop Subgroup 20A Sorghum Soybean, Except Forage, Hay, and Silage Sugarbeet	0 days
Brassica Head and Stem Vegetables, Crop Group 5-16 Brassica Leafy Greens, Crop Subgroup 4-16B Bulb Vegetables, Crop Group 3-07 Cucurbit Vegetables, Crop Group 9 Fruiting Vegetables, Crop Group 8-10 Grasses Grown for Seed Leaf Petiole Vegetables, Crop Subgroup 22B Leafy Greens, Crop Subgroup 4-16A Leaves of Root and Tuber Vegetables, Crop Group 2 Non-grass Animal Feeds, Crop Group 18 Quinoa Rice Root Vegetables, Crop Subgroup 1A Strawberry Sunflower, Crop Subgroup 20B Tobacco Tuberous and Corm Vegetables, Crop Subgroup 1C	30 days
All Other Crops Intended for Food and Feed	365 days

## 7.0 Seed Treatment Directions

### 7.1 Peanut

Crop			
Peanut			
Target Disease/Pest	Use Rate*		Application Instructions
	fl oz product/ 100 lb seed	mg ai/seed	
Seedborne and Soilborne Fungi Causing Decay, Damping-off, and Seedling Blight  Seedling Damping-off (Fusarium, Penicillium, Pythium, and Rhizoctonia)  Cylindrocladium black rot (CBR)  <i>Suppression Only:</i> White Mold/Stem Rot ( <i>Sclerotium rolfsii</i> )  <i>Suppression Only:</i> Aspergillus Crown Rot ( <i>Aspergillus flavus</i> and <i>Aspergillus niger</i> )	2.0	Azoxystrobin: 0.086 Pydiflumetofen: 0.053 Fludioxonil: 0.027 Sedaxane: 0.026 Mefenoxam: 0.005  Total: 0.197	For additional suppression of Aspergillus Crown Rot, Trebuset Peanuts Colorless may be mixed with a difenoconazole-containing seed-treatment product such as Dividend (EPA Reg. No. 100-740).
*Based on an average seed weight of 850 peanuts/lb			
<b>Resistance Management:</b> Refer to <b>Section 3.1</b> .			
USE RESTRICTIONS			
<ul style="list-style-type: none"> <li>• <b>Maximum Annual Plantings:</b> 2</li> <li>• <b>DO NOT</b> exceed 0.073 lb ai/A/calendar year of azoxystrobin-containing products as a seed treatment.</li> <li>• <b>DO NOT</b> exceed 0.18 lb ai/A/calendar year of pydiflumetofen-containing products through any combination of seed and foliar applications.</li> <li>• <b>DO NOT</b> exceed 0.023 lb ai/A/calendar year of fludioxonil-containing products as a seed treatment.</li> <li>• <b>DO NOT</b> exceed 0.046 lb ai/A/year of sedaxane-containing products as a seed treatment.</li> <li>• <b>DO NOT</b> exceed 0.071 lb ai/A/calendar year of mefenoxam- or metalaxyl-containing products as a seed treatment.</li> <li>• <b>DO NOT</b> exceed the maximum single application rate of 2 fl oz product/100 lb seed.</li> <li>• The maximum allowed number of treated seeds is 193,800 treated seeds per acre (equivalent to 228 pounds of treated seed per acre).</li> <li>• <b>Sections 4.0, 4.1, 5.0, and 6.0</b> contain additional product use restrictions.</li> </ul>			

## 8.0 STORAGE AND DISPOSAL

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

#### Pesticide Storage

Keep this product in its tightly closed original container, when not in use. Store in a cool, dry (preferably locked) area that is inaccessible to children and animals.

#### Pesticide Disposal

Pesticide wastes may be hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative of the nearest EPA Regional Office for guidance.

#### Container Handling (less than or equal to 5 gallons)

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Add water – at least 2% of the container volume, and up to 1/3 of the volume of water needed to make the proper slurry composition with a maximum of 1/4 of the container volume – and recap. Shake for 30 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. If used in application equipment, adjust the slurry volume application rate to account for any added rinsate water. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

#### Container Handling (greater than 5 gallons)

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Add water – at least 2% of the container volume, and up to 1/3 of the volume of water needed to make the proper slurry composition with a maximum of 1/4 of the container volume. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 60 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. If used in application equipment, adjust the slurry volume application rate to account for any added rinsate water. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

#### Container Handling (greater than 5 gallons)

Refillable container. Refill this container with pesticide only. Do not reuse the container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Add water – at least 2% of the container volume, and up to 1/3 of the volume of water needed to make the proper slurry composition with a maximum of 1/4 of the container volume. Replace and tighten closure. Agitate vigorously or recirculate the rinsate with a pump for at least 2 minutes, ensuring that the rinsate rinses the walls of the container. Empty the rinsate into application equipment or rinsate collection system, for later use or disposal. Repeat this procedure two more times. If used in application equipment, adjust the slurry volume application rate to account for any added rinsate water. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

**CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER.**

## 9.0 CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, LLC or Seller. To the extent permitted by applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of the product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and (2) Buyer and User assume the risk of any such use. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.**

To the extent permitted by applicable law, in no event shall SYNGENTA be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

ADEPIDYN®, Dividend®, Trebuset®, the ALLIANCE FRAME,   
the SYNGENTA Logo, and the PURPOSE ICON   
are Trademarks of a Syngenta Group Company

©2022 Syngenta

For non-emergency (e.g., current product information), call  
Syngenta Crop Protection at 1-800-334-9481.

Manufactured for:  
Syngenta Crop Protection, LLC  
P.O. Box 18300  
Greensboro, North Carolina 27419-8300

**SCP 1689A-L1A 0122**  
**4153336**

MEFENOXAM	GROUP 4	FUNGICIDE
PYDIFLUMETOFEN	GROUP 7	FUNGICIDE
SEDAXANE	GROUP 7	FUNGICIDE
AZOXYSTROBIN	GROUP 11	FUNGICIDE
FLUDIOXONIL	GROUP 12	FUNGICIDE



## Fungicide

ADEPIDYN® Technology<sup>1</sup>

Active Ingredients:	
Azoxystrobin <sup>2</sup>	11.0%
Pydiflumetofen <sup>3</sup>	6.85%
Fludioxonil <sup>4</sup>	3.47%
Sedaxane <sup>5</sup>	3.34%
Mefenoxam <sup>6</sup>	0.64%
Other Ingredients:	74.7%
Total:	100.00%

<sup>1</sup>Technology denotes the active ingredient Pydiflumetofen.

<sup>2</sup>CAS No. 131860-33-8

<sup>3</sup>CAS No. 1228284-64-7

<sup>4</sup>CAS No. 131341-86-1

<sup>5</sup>CAS No. 874967-67-6

<sup>6</sup>CAS No. 70630-17-0 and 69516-34-3

Trebuset Peanuts Colorless is a flowable concentrate for seed treatment containing 1.03 lb azoxystrobin, 0.64 lb pydiflumetofen, 0.32 lb fludioxonil, 0.31 lb sedaxane, and 0.060 lb mefenoxam per gallon.

See additional precautionary statements and directions for use inside booklet.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

EPA Reg. No. 100-1689 EPA Est. 100-NE-001

ADEPIDYN®, Trebuset®, and the SYNGENTA Logo are Trademarks of a Syngenta Group Company

©2022 Syngenta

Manufactured by:  
Syngenta Crop Protection, LLC  
P.O. Box 18300  
Greensboro, North Carolina 27419-8300

SCP 1689A-L1A 0122  
4153336

**15 gallons**  
Net Contents

## KEEP OUT OF REACH OF CHILDREN. CAUTION

### FIRST AID

**If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person. **If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

**HOTLINE NUMBER:** For 24-Hour Medical Emergency Assistance (Human or Animal) Or Chemical Emergency Assistance (Spill, Leak, Fire or Accident) Call **1-800-888-8372**.

### PRECAUTIONARY STATEMENTS

#### Hazards to Humans and Domestic Animals

#### CAUTION

Harmful if swallowed or absorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

#### Environmental Hazards

This product is toxic to fish, aquatic invertebrates, oysters, and shrimp. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. **DO NOT** apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwater or rinsate.

#### Groundwater Advisory

Azoxystrobin, a degradate of azoxystrobin, and mefenoxam are known to leach through soil into groundwater under certain conditions as a result of label use. Fludioxonil and pydiflumetofen have properties and characteristics associated with chemicals detected in groundwater. These chemicals may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

#### Surface Water Advisory

This product may impact surface water quality due to runoff of rainwater. This is especially true for poorly draining soils and soils with shallow groundwater. This product is classified as having a medium potential for reaching both surface water and aquatic sediment via runoff several months or more after application.

A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of pydiflumetofen from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours. Sound erosion control practices will reduce this product's potential to reach aquatic sediment via runoff.

#### Physical or Chemical Hazards

**DO NOT** mix or allow to come in contact with oxidizing agents. Hazardous chemical reaction may occur.

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Keep this product in its tightly closed original container, when not in use. Store in a cool, dry (preferably locked) area that is inaccessible to children and animals.

**Pesticide Disposal:** Pesticide wastes may be hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative of the nearest EPA Regional Office for guidance.

#### Container Handling: Non-refillable container.

Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Add water – at least 2% of the container volume, and up to 1/3 of the volume of water needed to make the proper slurry composition with a maximum of 1/4 of the container volume. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 60 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. If used in application equipment, adjust the slurry volume application rate to account for any added rinsate water. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

**CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER.**

**syngenta®**

