



# Trimec<sup>®</sup> CRABGRASS<sup>®</sup>

*Plus*  
**LAWN WEED KILLER**  
READY - T O - U S E

**ACTIVE INGREDIENTS:**

2,4-D, dimethylamine salt	0.313%
Quinclorac	0.104%
Dicamba, dimethylamine salt	0.029%
<b>OTHER INGREDIENTS:</b>	<b>99.554%</b>
<b>TOTAL 100.000%</b>	

**THIS PRODUCT CONTAINS:**

2,4-dichlorophenoxyacetic acid equivalent, 0.260%  
 3,7-dichloro-8-quinolinecarboxylic acid, 0.104%  
 3,6-dichloro-o-anisic acid equivalent, 0.024%  
 Isomer Specific by AOAC Method.

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**



**READ THE ENTIRE LABEL FIRST.  
OBSERVE ALL PRECAUTIONS AND  
FOLLOW DIRECTIONS CAREFULLY.**

**PRECAUTIONARY STATEMENTS**

**Hazards to Humans and Domestic Animals**

**CAUTION:** Causes moderate eye irritation. Harmful if swallowed. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash clothing before reuse. Wear long sleeved shirt, long pants, shoes and socks.

<b>First Aid:</b>	
<b>If in eyes:</b>	<ul style="list-style-type: none"> <li>Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first five minutes, then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If swallowed:</b>	<ul style="list-style-type: none"> <li>Call a poison control center or doctor immediately for treatment advice.</li> <li>Have person sip a glass of water if able to swallow.</li> <li>Do not induce vomiting unless told to by a poison control center or doctor.</li> <li>Do not give anything to an unconscious person.</li> </ul>
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment advice.	

**ENVIRONMENTAL HAZARDS:**

This pesticide is toxic to fish and aquatic invertebrates. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area.

This product has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

**DIRECTIONS FOR USE:**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**USE RESTRICTIONS:**

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until sprays have dried.
- Apply when daytime temperatures are between 50° and 90°F (do not apply above 85°F for bermudagrass). Do not treat when air temperatures exceed 90°F as damage to turfgrass may occur.
- Do not apply this product to food crops (gardens, fruits and vegetables), forage crops, or ornamental plants including flowers, trees, shrubs, hedges, woody ornamentals, groundcovers established in landscape plantings and other nontarget plants.
- Do not spray exposed roots of ornamentals and trees.
- This product is for spot treatment only.
- Do not use clippings from treated area as compost or mulch in vegetable gardens, flower beds, or around other desirable plants.

**PRODUCT FACTS:**

**What It Does:**

Kills major broadleaf and troublesome grass weeds including dandelion, chickweed, clover as well as crabgrass and foxtail. (See below for complete list.)

**Where to Use:**

Lawns: Kills the weeds not the lawn when used as directed.

**When to Use:**

Apply when daytime temperatures are between 50° and 90°F, (do not apply above 85°F for bermudagrass).

**Questions & Comments:**

Call 1-800-884-3179 or visit our website at [www.GordonsUSA.com](http://www.GordonsUSA.com)

**SHAKE WELL BEFORE USING.**

**HOW IT WORKS:**

Trimec<sup>®</sup> Crabgrass Plus Lawn Weed Killer Ready-To-Use contains three proven weed killers that target lawn weeds and crabgrass. The herbicides enter the lawn weeds through their leaves and move inside the weeds, even down into the roots. And lawns listed on this label will not be harmed when used as directed.

**WHERE TO USE:**

Trimec<sup>®</sup> Crabgrass Plus Lawn Weed Killer Ready-To-Use is a ready-to-use (RTU) product intended for the treatment of individual weeds, patches of weeds or sparse weeds scattered throughout home lawns. Trimec<sup>®</sup> Crabgrass Plus Lawn Weed Killer Ready-To-Use can be applied to the following established turfgrass species:

Cool season turfgrass	Warm season turfgrass
Kentucky bluegrass	Bermudagrass, common and hybrid
Ryegrass, perennial and annual	Zoysiagrass
Tall fescue	Buffalograss
Mixtures of cool season grasses containing fine fescues (red, hard and chewings) and bentgrass	

DO NOT apply Trimec<sup>®</sup> Crabgrass Plus Lawn Weed Killer Ready-To-Use to the following:

Warm season turfgrass	Other areas
St. Augustinegrass	Lawns with desirable clovers or legumes
Bahiagrass	Dichondra
Centipedegrass	Ornamental plants (flowers, trees, groundcovers, landscape beds and shrubs)
Carpetgrass	Turfgrass species that are not listed on this label.
Seashore paspalum	Gardens including vegetables, fruit trees, vines and berries.

**WEEDS CONTROLLED**

**GRASS WEEDS:**

- crabgrass (large and smooth)
- foxtail (green, yellow and giant)
- kikuyugrass and
- broadleaf signalgrass

**BROADLEAF WEEDS:**

Annual yellow sweetclover	Fiddleneck	Rough cinquefoil
Aster	Florida pusley	Rough fleabane
Austrian fieldcress	Frenchweed	Russian pigweed
Bedstraw	Galinsoga	Russian thistle
Beggarticks	Goathead	Scarlet pimpernel
Betony, Florida	Goldenrod	Scotch thistle
Bindweed, field	Ground ivy	Sheep sorrel
Bird vetch	Gumweed	Shepherd's purse
Bitter wintercress	Hairy fleabane	Slender plantain
Bittercress, hairy	Hawkweed	Smallflower galinsoga
Bitterweed	Healall	Smooth dock
Black-eyed Susan	Heartleaf drymary	Smooth clover
Black medic	Heathaster	Sorrel
Black mustard	Hedge bindweed	Sowthistle
Blackseed plantain	Hedge mustard	Spanish needles
Blessed thistle	Hemp	Speedwell
Bloodflower milkweed	Henbit	Spiny amaranth
Blue lettuce	Hoary cress	Spiny cocklebur
Blue vervain	Hoary plantain	Spiny sowthistle
Bracted plantain	Hoary vervain	Spotted catsear
Brassbuttons	Horsenettle	Spotted spurge
Bristly oxtongue	Jimsonweed	Spurweed
Broadleaf dock	Knawel	Stinging nettle
Broadleaf plantain	Knotweed	Strawberry, India mock
Broomweed	Kochia	Tall nettle
Buckhorn	Lambsquarters	Tall vervain
Buckhorn plantain	Lespedeza	Tansy ragwort
Bulbous buttercup	Mallow	Tansy mustard
Bull thistle	Matchweed	Tanweed
Bullnettle	Mexicanweed	Thistle
Burclover	Milk vetch	Trailing crownvetch
Burdock	Morningglory	Tumble mustard
Burning nettle	Mousser hawkweed	Tumble pigweed
Burweed	Mugwort	Velvetleaf
Buttercup	Musk thistle	Venice mallow
Buttonweed	Mustard	Virginia buttonweed
Canada thistle	Narrowleaf plantain	Virginia creeper
Carolina geranium	Narrowleaf vetch	Virginia pepperweed
Carpetweed	Nettle	Wavyleaf bullthistle
Catchweed bedstraw	Orange hawkweed	Western clematis
Catnip	Oriental cocklebur	Western salsify
Catsear	Oxalis	White mustard
Chickweed, common	Parsley-piert	Wild aster
Chickweed, mouseear	Parsnip	Wild buckwheat
Chicory	Pearlwort	Wild carrot
Cinquefoil	Pennycress	Wild four-o'clock
Clover, crimson	Pennywort	Wild garlic
Clover, hop	Peppergrass	Wild geranium
Clover, red	Pepperweed	Wild lettuce
Clover, strawberry	Pigweed	Wild marigold
Clover, sweet	Pineywoods bedstraw	Wild mustard
Clover, white	Plains coreopsis (tickseed)	Wild onion
Cockle	Plantain	Wild parsnip
Cocklebur	Poison ivy	Wild radish
Common mullein	Poison oak	Wild rape
Creeping Jenny (Creeping Charlie)	Pokeweed	Wild strawberry
Cudweed	Poorjoe	Wild sweet potato
Curly dock	Prairie sunflower	Wild vetch
Daisy, English	Prickly lettuce	Wild violet
Daisy fleabane	Prickly sida	Woodsorrel
Daisy, oxeye	Prostrate knotweed	Woolly croton
Dandelion	Prostrate pigweed	Woolly morningglory
Dichondra	Prostrate spurge	Woolly plantain
Dogbane	Prostrate vervain	Wormseed
Dogfennel	Puncturevine	Yarrow
Dollarweed	Purslane, common	Yellow rocket
False dandelion	Ragweed	Yellowflower pepperweed
Falseflax	Red sorrel	
False sunflower	Redroot pigweed	
	Redstem filaree	

**WHEN TO USE:**

Apply this product when weeds are small and actively growing.

**CRABGRASS (large and smooth):**

- **Spring:** Apply this product from crabgrass emergence until the crabgrass is 3-4 inches in height. This period begins the first time that you can see crabgrass in your lawn. Re-treat at 21 days later for dense populations.

**FOXTAIL (GREEN, GIANT AND YELLOW):**

- **Spring and early summer:** Apply this product to foxtail during the early growth stages or when foxtail is 1-4 inches in height. Re-treat at 21 days later for dense populations.

**BROADLEAF WEEDS:**

- **Spring:** Apply when **dandelion, plantain, buttercup, speedwell, knotweed** and other broadleaf weeds are young and actively growing with adequate soil moisture for best results.
- **Fall:** Apply when **henbit, chickweed, white clover, ground ivy, lawn burweed** and other broadleaf weeds are young and actively growing with adequate soil moisture for best results.

**HOW TO USE:**

**TRIGGER SPRAYERS:**

1. Shake well.
2. Adjust the nozzle to coarse spray pattern to minimize wind drift. Spray the weeds only. Spray to lightly cover broadleaf and grass weeds. Excessive spraying can injure some lawns. Do not apply when daytime temperatures are above 90°F as lawn injury can occur (do not apply above 85°F for bermudagrass).

**USE PRECAUTIONS:**

**For newly seeded areas:**

- Do not apply within 4 weeks after seedling emergence of Kentucky bluegrass, fine fescue mixtures and perennial ryegrass. Apply this product to newly seeded grasses when well-established (approximately 6 weeks after seedling emergence) or after the third mowing.

**For newly sodded, sprigged, or plugged areas:**

- The application of this product to newly sodded, sprigged, or plugged grasses should be delayed until 3 to 4 weeks after the sodding, sprigging, or plugging operations.

**Seeding:**

- Turf species listed on this label can be seeded into the treated areas at four (4) weeks after the application of this product.

**Irrigation:**

- Do not apply this product immediately before rainfall or irrigation. For best results, do not irrigate or water the turfgrass within 24 hours after application.
- Avoid applications of this product when turfgrasses are under stress since injury may result.

**Mowing:**

- Delay mowing 2 days before and until 2 days after the application of this product.

**STORAGE AND DISPOSAL**

**PESTICIDE STORAGE:** Keep from freezing. Store in original container in a locked storage area inaccessible to children and pets.

**PESTICIDE DISPOSAL AND CONTAINER HANDLING:** Non-refillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

**LIMITED WARRANTY AND DISCLAIMER**

FOR USE ONLY AS DIRECTED. THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

Trimec® and Gordon® logo are registered trademarks of PBI-Gordon Corporation.

755/1-2019 AP110216  
EPA REG. NO. 2217-906



Employee-Owned

MANUFACTURED BY  
PBI/GORDON CORPORATION  
P.O. BOX 860350  
SHAWNEE, KANSAS 66286  
www.GordonsUSA.com

ATTENTION: This specimen label is provided for informational use only. This product may not yet be available for sale in your state or area. The information found in this label may differ from the information found on the product label you are using. Always follow the instructions for use and precautions on the label of the product you are using.