



Trimec[®] CRABGRASS[®]

PLUS LAWN WEED KILLER

CONCENTRATE

ACTIVE INGREDIENTS:

2,4-D, dimethylamine salt	6.42%
Quinclorac	2.13%
Dicamba, dimethylamine salt	0.60%
OTHER INGREDIENTS:	90.85%
	TOTAL 100.00%

THIS PRODUCT CONTAINS:

0.456 lb 2,4-dichlorophenoxyacetic acid equivalent per gallon or 5.33%
 0.182 lb 3,7-dichloro-8-quinolinecarboxylic acid per gallon or 2.13%
 0.043 lb 3,6-dichloro-o-anisic acid equivalent per gallon or 0.50%
 Isomer Specific By AOAC Methods.

KEEP OUT OF REACH OF CHILDREN CAUTION



**READ THE ENTIRE LABEL FIRST.
OBSERVE ALL PRECAUTIONS AND
FOLLOW DIRECTIONS CAREFULLY.**

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Harmful if swallowed. When using this product, wear long-sleeved shirt, long pants, socks, shoes, and rubber gloves.

Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower.

User Safety Recommendations

- Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to by a poison control center or doctor. • Do not give anything to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment advice.

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

1. Use Restrictions

- For residential lawn use only.
- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until sprays have dried.
- Use on established lawns. Do not use on Bahiagrass, Bentgrass (colonial and creeping), Carpetgrass, Centipedegrass, Dichondra, Gardens (including vegetables, fruit trees, vines and berries), Ornamental plants (flowers, trees, groundcovers, landscape beds and shrubs), Seashore Paspalum, St. Augustinegrass, and turfgrass species that are not listed on this label.)
- Application to bermudagrass may cause temporary yellowing or discoloration but full recovery can be expected. Do not apply to bermudagrass when daytime temperatures exceed 85°F.
- Do not use clippings from the treated areas as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens.
- Do not apply this product to food crops (gardens, fruits, and vegetables), forage crops, or ornamental plants including flowers, trees, shrubs, hedges, woody ornamentals, groundcovers established in landscape planting and other nontarget plants.
- Do not spray exposed roots of ornamentals and trees.

Spray Drift Management

It is the responsibility of the applicator to avoid spray drift at the application site. Do not allow the herbicide solution to mist, drip, drift, or splash onto desirable broadleaf plants, as small amounts of this product can damage sensitive plants near the treated area. If desirable plants are accidentally sprayed, immediately rinsing leaves with water may reduce or eliminate plant damage.

2. How Much To Apply

For Bermudagrass (common and hybrid), Buffalograss, Kentucky Bluegrass, mixtures of cool season grasses containing fine fescues (red, hard and chewings), Ryegrass (perennial and annual), Tall Fescue, and Zoysiagrass

Spot Treatments: Mix 2.5 fl.oz. (5 Tbsp) in 1 gallon of water to treat 400 sq.ft.

Entire Lawn Treatments:

- Pump Style Sprayer: Mix 6.4 fl.oz. in 0.5 to 1 gallon of water to treat 1,000 sq.ft.
- Hose-End Sprayer: Apply 6.4 fl.oz. per 1,000 sq.ft.
- 12 Volt & Tow-Behind Sprayers: Mix 6.4 fl.oz. in 0.5 to 1 gallon of water to treat 1,000 sq.ft.

Over-application or rates above those specified on this label can cause turf injury.

3. When To Use

Apply this product when weeds are small and actively growing.

Weed Species	Timing	Comments
Crabgrass (large and smooth):	Apply in spring from crabgrass emergence until the crabgrass is 3 to 4 inches in height.	Re-treat at the same rate at 21 days later for dense populations or for continued crabgrass emergence.
Foxtail (green, giant and yellow):	Apply in spring and early summer during the early growth stages or when foxtail is 1 to 4 inches in height.	Re-treat at the same rate at 21 days later for dense populations.

(continued on next page)

Weed Species (cont.)	Timing (cont.)	Comments (cont.)
Broadleaf signalgrass	Apply in spring and early summer during the early growth stages or during the 3 to 7 leaf stage.	Re-treat at the same rate at 21 days later if needed.
Dandelion, plantain, buttercup, speedwell, knotweed and other listed broadleaf weeds	Apply in spring when broadleaf weeds are small and actively growing.	Apply when broadleaf weeds are young and actively growing with adequate soil moisture.
Henbit, chickweed, white clover, ground ivy, lawn burweed and other listed broadleaf weeds	Apply in fall when broadleaf weeds are young and actively growing.	Apply when broadleaf weeds are young and actively growing with adequate soil moisture.
The maximum application rate is 6.4 fl.oz. of product per 1,000 sq.ft. per application. The maximum number of applications is limited to 2 per year with a minimum of 21 days between applications. The maximum seasonal rate is 12.8 fl.oz. of product per 1,000 sq.ft.		

4. How To Apply

4.1 Pump Style Sprayers

Ideal for spraying individual weeds.

1. Add the appropriate amount of water and concentrate to the sprayer tank.
2. Close sprayer, shake well, and pump handle to pressurize.
3. Adjust nozzle to deliver a coarse spray pattern.
4. To broadcast over large areas: apply evenly to treated area.
5. To treat individual weeds: spray to wet weed leaf surfaces.
6. Re-pressurize the sprayer as needed to maintain a good spray pattern.

4.2 12 Volt & Tow-Behind Boom Sprayers

Ideal for treating large country and suburban lawns.

1. To calibrate your application equipment consult your equipment owner's manual for speed and pressure settings required to deliver 0.5 to 1 gallon of spray solution to treat 1,000 sq.ft. For example, at 4 mph a sprayer might deliver 1/2 gallon per 1,000 sq.ft. at 40 psi.
2. To apply evenly, drive at a constant speed that delivers the prescribed amount of spray required for the area to be treated.
3. Mix the appropriate amount of product and water in the sprayer using the following tables as a guide.

Tank Size	Spray Volume	Coverage Area	Concentrate To Add To Tank
15 gal	0.5 gal/1000 sq.ft.	30,000 sq.ft.	192 fl.oz. (1.5 gal)
	1 gal/1000 sq.ft.	15,000 sq.ft.	96 fl.oz. (0.75 gal)
25 gal	0.5 gal/1000 sq.ft.	50,000 sq.ft.	320 fl.oz. (2.5 gal)
	1 gal/1000 sq.ft.	25,000 sq.ft.	160 fl.oz. (1.25 gal)

For spot treatments with 12 Volt & Pull Behind Sprayers

See rates under "How Much To Apply," Spot Treatments.

4.3 Dial Style Hose-End Sprayers

Ideal for treating medium-sized lawns.

1. Fill sprayer with sufficient concentrate to cover the total area to be treated. **DO NOT ADD WATER.**
2. Close sprayer and set dial to 2.5 oz.
3. Connect sprayer to hose.
4. Extend hose to furthest point from faucet. Walking back towards the faucet, begin spraying, avoiding contact with the treated area.
5. Pour any unused product back into this container.

4.4 Concentrate Plus Water Hose-End Sprayers

Ideal for treating medium-sized lawns.

1. Fill sprayer with sufficient concentrate to cover the total area to be treated. Pour product into sprayer jar to the "Fluid Oz." level desired (See sprayer jar for markings)
2. Add the appropriate amount of water to the "Gallons" level indicated on the side of the sprayer jar.
3. Close sprayer, connect to garden hose, shake well.
4. Extend hose to furthest point from faucet. Walking back towards the faucet, begin spraying, avoiding contact with the treated area.

5. Use Precautions

- Apply when weeds are young and actively growing.
- Delay mowing 2 days before and until 2 days after the application of this product.

- If the lawn is dry, water 24 hours before using this product.
- Do not water lawn within 24 hours of treatment.
- For new lawns, apply only when grass is well-established (approximately 6 weeks after seedling emergence) or after the third mowing.
- Bare spots may be seeded 3 to 4 weeks after application.
- Spray when air temperatures are cooler than 90°F.
- Avoid applications during the spring transition of the warm season grasses and during periods of extremely high temperatures.
- Application to bermudagrass may cause temporary yellowing or discoloration but full recovery can be expected. Do not apply to bermudagrass when daytime temperatures exceed 85°F.

6. Grass and Broadleaf Weeds Killed

GRASS WEEDS		
Crabgrass, large and smooth	Foxtail, green, yellow and giant	Signalgrass, broadleaf

BROADLEAF WEEDS		
Annual yellow sweetclover	False sunflower	Redroot pigweed
Aster	Fiddleneck	Redstem filaree
Austrian fieldcress	Florida pusley	Rough cinquefoil
Bedstraw	Frenchweed	Rough fleabane
Beggarticks	Galinsoga	Russian pigweed
Betony, Florida	Goathead	Russian thistle
Bindweed, field	Goldenrod	Scarlet pimpernel
Bird vetch	Ground ivy	Scotch thistle
Bitter wintercress	Gumweed	Sheep sorrel
Bittercress, hairy	Hairy fleabane	Shepherd's purse
Bitterweed	Hawkweed	Slender plantain
Black-eyed Susan	Healall	Smallflower galinsoga
Black medic	Heartleaf drymary	Smooth dock
Black mustard	Heathaster	Smooth pigweed
Blackseed plantain	Hedge bindweed	Sorrel
Blessed thistle	Hedge mustard	Sowthistle
Bloodflower milkweed	Hemp	Spanish needles
Blue lettuce	Henbit	Speedwell
Blue vervain	Hoary cress	Spiny amaranth
Bracted plantain	Hoary plantain	Spiny cocklebur
Brassbuttons	Hoary vervain	Spiny sowthistle
Bristly oxtongue	Horsenettle	Spotted catsear
Broadleaf dock	Jimsonweed	Spotted spurge
Broadleaf plantain	Knawel	Spurweed
Broomweed	Knotweed	Stinging nettle
Buckhorn	Kochia	Strawberry, India mock
Buckhorn plantain	Lambsquarters	Tall nettle
Bulbous buttercup	Lespedeza	Tall vervain
Bull thistle	Mallow	Tansy ragwort
Bullnettle	Matchweed	Tansy mustard
Burclover	Mexicanweed	Tanweed
Burdock	Milk vetch	Thistle
Burning nettle	Morningglory	Trailing crownvetch
Burweed	Mouseear hawkweed	Tumble mustard
Buttercup	Mugwort	Tumble pigweed
Buttonweed	Musk thistle	Velvetleaf
Canada thistle	Mustard	Venice mallow
Carolina geranium	Narrowleaf plantain	Virginia buttonweed
Carpentweed	Narrowleaf vetch	Virginia creeper
Catchweed bedstraw	Nettle	Virginia pepperweed
Catnip	Orange hawkweed	Wavyleaf bullthistle
Catsear	Oriental cocklebur	Western clematis
Chickweed, common	Oxalis	Western salsify
Chickweed, mouseear	Parsley-piert	White mustard
Chicory	Parsnip	Wild aster
Cinquefoil	Pearlwort	Wild buckwheat
Clover, crimson	Pennycress	Wild carrot
Clover, hop	Pennywort	Wild four-o'clock
Clover, red	Peppergrass	Wild garlic
Clover, strawberry	Pepperweed	Wild geranium
Clover, sweet	Pigweed	Wild lettuce
Clover, white	Pineywoods bedstraw	Wild marigold
Cockle	Plains coreopsis (tickseed)	Wild mustard
Cocklebur	Plantain	Wild onion
Common mullein	Poison ivy	Wild parsnip
Creeping Jenny (Creeping Charlie)	Poison oak	Wild radish
Cudweed	Pokeweed	Wild rape
Curly dock	Poorjoe	Wild strawberry
Daisy, English	Prairie sunflower	Wild sweet potato
Daisy fleabane	Prickly lettuce	Wild vetch
Daisy, oxeye	Prickly sida	Wild violet
Dandelion	Prostrate knotweed	Woodsorrel
Dichondra	Prostrate pigweed	Woolly croton
Dogbane	Prostrate spurge	Woolly morningglory
Dogfennel	Prostrate vervain	Woolly plantain
Dollarweed	Puncturevine	Wormseed
False dandelion	Purslane, common	Yarrow
Falseflax	Ragweed	Yellow rocket
	Red sorrel	Yellowflower pepperweed

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Keep from freezing. Store in original container in a locked storage area inaccessible to children and pets.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

GORDON'S® logo and TRIMEC® are registered trademarks of PBI-Gordon Corporation.

761/10-2022 AP040616
EPA REG. NO. 2217-896



Employee-Owned

MANUFACTURED BY
PBI/GORDON CORPORATION
P.O. BOX 860350
SHAWNEE, KANSAS 66286
www.GordonsUSA.com

ATTENTION: This specimen label is provided for informational use only. This product may not yet be available for sale in your state or area. The information found in this label may differ from the information found on the product label you are using. Always follow the instructions for use and precautions on the label of the product you are using.