



Trimec

LAWN WEED KILLER

ACTIVE INGREDIENTS:

| | |
|--------------------------------------|---------------|
| 2,4-D, dimethylamine salt | 7.59% |
| Mecoprop-p, dimethylamine salt | 1.83% |
| Dicamba, dimethylamine salt | 0.84% |
| OTHER INGREDIENTS: | 89.74% |
| TOTAL 100.00% | |

THIS PRODUCT CONTAINS:

0.54 lb 2,4-dichlorophenoxyacetic acid equivalent per gallon or 6.30%
 0.13 lb (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid equivalent per gallon or 1.51%
 0.06 lb 3,6-dichloro-o-anisic acid equivalent per gallon or 0.70%

Isomer Specific by AOAC Method.

2,4-D, dimethylamine salt (CAS Registry No. 2008-39-1)
 Mecoprop-p, dimethylamine salt (CAS Registry No. 66423-09-4)
 Dicamba, dimethylamine salt (CAS Registry No. 2300-66-5)

KEEP OUT OF REACH OF CHILDREN CAUTION



**READ THE ENTIRE LABEL FIRST.
OBSERVE ALL PRECAUTIONS AND
FOLLOW DIRECTIONS CAREFULLY.**

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if absorbed through skin. Harmful if swallowed. Avoid contact with skin, eyes, or clothing. Causes moderate eye irritation. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Wear long-sleeved shirt, long pants, socks, shoes and gloves. Remove and wash contaminated clothing before reuse.

| First Aid | |
|--|--|
| If in eyes: | <ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. |
| If swallowed: | <ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person. |
| If on skin or on clothing: | <ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a poison control center or doctor for treatment advice. |
| If inhaled: | <ul style="list-style-type: none"> Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for treatment advice. |
| HOT LINE NUMBER: Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment information. | |

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

1. Use Restrictions

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until sprays have dried.
- Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.
- Use on established lawns. Will not harm lawn grasses when used as directed (Do not use on carpetgrass, improved varieties of St. Augustinegrass, such as 'Floritam,' dichondra or desirable clovers)
- The turf tolerance of centipedegrass and St. Augustinegrass can range from sensitive to tolerant. Do not apply this product to 'Floritam' St. Augustinegrass in Florida.
- Do not apply as a fine mist because of potential for injury to desirable plants.
- Do not use this product on or near desirable plants, including contact of spray on exposed root systems or adventitious shoots within the drip line of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day. Rinse sprayer over lawn area only.

2. How Much To Apply

Bahiagrass, Bermudagrass, Fescues, Kentucky Bluegrass, Perennial Ryegrass & Zoysiagrass:

Spot Treatments: Mix 1 fl.oz. (2 Tbsp) in 1 gallon of water to treat 250 sq.ft.

Entire Lawn Treatments:

- Pump Style Sprayer: Mix 4 fl.oz. in 0.5 to 1 gallon of water to treat 1,000 sq.ft.
- Hose-End Sprayer: Apply 4 fl.oz. per 1,000 sq.ft.
- 12 Volt & Tow-Behind Sprayers: Mix 4 fl.oz. in 0.5 to 1 gallon of water to treat 1,000 sq.ft.

Table 1: Spot Treatment of Bahiagrass, Bermudagrass, Fescues, Kentucky Bluegrass, Perennial Ryegrass, & Zoysiagrass

| Tank Size | Concentrate to add to tank | Coverage Area |
|-----------|----------------------------|---------------|
| 1 gal | 1 fl.oz. | 250 sq.ft. |

Table 2: Entire Lawn Treatment of Bahiagrass, Bermudagrass, Fescues, Kentucky Bluegrass, Perennial Ryegrass, & Zoysiagrass

| Tank Size | Concentrate to add to tank | Coverage Area |
|-----------|----------------------------|---------------|
| 1 gal | 4 fl.oz. | 1,000 sq.ft. |
| 15 gal | 60 fl.oz. | 15,000 sq.ft. |
| 25 gal | 100 fl.oz. | 25,000 sq.ft. |

Centipedegrass & St. Augustinegrass:

Spot treatments: Mix 0.5 fl.oz. (1 Tbsp) in 1 gallon of water to treat 250 sq.ft.

Entire Lawn Treatments:

- Pump Style Sprayer: Mix 2 fl.oz. in 0.5 to 1 gallon of water to treat 1,000 sq.ft.
- Hose-End Sprayer: Apply 2 fl.oz. per 1,000 sq.ft.
- 12 Volt & Tow-Behind Sprayers: Mix 2 fl.oz. in 0.5 to 1 gallon of water to treat 1,000 sq.ft.

| Tank Size | Concentrate to add to tank | Coverage Area |
|-----------|----------------------------|---------------|
| 1 gal | 1 fl.oz. | 500 sq.ft. |

| Tank Size | Concentrate to add to tank | Coverage Area |
|-----------|----------------------------|---------------|
| 1 gal | 4 fl.oz. | 2,000 sq.ft. |
| 15 gal | 60 fl.oz. | 30,000 sq.ft. |
| 25 gal | 100 fl.oz. | 50,000 sq.ft. |

3. How To Apply

3.1 Pump Style Sprayers

Ideal for spraying individual weeds

1. Add the appropriate amount of water and concentrate to the sprayer tank.
2. Close sprayer, shake well, and pump handle to pressurize.
3. Adjust nozzle to deliver a coarse spray pattern.
4. To broadcast over large areas: apply evenly to treated area.
5. To treat individual weeds: spray to wet weed leaf surfaces.
6. Re-pressurize the sprayer as needed to maintain a good spray pattern.

3.2 12 Volt & Tow-Behind Boom Sprayers

Ideal for treating large country and suburban lawns

1. To calibrate your application equipment consult your equipment owner’s manual for speed and pressure settings required to deliver 0.5 to 1 gallon of spray solution to treat 1,000 sq.ft.. For example, at 4 mph a sprayer might deliver 1/2 gallon per 1,000 sq.ft. at 40 psi.
2. To apply evenly, drive at a constant speed that delivers the prescribed amount of spray required for the area to be treated.
3. Mix the appropriate amount of product and water in the sprayer using the following tables as a guide.

| Tank Size | Spray volume | Coverage Area | Concentrate to add to tank |
|-----------|---------------------|---------------|----------------------------|
| 15 gal | 0.5 gal/1000 sq.ft. | 30,000 sq.ft. | 120 fl.oz. |
| | 1 gal/1000 sq.ft. | 15,000 sq.ft. | 60 fl.oz. |
| 25 gal | 0.5 gal/1000 sq.ft. | 50,000 sq.ft. | 200 fl.oz. |
| | 1 gal/1000 sq.ft. | 25,000 sq.ft. | 100 fl.oz. |

| Tank Size | Spray volume | Coverage Area | Concentrate to add to tank |
|-----------|---------------------|---------------|----------------------------|
| 15 gal | 0.5 gal/1000 sq.ft. | 30,000 sq.ft. | 60 fl.oz. |
| | 1 gal/1000 sq.ft. | 15,000 sq.ft. | 30 fl.oz. |
| 25 gal | 0.5 gal/1000 sq.ft. | 50,000 sq.ft. | 100 fl.oz. |
| | 1 gal/1000 sq.ft. | 25,000 sq.ft. | 50 fl.oz. |

For spot treatments with 12 Volt & Pull Behind Sprayers

See rates under How Much To Apply.

3.3 Dial Style Hose-End Sprayers

Ideal for treating medium-sized lawns

1. Fill sprayer with sufficient concentrate to cover the total area to be treated. **DO NOT ADD WATER.**
2. Close sprayer and set dial to 1 oz.

3. Connect sprayer to hose.
4. Extend hose to furthest point from faucet. Walking back towards the faucet, begin spraying, avoiding contact with the treated area.
5. Pour any unused product back into this container.

3.4 Concentrate Plus Water Hose-End Sprayers

Ideal for treating medium-sized lawns

1. Fill sprayer with sufficient concentrate to cover the total area to be treated. Pour product into sprayer jar to the “Fluid Oz.” level desired (See sprayer jar for markings).
2. Add the appropriate amount of water to the “Gallons” level indicated on the side of the sprayer jar. Close sprayer, connect to garden hose, shake well.
3. Extend hose to furthest point from faucet. Walking back towards the faucet, begin spraying, avoiding contact with the treated area.

4. Use Precautions

- Apply when weeds are young and actively growing.
- Apply 1 to 3 days after mowing.
- If the lawn is dry, water 24 hours before using this product.
- Do not water lawn within 24 hours after treatment.
- Certain hard-to-kill weeds may need to be spot treated in 30 days.
- For new lawns, apply only when grass has reached a height of at least 2 inches.
- Bare spots may be seeded 3 to 4 weeks after application.
- Spray when air temperatures are between 60° and 90°F with good soil moisture.
- Avoid applications during the spring transition of the warm season grasses and during periods of extremely high temperatures.

5. Limitations

Broadcast applications to residential turfgrass:

The maximum application rate is 4.0 fl.oz. of product per 1,000 sq.ft. per application. The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 8.0 fl.oz. of product per 1,000 sq.ft.

Spot treatments to residential turfgrass:

Spot treatment is defined as a treatment area no greater than 1,000 sq.ft. per acre. The maximum application rate is 4.0 fl.oz. per 1,000 sq.ft. per application. The maximum number of spot treatments is limited to 2 per year with a minimum of 30 days between applications.

| | | |
|------------------------------|------------------------------------|--------------------------|
| Amaranth, purple | Catnip | Falseflax |
| Amaranth, slender | Catsear, spotted (false dandelion) | False sunflower |
| Aster, heath | Chickweed, common | Fiddleneck |
| Aster, white heath | Chickweed, mouseear | Filaree, redstem |
| Aster, purple | Chickweed, sticky | Filaree, whitestem |
| Aster, white prairie | Chicory | Fleabane, daisy (annual) |
| Austrian fieldcress | Cinquefoil | Fleabane, rough |
| Bedstraw, Catchweed | Clover, crimson | Galinsoga, hairy |
| Bedstraw, smooth | Clover, alyce | Galinsoga, smallflower |
| Bedstraw, Northern | Clover, hop | Goldenrod |
| Bedstraw, pinewoods | Clover, rabbit foot | Ground ivy |
| Beggarticks, hairy | Clover, red | Gumweed |
| Betony, Florida | Clover, strawberry | Hairy fleabane |
| Bindweed, Field | Clover, yellow sweet | Hawkweed, meadow |
| Bindweed, Hedge | Clover, white sweet | Hawkweed, mouseear |
| Bitter Wintercress | Clover, white | Hawkweed, orange |
| Bittercress, hairy | Cockle | Hawkweed, yellow |
| Bitterweed | Cocklebur | Healall |
| Black-eyed Susan | Cocklebur, oriental | Heartleaf drymary |
| Black Medic | Cocklebur, spiny | Hemp |
| Blood flower milkweed | Common mullein | Henbit |
| Blue lettuce | Creeping Jenny (Creeping Charlie) | Hoary cress |
| Brassbuttons | Cudweed | Horsenettle |
| Bristly oxtongue | Daisy, English | Jimsonweed |
| Broomweed | Daisy, oxeye | Knawel |
| Buckhorn | Dandelion, common | Knotweed, prostrate |
| Bullnettle | Dandelion, false | Kochia |
| Burclover | Deadnettle | Lambsquarters |
| Burdock | Dichondra | Lespedeza |
| Burweed, lawn (spurweed) | Dock, broadleaf | Mallow, alkali |
| Buttercup, bulbous | Dock, curly | Mallow, bristly |
| Buttercup, creeping | Dogbane | Mallow, common |
| Buttonweed, common (poorjoe) | Dogfennel | Mallow, dwarf |
| Buttonweed, virginia | Dollarweed (Pennywort) | Mallow, venice |
| Carolina geranium (wild) | Dove weed | Matchweed |
| Carpetweed | Elderberry | Mexicanweed |

(cont. on next page)

| 6. Weeds Killed (cont.) | | |
|------------------------------------|---------------------------------|------------------------------|
| Morningglory, bigroot | Pokeweed | Thistle, platte |
| Morningglory, ivyleaf | Prairie Sunflower | Thistle, plumeless |
| Morningglory, mexican | Prickly lettuce (compass plant) | Thistle, russian |
| Morningglory, red | Prickly sida | Thistle, scotch |
| Morningglory, tall | Prostrate knotweed | Thistle, wavyleaf |
| Morningglory, wolly | Puncturevine | Thistle, yellow (horrible) |
| Mugwort | Purslane, common | Trailing crownvetch |
| Mustard, ball | Pusley, brazil | Velvetleaf |
| Mustard, black | Pusley, florida | Vervain, blue |
| Mustard, blue | Pusley, large flower | Vervain, hoary |
| Mustard, hedge | Ragweed, bur | Vervain, prostrate |
| Mustard, Indian | Ragweed, common | Vervain, tall |
| Mustard, tansy | Ragweed, lancleaf | Vetch, bird |
| Mustard, tumble | Ragweed, western | Vetch, hairy |
| Mustard, white | Redstem filaree | Vetch, milk |
| Mustard, wormseed | Rough cinquefoil | Vetch, narrowleaf |
| Nettle, stinging (burning) | Scarlet pimpernel | Vetch, wild |
| Nettle, tall | Shepherd's purse | Virginia creeper |
| Nettle, wood | Smooth chaff-flower | Western clematis |
| Parsley-piert | Smoothdock | Western salsify |
| Parsnip | Sorrel, red (sheep) | Wild aster |
| Pearlwort | Sowthistle, annual | Wild buckwheat |
| Pennycress | Sowthistle, spiny | Wild carrot |
| (frenchweed) | Spanish needles | Wild four-o'clock |
| Pepperweed, perennial | Speedwell, bi-lobed | Wild garlic |
| Pepperweed, virginia (Peppergrass) | Speedwell, common | Wild lettuce |
| Pigweed, prostrate | Speedwell, corn | Wild marigold |
| Pigweed, redroot | Speedwell, ivyleaf | Wild mustard (yellow rocket) |
| Pigweed, russian | Speedwell, persian | Wild onion |
| Pigweed, smooth | Speedwell, purslane | Wild parsnip |
| Pigweed, tumble | Speedwell, slender | Wild radish |
| Pineapple weed | Speedwell, snow | Wild rape |
| Plains coreopsis (tickseed) | Speedwell, thymeleaf | Wild strawberry |
| Plantain, blackseed | Spiny amaranth | Wild sweet potato |
| Plantain, bracted | Spurge, spotted | Wild violet |
| Plantain, broadleaf | Spurge, prostrate | Woodsorrel, creeping |
| Plantain, buckhorn | Spurweed | Woodsorrel, violet |
| Plantain, hoary | Strawberry, India mock | Woodsorrel, yellow (Oxalis) |
| Plantain, narrowleaf | Tansy ragwort | Woolly croton |
| Plantain, slender | Tanweed | Wormseed |
| Plantain, wolly | Thistle, blessed | Yarrow, common |
| Poison ivy | Thistle, bull | Yarrow, western |
| Poison oak | Thistle, canada | Yarrow, yellow |
| | Thistle, flodman (prairie) | |
| | Thistle, musk | |

809/10-2022 AP061214
EPA REG. NO. 2217-539-33955



MANUFACTURED AND DISTRIBUTED BY
PBI/GORDON CORPORATION
P.O. BOX 860350
SHAWNEE, KANSAS 66286
www.GordonsUSA.com

ATTENTION: This specimen label is provided for informational use only. This product may not yet be available for sale in your state or area. The information found in this label may differ from the information found on the product label you are using. Always follow the instructions for use and precautions on the label of the product you are using.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Keep from freezing. Store in original container in a locked storage area inaccessible to children and pets.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling, if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

TRIMEC® is a registered trademark of PBI-Gordon Corporation.