



# Trimec NUTSEdge<sup>®</sup>

**PLUS**  
**LAWN WEED KILLER**  
**CONCENTRATE**

**Kills Yellow Nutsedge &  
200+ Broadleaf Weeds\***

**Treats Northern & Southern Grasses**

**ACTIVE INGREDIENTS:**

2,4-D, dimethylamine salt . . . . .	6.10%
Mecoprop-p, dimethylamine salt . . . . .	2.20%
Dicamba, dimethylamine salt . . . . .	0.57%
Sulfentrazone . . . . .	0.15%

<b>OTHER INGREDIENTS:</b> . . . . .	90.98%
<b>TOTAL</b>	<b>100.00%</b>

**THIS PRODUCT CONTAINS:**

- 0.44 lb 2,4-D acid equivalent per gallon or 5.08%
- 0.16 lb Mecoprop-p acid equivalent per gallon or 1.82%
- 0.04 lb Dicamba acid equivalent per gallon or 0.47%
- 0.01 lb Sulfentrazone per gallon or 0.15%

Isomer Specific by AOAC Method.

\*As listed in attached booklet.

## KEEP OUT OF REACH OF CHILDREN CAUTION



**READ THE ENTIRE LABEL FIRST.  
OBSERVE ALL PRECAUTIONS AND  
FOLLOW DIRECTIONS CAREFULLY.**

### PRECAUTIONARY STATEMENTS

**Hazards to Humans and Domestic Animals**

**CAUTION:** Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wear long-sleeved shirt and long pants, shoes plus socks and gloves. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

First Aid	
<b>If swallowed:</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to by a poison control center or doctor.</li> <li>• Do not give anything to an unconscious person.</li> </ul>
<b>If in eyes:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment advice.	

**Environmental Hazards**

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**SHAKE WELL BEFORE USING.**

**PRECAUTIONS AND RESTRICTIONS:**

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.
- Do not allow people or pets to enter the treated area until sprays have dried.

**1. Weeds Controlled**

**GRASSY WEEDS:**

Yellow Nutsedge

**BROADLEAF WEEDS:**

Trimec<sup>®</sup> Nutsedge Plus Lawn Weed Killer Concentrate will control or suppress the following broadleaf weeds.

Broadleaf Weeds		
Amaranth, purple	Clover, red	Hoary cress
Amaranth, slender	Clover, strawberry	Horsenettle
Aster, heath	Clover, yellow sweet	Horseweed
Aster, white heath	Clover, white sweet	Jimsonweed
Aster, purple	Clover, white	Johnny Jump-up violet
Aster, white prairie	Cockle	Knawel
Artichoke	Cocklebur	Knotweed, prostrate
Austrian fieldcress	Cocklebur, oriental	Kochia
Bedstraw, Catchweed	Cocklebur, spiny	Lambsquarters
Bedstraw, smooth	Common mullein	Lespedeza
Bedstraw, Northern	Corn Spurry	Mallow, alkali
Bedstraw, pineywoods	Creeping Beggarweed	Mallow, bristly
Beggarticks, hairy	Creeping Jenny	Mallow, common
Betony, Florida	(Creeping Charlie)	Mallow, dwarf
Bindweed, Field	Cupid shaving Brush	Mallow, venice
Bindweed, Hedge	Cudweed	Marcela
Bitter Wintercress	Daisy, English	Matchweed
Bittercress, hairy	Daisy, oxeye	Mexicanweed
Bitterweed	Dandelion, common	Moneywort
Black-eyed Susan	Dandelion, false	Morningglory, bigroot
Black Medic	Day flower	Morningglory, ivyleaf
Blood flower milkweed	Deadnettle	Morningglory, mexican
Blue lettuce	Dichondra	Morningglory, red
Brassbuttons	Dock, broadleaf	Morningglory, tall
Bristly oxtongue	Dock, curly	Morningglory, wolly
Broomweed	Dogbane	Mugwort
Buckhorn	Dogfennel	Mustard, ball
Bullnettle	Dollarweed (Pennywort)	Mustard, black
Burclover	Dove weed	Mustard, blue
Burdock	Elderberry	Mustard, hedge
Burweed, lawn	Falseflax	Mustard, Indian,
(spurweed)	False sunflower	Mustard, tansy
Buttercup, bulbous	Fiddleneck	Mustard, tumble
Buttercup, creeping	Filaree, redstem	Mustard, white
Buttonweed, common	Filaree, whitestem	Mustard, wormseed
(poorjoe)	Fleabane, daisy (annual)	Nettle, stinging (burning)
Buttonweed, virginia	Fleabane, rough	Nettle, tall
Carolina geranium (wild)	Galinsoga, hairy	Nettle, wood
Carpetweed	Galinsoga, smallflower	Parsley-piert
Catnip	Goldenrod	Parsnip
Catsear, spotted (false	Ground ivy	Pearlwort
dandelion)	Groundsel	Pennycress
Chamber bitter	Gumweed	(frenchweed)
Chickweed, common	Hairy fleabane	Pepperweed, perennial
Chickweed, mouseear	Hawkweed, meadow	Pepperweed, virginia
Chickweed, sticky	Hawkweed, mouseear	(Peppergrass)
Chicory	Hawkweed, orange	Pigweed, prostate
Cinquefoil	Hawkweed, yellow	Pigweed, redroot
Clover, crimson	Healall	Pigweed, russian
Clover, alyce	Heartleaf drymary	Pigweed, smooth
Clover, hop	Hemp	Pigweed, tumble
Clover, rabbit foot	Henbit	

(cont. on next page)

Broadleaf Weeds (cont.)		
Pineapple weed	Smoothdock	Vervain, prostrate
Plains coreopsis (tickseed)	Sorrel, red (sheep)	Vervain, tall
Plantain, blackseed	Sowthistle, annual	Vetch, bird
Plantain, bracted	Sowthistle, spiny	Vetch, hairy
Plantain, broadleaf	Spanish needles	Vetch, milk
Plantain, buckhorn	Speedwell, bi-lobed	Vetch, narrowleaf
Plantain, hoary	Speedwell, common	Vetch, wild
Plantain, narrowleaf	Speedwell, corn	Virginia creeper
Plantain, slender	Speedwell, ivyleaf	Western clematis
Plantain, woolly	Speedwell, persian	Western salsify
Poison ivy	Speedwell, purslane	Wild aster
Poison oak	Speedwell, slender	Wild buckweed
Pokeweed	Speedwell, snow	Wild carrot
Prairie Sunflower	Speedwell, thymeleaf	Wild four-o'clock
Prickly lettuce (compass plant)	Spiny amaranth	Wild garlic
Prickly sida	Spurge, spotted	Wild lettuce
Prostrate knotweed	Spurge, prostrate	Wild marigold
Puncturevine	Spurweed	Wild mustard (yellow rocket)
Purslane, common	Star of Bethlehem	Wild onion
Pusley, brazil	Strawberry, India mock	Wild parsnip
Pusley, florida	Tansy ragwort	Wild radish
Pusley, large flower	Tanweed	Wild rape
Ragweed, bur	Thistle, blessed	Wild strawberry
Ragweed, common	Thistle, bull	Wild sweet potato
Ragweed, lanleaf	Thistle, canada	Wild violet
Ragweed, western	Thistle, flodman (prairie)	Woodsorrel, creeping
Redstem filaree	Thistle, musk	Woodsorrel, violet
Rough cinquefoil	Thistle, platte	Woodsorrel, yellow (Oxalis)
Scarlet pimpernel	Thistle, scotch	Woolly croton
Shepherd's purse	Thistle, wavyleaf	Wormseed
Smartweed, ladysthumb	Thistle, yellow (horrible)	Yarrow, common
Smartweed, pale	Trailing crownvetch	Yarrow, western
Smartweed, pennsylvania	Velvetleaf	Yarrow, yellow
Smooth chaff-flower	Vervain, blue	
	Vervain, hoary	

**2. Where To Use**

This product is intended for use by homeowners on lawns including:

**Cool season turfgrass:**

- Kentucky bluegrass
- Perennial ryegrass
- Fescue spp., including tall, red and fine leaf fescues
- Bentgrass

**Warm season turfgrass:**

- Bermudagrass
- Bahiagrass
- Zoysiagrass
- St. Augustinegrass\*
- Centipedegrass

\*NOTE: Do not use on 'Floratum' or other improved varieties of St. Augustinegrass in Florida. Do not use this product on carpetgrass, dichondra, legumes, or lawns containing desirable clovers.

**3. When To Use**

Apply when daytime temperatures are between 50° and 85°F

**SPRING:** Spray when weeds are young and actively growing with adequate soil moisture for best results.

**FALL:** Spray when weeds are young and actively growing with adequate soil moisture for best results. Fall applications will control weeds that may otherwise go dormant through the winter and resprout the following spring.

**Limitations on broadcast applications to residential turfgrass**

The maximum application rate is 5.0 fl.oz. of product per 1,000 sq.ft. per application. The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 10 fl.oz. of product per 1,000 sq.ft.

**3.1 Spot Treatment**

To treat small areas or individual weeds use a pressure sprayer for better control. Do not use a hose-end sprayer for Spot Treatments. MIX 1.25 FL.OZ. (2.5 TABLESPOONS) OF PRODUCT IN 1 GALLON OF WATER. Shake well before spraying. Spray the target weeds thoroughly. Nozzle adjustments may be necessary to deliver small spray droplets of the target weeds for uniform coverage.

**Limitations on spot treatments:**

Spot treatment is defined as a treatment area no greater than 1,000 sq.ft. per acre. The maximum application rate is 5.0 fl.oz. per 1,000 sq.ft. per application. The maximum number of spot treatments is limited to 2 per year with a minimum of 30 days between applications.

**3.2 Entire Lawn Treatments**

Three easy steps to mix and apply with hand-operated sprayers:

- 1) Measure the total lawn area to be treated. This is done by multiplying the length of the area by the width to obtain total square feet. Prepare only enough spray solution to treat the area calculated.
- 2) Select the application rate from Table 1 below. Carefully measure the proper amount of product to treat 500 sq.ft. If more or less than 500 sq.ft. is to be treated, adjust the amount of product accordingly.
- 3) Shake the spray solution before and during application. Use a spray pattern that gives small droplets without any fine mist.

**FOR HAND-OPERATED SPRAYERS (BACKPACK, KNAPSACK, TANK-TYPE, OR PUMP-UP)**

Table 1. Amount of product to be diluted in 1 gallon water required to treat 500 sq.ft. of lawn with pressure sprayers.	
Turfgrass	Amount of Product
St. Augustinegrass*, Bermudagrass, Bentgrass, and Centipedegrass	1.25 fl.oz. (2.5 Tbsp)
Kentucky bluegrass, Fescue spp., Perennial ryegrass, Zoysiagrass and Bahiagrass	2.5 fl.oz. (5 Tbsp)
*Do not use on 'Floratum' or other improved varieties of St. Augustinegrass in Florida. For pressure sprayers, mix the indicated amount of product listed above with 1 gallon of water for each 500 sq.ft. to be treated.	

**FOR DIAL-IN HOSE END SPRAYERS WITH NO PRE-MIXING**

- 1) From the Table below, determine amount of Trimec® Nutsedge Plus Lawn Weed Killer Concentrate to be applied for each 1,000 sq.ft. of lawn.
- 2) Fill sprayer jar with enough concentrate to cover area to be treated. Do not add additional water to the sprayer jar.
- 3) Set dial on sprayer to the "Rate per Gallon" setting from the Table below.
- 4) Attach sprayer to hose. Extend hose to the farthest area of lawn to be treated and work back toward faucet so that you do not walk over treated areas. Apply evenly to lawn.
- 5) Unused product can be returned to the original container when completed.

Table 2. Application rates for turfgrass with hose-end sprayers		
Turfgrass	Dial settings or Rate per gallon	Amount of herbicide applied per 1000 sq.ft.
St. Augustinegrass*, Bentgrass, Bermudagrass, and Centipedegrass	2 Tablespoons (6 teaspoons) (1 ounce)	2.5 fl.oz.
Kentucky bluegrass, Fescue spp., Perennial Ryegrass, Zoysiagrass and Bahiagrass	4 Tablespoons (12 teaspoons) (2 ounces)	5.0 fl.oz.
*Do not use on 'Floratum' or other improved varieties of St. Augustinegrass in Florida.		

An example for a hose-end sprayer is shown below:

- For Kentucky bluegrass: To treat 1,000 sq.ft. of lawn, measure 5.0 fl.oz. of product into sprayer jar. **Do not add water.** Set dial on 4 Tbsp and spray uniformly.
- **To BEGIN spraying:** Attach hose to sprayer. Extend the garden hose to the farthest section of lawn to be treated. Rotate sprayer nozzle tip to direct spray up or down. Turn on water and squeeze trigger or rotate on/off knob to begin.
- **AFTER spraying:** Unused product or concentrate can be poured back into the original container. Keep the dial setting and clean the sprayer by rinsing the sprayer jar with water for one minute.

Check the directions you received when you purchased your sprayer for further details on the proper use of your dial sprayer.

**FOR CONCENTRATE-PLUS-WATER SPRAYERS OR HOSE-END SPRAYERS WITH PRE-MIXING**

For sprayers that require the addition of water as well as concentrate prior to application:

- 1) From the following Table 3, determine amount of concentrate required to treat each 1,000 sq.ft. of lawn.
- 2) Pour concentrate into sprayer jar to the required "fl.oz." level.
- 3) Then add the appropriate amount of water to the "gals." level. Replace sprayer top on jar and shake well before spraying.

- 4) Attach sprayer to hose. Extend hose to the farthest area of lawn to be treated and work back toward faucet so that you do not walk over treated areas. Apply evenly to lawn.

**Table 3. Application rates for turfgrass with hose-end sprayers that require pre-mixing.**

Size of area to be treated	St. Augustinegrass, Bermudagrass, Bentgrass, Centipedegrass	Kentucky bluegrass, Fescue spp., Perennial ryegrass, Zoysiagrass and Bahiagrass	Gallons of water
1,000 sq.ft	2.5 fl.oz.	5.0 fl.oz.	2 gal
5,000 sq.ft.	12.5 fl.oz.	25.0 fl.oz.	10 gal
6,000 sq.ft.	15.0 fl.oz.	30.0 fl.oz.	12 gal

\*Do not use on 'Floritam' or other improved varieties of St. Augustinegrass in Florida.

**EXAMPLE:** To treat 1,000 sq.ft. of fescue pour in concentrate to the 5 fl.oz. mark and add water to the 2 gallon mark. Check the directions you received when you purchased your sprayer for further details on the proper use of your sprayer.

**4. Use Tips**

- 1) Spray during growing season when weeds are actively growing.
- 2) For best results, mow lawn 2 to 3 days before treating weeds.
- 3) To avoid risk of lawn injury, do not spray drought-stressed lawns.
- 4) Do not water within 6 hours after treatment. Poor weed control may result if rainfall occurs within 3 hours of application.
- 5) Do not apply this product to newly seeded grasses until well-established or after the third mowing.
- 6) Treated areas may be reseeded 3 to 4 weeks after application.
- 7) Adjust nozzle to produce a coarse spray. Coarse sprays are less likely to drift. Avoid fine mists.
- 8) If certain tough weeds are not dead within 3 weeks, repeat application using spot treatment directions.

**PROHIBITIONS:**

- Do not use on 'Floritam' or other improved varieties of St. Augustinegrass in Florida.
- Do not use this product on carpetgrass, dichondra, legumes, or lawns containing desirable clovers.
- Do not treat when air temperatures exceed 85°F as damage to grass may occur.
- Application at rates above specified on this label may cause injury to turfgrass.
- Do not apply this product to food crops (gardens), forage crops, or ornamental plants including flowers, trees, shrubs, hedges, woody ornamentals, or groundcovers established in landscape plantings.
- Do not spray exposed roots of ornamentals and trees.

**Other Use Precautions:**

- Do not apply as a fine mist because of potential for injury to desirable plants. Do not use this product on or near desirable plants, including within the drip line of the roots of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day. Rinse applicator over lawn area only.

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage and disposal.

**PESTICIDE STORAGE:** Store in original container in a locked storage area inaccessible to children or pets.

**PESTICIDE DISPOSAL AND CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

**LIMITED WARRANTY AND DISCLAIMER**

FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS

FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

Timec® is a registered trademark of PBI-Gordon Corporation.

855/6-2013 AP082611  
EPA REG. NO. 2217-880



**MANUFACTURED BY**  
**PBI/GORDON CORPORATION**  
**1217 WEST 12TH STREET**  
**KANSAS CITY, MISSOURI 64101**  
[www.GordonsUSA.com](http://www.GordonsUSA.com)

ATTENTION: This specimen label is provided for informational use only. This pesticide product may not yet be available for sale in your state or area. The information found in this label may differ from the information found on the product label you are using. Always follow the instructions for use and precautions on the label of the pesticide you are using.