



Trimec

READY SPRAY

LAWN WEED KILLER

ACTIVE INGREDIENTS:

2,4-D, dimethylamine salt	5.56%
Mecoprop-p, dimethylamine salt	1.34%
Dicamba, dimethylamine salt	0.62%

OTHER INGREDIENTS:	92.48%
TOTAL	100.00%

THIS PRODUCT CONTAINS:

- 4.62% 2,4-dichlorophenoxyacetic acid equivalent (CAS 2008-39-1)
 - 1.11% (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid equivalent (CAS 66423-09-4)
 - 0.51% 3,6-dichloro-o-anisic acid equivalent (CAS 2300-66-5)
- Isomer Specific by AOAC Method.

KEEP OUT OF REACH OF CHILDREN CAUTION



**READ THE ENTIRE LABEL FIRST.
OBSERVE ALL PRECAUTIONS AND
FOLLOW DIRECTIONS CAREFULLY.**

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Harmful if absorbed through skin or swallowed. Avoid contact with skin, eyes or clothing. Wear protective eyewear. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

First Aid	
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also call toll-free 1-877-800-5556 for emergency medical treatment information.	

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

PRECAUTIONS AND RESTRICTIONS:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.
- Do not allow people or pets to enter the treated area until sprays have dried.

IMPORTANT: Do not exceed specified dosages for any area.

FOR USE ON LAWNS:

This product is designed for use through this sprayer for controlling broadleaf weeds in bahiagrass, bermudagrass, Kentucky bluegrass, fescue, perennial ryegrass, zoysiagrass, centipedegrass, and St. Augustinegrass lawns, vacant lots, playgrounds and other similar areas.

For control of dandelion, chickweed, plantain, clover, and other listed weeds, apply when weeds are actively growing. Spray weeds such as wild garlic and wild onion when they appear in late fall and again in early spring.

Water lawn thoroughly 1 to 2 days before application. Do not water for 24 hours after application. Do not mow for one week before treating.

Spray entire lawn, wetting weeds and lawn grasses, do not saturate. Adjust water pressure at faucet to give coarse spray droplets. Do not apply when temperatures are expected to exceed 85°F within 24 hours following application.

Temporary injury may occur on bentgrass, bermudagrass, and red fescue lawn grasses. Do not apply to carpetgrass, dichondra, Lippia lawns, and lawns where desirable clovers are present.

Do not apply this product to improved varieties of 'Floritam' St. Augustinegrass in Florida.

Newly seeded lawns may be treated after they have been mowed at least three times.

Other Use Precautions:

- Do not apply as a fine mist because of potential for injury to desirable plants.
- Do not use this product on or near desirable plants, including contact of spray on exposed root systems or adventitious shoots within the drip line of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.

HOW TO USE THE TRIMEC® READY SPRAY

Shake well before using. Connect attached spray nozzle to garden hose. Turn the water control "ON/OFF" valve on top of spray nozzle to the "OFF" position. Turn on water at faucet. Extend hose to the farthest area of lawn to be treated and work back towards faucet so that you do not walk over treated areas. Using a coin or key, rotate the small product control valve located near front of nozzle forward to the "OPEN" position.

To begin spraying, point spray nozzle towards lawn and turn the water control "ON/OFF" valve to the "ON" position. The water automatically mixes with the product. Walk at a steady pace while spraying using an even sweeping motion, slightly overlapping treated areas. Refer to graduated scale on side of container to guide coverage.

To stop spraying, turn water control "ON/OFF" valve to the "OFF" position. Rotate small product control valve back to "CLOSED" position. Turn off water at faucet. To relieve pressure, turn water control "ON/OFF" valve to the "ON" position and point sprayer away from any desirable plants before removing unit from hose.

One quart of Trimec® Ready Spray Lawn Weed Killer will cover 5,000 square feet of lawn for bahiagrass, Kentucky bluegrass, bermudagrass, fescue, perennial ryegrass, and zoysiagrass.

For centipedegrass and St. Augustinegrass, one quart of Trimec® Ready Spray Lawn Weed Killer will treat 8,000 square feet of lawn.

Limitations on broadcast applications to residential turfgrass:

The maximum application rate is 6.4 fl.oz. of product per 1,000 sq.ft. per application. The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 12.8 fl.oz. of product per 1,000 sq.ft.

WEEDS CONTROLLED:

This product is effective on hard to control broadleaf weed species such as dandelion, white clover, sheep sorrel and spurge. In addition, Trimec® Ready Spray Lawn Weed Killer controls susceptible weeds including chickweed, henbit, and plantain. A list is presented as follows:

WEEDS		
Amaranth, purple	Dichondra	Pennycress
Amaranth, slender	Dock, broadleaf	(frenchweed)
Aster, heath	Dock, curly	Pepperweed, perennial
Aster, white heath	Dogbane	Pepperweed, Virginia
Aster, purple	Dogfennel	(Peppergrass)
Aster, white prairie	Dollarweed (Pennywort)	Pigweed, prostrate
Austrian fieldcress	Dove weed	Pigweed, redroot
Bedstraw, Catchweed	Elderberry	Pigweed, russian
Bedstraw, smooth	Falseflax	Pigweed, smooth
Bedstraw, Northern	False sunflower	Pigweed, tumble
Bedstraw, pineywoods	Fiddleneck	Pineapple weed
Beggarticks, hairy	Filaree, redstem	Plains coreopsis
Betony, Florida	Filaree, whitestem	(tickseed)
Bindweed, Field	Fleabane, daisy	Plantain, blackseed
Bindweed, Hedge	(annual)	Plantain, bracted
Bitter Wintercress	Fleabane, rough	Plantain, broadleaf
Bittercress, hairy	Galinisoga, hairy	Plantain, buckhorn
Bitterweed	Galinisoga, smallflower	Plantain, hoary
Black-eyed Susan	Goldenrod	Plantain, narrowleaf
Black Medic	Ground ivy	Plantain, slender
Blood flower milkweed	Gumweed	Plantain, woolly
Blue lettuce	Hairy fleabane	Poison ivy
Brassbuttons	Hawkweed, meadow	Poison oak
Bristly oxtongue	Hawkweed, mouseear	Pokeweed
Broomweed	Hawkweed, orange	Prairie Sunflower
Buckhorn	Hawkweed, yellow	Prickly lettuce
Bullnettle	Healall	(compass plant)
Burclover	Heartleaf drymary	Prickly sida
Burdock	Hemp	Prostrate knotweed
Burweed, lawn	Henbit	Puncturevine
(spurweed)	Hoary cress	Purslane, common
Buttercup, bulbous	Horsenettle	Pusley, brazil
Buttercup, creeping	Jimsonweed	Pusley, Florida
Buttonweed, common	Knawel	Pusley, large flower
(poorjoe)	Knotweed, prostrate	Ragweed, bur
Buttonweed, Virginia	Kochia	Ragweed, common
Carolina geranium	Lambsquarters	Ragweed, lancleaf
(wild)	Lespedeza	Ragweed, western
Carpetweed	Mallow, alkali	Redstem filaree
Catnip	Mallow, bristly	Rough cinquefoil
Catsear, spotted (false	Mallow, common	Scarlet pimpernel
dandelion)	Mallow, dwarf	Shepard's purse
Chickweed, common	Mallow, venice	Smooth chaff-flower
Chickweed, mouseear	Matchweed	Smoothdock
Chickweed, sticky	Mexicanweed	Sorrel, red (sheep)
Chicory	Morningglory, bigroot	Sowthistle, annual
Cinquefoil	Morningglory, ivyleaf	Sowthistle, spiny
Clover, crimson	Morningglory, Mexican	Spanish needles
Clover, alyce	Morningglory, red	Speedwell, bi-lobed
Clover, hop	Morningglory, tall	Speedwell, common
Clover, rabbit foot	Morningglory, wolly	Speedwell, corn
Clover, red	Mugwort	Speedwell, ivyleaf
Clover, strawberry	Mustard, ball	Speedwell, Persian
Clover, yellow sweet	Mustard, black	Speedwell, purslane
Clover, white sweet	Mustard, blue	Speedwell, slender
Clover, white	Mustard, hedge	Speedwell, snow
Cockle	Mustard, Indian,	Speedwell, thymeleaf
Cocklebur	Mustard, tansy	Spiny amaranth
Cocklebur, oriental	Mustard, tumble	Spurge, spotted
Cocklebur, spiny	Mustard, white	Spurge, prostrate
Common mullein	Mustard, wormseed	Spurweed
Creeping Jenny	Nettle, stinging	Strawberry, India mock
(Creeping Charlie)	(burning)	Tansy ragwort
Cudweed	Nettle, tall	Tanweed
Daisy, English	Nettle, wood	Thistle, blessed
Daisy, oxeye	Parsley-piert	Thistle, bull
Dandelion, common	Parsnip	Thistle, Canada
Dandelion, false	Pearlwort	(cont. on next column)
Deadnettle		

WEEDS (cont.)		
Thistle, flodman (prairie)	Vetch, hairy	Wild onion
Thistle, musk	Vetch, milk	Wild parsnip
Thistle, platte	Vetch, narrowleaf	Wild radish
Thistle, plumeless	Vetch, wild	Wild rape
Thistle, Russian	Virginia creeper	Wild strawberry
Thistle, scotch	Western clematis	Wild sweet potato
Thistle, wavyleaf	Western salsify	Wild violet
Thistle, yellow (horrible)	Wild aster	Woodsorrel, creeping
Trailing crownvetch	Wild buckwheat	Woodsorrel, violet
Velvetleaf	Wild carrot	Woodsorrel, yellow (Oxalis)
Vervain, blue	Wild four-o'clock	Woolly croton
Vervain, hoary	Wild garlic	Wormseed
Vervain, prostrate	Wild lettuce	Yarrow, common
Vervain, tall	Wild marigold	Yarrow, western
Vetch, bird	Wild mustard (yellow rocket)	Yarrow, yellow

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container in a locked storage area inaccessible to children or pets. Keep from freezing.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Triple rinse container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

Gordon's® logo and Trimec® are registered trademarks of PBI-Gordon Corporation.

827/8-2016 AP060716
EPA REG. NO. 2217-540



MANUFACTURED BY
PBI/GORDON CORPORATION
1217 WEST 12TH STREET
KANSAS CITY, MISSOURI 64101
www.GordonsUSA.com

ATTENTION: This specimen label is provided for informational use only. This product may not yet be available for sale in your state or area. The information found in this label may differ from the information found on the product label you are using. Always follow the instructions for use and precautions on the label of the product you are using.