



Trimec SPEED

LAWN WEED KILLER CONCENTRATE

ACTIVE INGREDIENTS:

2,4-D, 2-ethylhexyl ester	4.01%
Mecoprop-p acid	0.49%
Dicamba acid	0.27%
Carfentrazone-ethyl	0.16%

OTHER INGREDIENTS:	95.07%
	TOTAL 100.00%

THIS PRODUCT CONTAINS:

0.226 lb 2,4-dichlorophenoxyacetic acid equivalent per gallon or 2.66%
0.042 lb (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per gallon or 0.49%
0.023 lb 3,6-dichloro-o-anisic acid equivalent per gallon or 0.27%
0.014 lb Ethyl α,2-dichloro-5-[4(difluoromethyl)-4,5-dihydro-3-methyl-5-oxo-1H-1,2,4-triazol-1-yl]-4-fluorobenzenepropanoate per gallon or 0.16%

KEEP OUT OF REACH OF CHILDREN CAUTION



**READ THE ENTIRE LABEL FIRST.
OBSERVE ALL PRECAUTIONS AND
FOLLOW DIRECTIONS CAREFULLY.**

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if absorbed through the skin. Avoid contact with skin, eyes, or clothing. When using this product, wear long-sleeved shirt and long pants, socks, shoes, and gloves. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

First Aid	
If on skin:	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment advice.	

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

Physical and Chemical Hazards

Combustible. Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

1. Product Description

Trimec® Speed Lawn Weed Killer Concentrate is a selective herbicide that provides post-emergent control of listed broadleaf weeds in established residential lawns. Trimec Speed Lawn Weed Killer Concentrate may be applied to lawns containing Kentucky bluegrass, perennial ryegrass, fescues, common and hybrid bermudagrass, bahiagrass, bentgrass, buffalograss, centipedegrass, zoysiagrass, and St. Augustinegrass (except Floratam). Trimec Speed Lawn Weed Killer Concentrate will control or suppress more than 250 broadleaf weeds, including dandelion, clover, henbit, oxalis, chickweed, knotweed, spurge, plantain, and many others as listed. Results will be visible within hours of application. Symptoms may include leaf reddening, leaf and stem curling and twisting, and chlorosis.

2. Use Restrictions

- For residential use only
- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until sprays have dried.
- Do not apply as a fine mist because of potential for injury to desirable plants. Do not use this product on or near desirable plants, including contact of spray on exposed root systems or adventitious shoots within the drip line of desirable trees and shrubs, since injury may result.
- Do not use on Carpetgrass, Dichondra, food crops (gardens, fruits, and vegetables), forage crops, or ornamental plants (flowers, trees, shrubs, hedges, woody ornamentals, groundcovers, groundcovers established in landscape plantings), 'Floratam' St. Augustinegrass, and turfgrass species that are not listed on this label.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day. Rinse applicator over lawn area only.

3. How Much To Apply

Over-application or rates above those specified on this label can cause turf injury.

For Kentucky Bluegrass, Bahiagrass, Common Bermudagrass, Fine Fescues, Tall Fescue, Perennial Ryegrass, and Zoysiagrass

Spot Treatments: Mix 3 fl.oz. (6 Tbsp) in 1 gallon of water to treat 500 sq.ft.

Entire Lawn Treatments:

- Pump Style Sprayer: Mix 6 fl.oz. in 0.5 to 1 gallon of water to treat 1000 sq.ft.
- Hose-End Sprayer: Apply 6 fl.oz. per 1000 sq.ft.
- 12 volt & Tow-Behind Sprayers: Mix 6 fl.oz. in 0.5 to 1 gallon of water to treat 1000 sq.ft.

For Hybrid Bermudagrass, Buffalograss, Centipedegrass, St. Augustinegrass (do not use on 'Floratam' St. Augustinegrass), and Bentgrass.

Spot Treatments: Mix 1.5 fl.oz. (3 Tbsp) in 1 gallon of water to treat 500 sq.ft.

Entire Lawn Treatments:

- Pump Style Sprayer: Mix 3 fl.oz. in 0.5 to 1 gallon of water to treat 1000 sq.ft.
- Hose-End Sprayer: Apply 3 fl.oz. per 1000 sq.ft.
- 12 volt & Tow-Behind Sprayers: Mix 3 fl.oz. in 0.5 to 1 gallon of water to treat 1000 sq.ft.

The maximum rate is 6 fl.oz. of product per 1000 sq.ft per application and the maximum number of broadcast applications is 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 12 fl.oz. of product per 1000 sq.ft. excluding spot treatments.

4. When To Use

Spray when broadleaf weeds are young and actively growing for best results. This product may be applied in the spring, summer, or fall. Spring and fall treatments under adequate soil moisture conditions are preferred to the summer treatments. Generally, summer broadcast applications to older, drought stressed weeds are less effective.

Use broadcast (entire areas) applications or follow-up applications at 30 day intervals for more mature weeds, for dense infestations, and for adverse environmental conditions. Spot treatments during the summer may be appropriate for sparse infestations, as a follow-up treatment, or any time broadleaf weeds are susceptible.

- **For Established Lawns:** Apply when daytime temperatures are below 90°F.
- **For Newly Seeded Areas:** Apply this product to newly seeded grasses when well-established (approximately 6 weeks after seedling emergence) or after the third mowing.
- **For Newly Sodded, Sprigged or Plugged areas:** Apply this product 3 to 4 weeks after the sodding, sprigging, or plugging operations.
- **Reseeding:** Treated areas can be reseeded at 2 weeks after the application of this product.
- **Irrigation:** Adequate soil moisture and favorable growing conditions enhance the performance of this product. If dry conditions exist, schedule irrigations (watering) before and after the application. Avoid applications of this product when turf is under stress since injury may result.
- **Mowing:** Avoid mowing within 1 to 2 days before and after the application of this product.

5. How To Apply

5.1 Pump Style Sprayers Ideal for spraying individual weeds

1. Add the appropriate amount of water and concentrate to the sprayer tank.
2. Close sprayer, shake well, and pump handle to pressurize.
3. Adjust nozzle to deliver a coarse spray pattern.
4. To broadcast over large areas: apply evenly to treated area
5. To treat individual weeds: spray to wet weed leaf surfaces.
6. Re-pressurize the sprayer as needed to maintain a good spray pattern.

5.2 12 Volt & Tow-Behind Boom Sprayers

Ideal for treating large country and suburban lawns

1. To calibrate your application equipment consult your equipment owner's manual for speed and pressure settings required to deliver 0.5 to 1 gallon of spray solution to treat 1000 sq.ft.. For example, at 4 mph a sprayer might deliver 1/2 gallon per 1000 sq.ft. at 40 psi.
2. To apply evenly, drive at a constant speed that delivers the prescribed amount of spray required for the area to be treated.
3. Mix the appropriate amount of product and water in the sprayer using the following tables as a guide.

Tank Size	Spray Volume	Coverage Area	Concentrate to add to tank
15 gal	0.5 gal/1000 sq.ft.	30,000 sq.ft.	90 to 180 fl. oz. (0.7 to 1.4 gal.)
	1 gal/1000 sq.ft.	15,000 sq.ft.	45 to 90 fl. oz. (0.4 to 0.7 gal)
25 gal	0.5 gal/1000 sq.ft.	50,000 sq.ft.	150 to 300 fl.oz. (1.2 to 2.3 gal)
	1 gal/1000 sq.ft.	25,000 sq.ft.	75 to 150 fl.oz. (0.6 to 1.2 gal)

For spot treatments with 12 Volt & Pull Behind Sprayers

See rates under "How Much To Apply," Spot Treatments.

5.3 Dial Style Hose-End Sprayers

Ideal for treating medium-sized lawns

1. Fill sprayer with sufficient concentrate to cover the total area to be treated. DO NOT ADD WATER.
2. Close sprayer and set dial to 2.5 oz.
3. Connect sprayer to hose
4. Extend hose to furthest point from faucet. Walking back towards the faucet, begin spraying, avoiding contact with the treated area.
5. Pour any unused product back into this container.

6. Use Precautions

- Delay mowing 2 days before and until 2 days after the application of this product.
- If the lawn is dry, water 24 hours before using this product.
- For best results, delay watering lawn until 2 hours after treatment.
- For new lawns, apply only when grass is well-established (approximately 6 weeks after seedling emergence) or after the third mowing.
- Bare spots may be seeded 3 to 4 weeks after application.
- Spray when air temperatures are cooler than 90°F.
- Avoid applications during the spring transition of the warm season grasses and during periods of extremely high temperatures.

7. Broadleaf Weeds Controlled

Amaranth, purple	Fleabane, rough	Pusley, large flower
Amaranth, slender	Galinsoga, hairy	Ragweed, bur
Aster, heath	Galinsoga, smallflower	Ragweed, common
Aster, white heath	Goldenrod	Ragweed, lance-leaf
Aster, purple	Ground ivy	Ragweed, western
Aster, white prairie	Groundsel	Redstem filaree
Artichoke	Gumweed	Rough cinquefoil
Austrian fieldcress	Hairy fleabane	Scarlet pimpernel
Bedstraw, catch-weed	Hawkweed, meadow	Shepherd's purse
Bedstraw, smooth	Hawkweed, mouse-ear	Smartweed,
Bedstraw, northern	Hawkweed, orange	ladysthumb
Bedstraw, pineywoods	Hawkweed, yellow	Smartweed, pale
Beggar-ticks, hairy	Healall	Smartweed,
Betony, Florida	Heart-leaf drymary	Pennsylvania
Bindweed, field	Hemp	Smooth chaff-flower
Bindweed, hedge	Henbit	Smooth dock
Bitter Wintercress	Hoary-cress	Sorrel, red (sheep)
Bitter-cress, hairy	Horsenettle	Sowthistle, annual
Bitterweed	Horseweed	Sowthistle, spiny
Black-eyed Susan	Jimsonweed	Spanish needles
Black medic	Johnny jump-up violet	Speedwell, bi-lobed
Blood-flower milkweed	Knawel	Speedwell, common
Blue lettuce	Knotweed, prostrate	Speedwell, corn
Brassbuttons	Kochia	Speedwell, ivyleaf
Bristly oxtongue	Lambsquarters	Speedwell, Persian
Broomweed	Lespedeza	Speedwell, purslane
Buckhorn	Mallow, alkali	Speedwell, slender
Bull-nettle	Mallow, bristly	Speedwell, snow
Bur-clover	Mallow, common	Speedwell, thyme-leaf
Burdock	Mallow, dwarf	Spiny amaranth
Burweed, lawn	Mallow, Venice	Spurge, spotted
Buttercup, bulbous	Marcela	Spurge, prostrate
Buttercup, creeping	Matchweed	Star of Bethlehem
Buttonweed, common (poorjoe)	Mexican-weed	Strawberry,
Buttonweed, Virginia	Moneywort	Indian cock
Carolina geranium	Morningglory, bigroot	Tansy ragwort
Carpetweed	Morningglory, ivyleaf	Tanweed
Catnip	Morningglory, red	Thistle, blessed
Catsear, spotted (false dandelion)	Morningglory, tall	Thistle, bull
Chamber bitter	Morningglory, wolly	Thistle, Canada
Chickweed, common	Mugwort	Thistle, Flodman's
Chickweed, mouse-ear	Mustard, ball	Thistle, musk
Chickweed, sticky	Mustard, black	Thistle, plumeless
Chicory	Mustard, blue	Thistle, prairie
Cinquefoil	Mustard, hedge	Thistle, Russian
Clover, crimson	Mustard, Indian	Thistle, Scotch
Clover, Alyce	Mustard, tansy	Thistle, wavy-leaf
Clover, hop	Mustard, tumble	Thistle, yellow
Clover, rabbit-foot	Mustard, white	Trailing crown-vetch
Clover, red	Mustard, wormseed	Velvetleaf
Clover, strawberry	Nettle, stinging	Vervain, blue
Clover, yellow sweet	Nettle, tall	Vervain, hoary
Clover, white sweet	Nettle, wood	Vervain, prostrate
Clover, white	Parsley-piert	Vervain, tall
Cockle	Parsnip	Vetch, bird
Cocklebur	Pearlwort	Vetch, hairy
Cocklebur, spiny	Pennycress	Vetch, milk
Common mullein	Pepperweed, perennial	Vetch, narrow-leaf
Corn spurry	Pepperweed, Virginia (Peppergrass)	Vetch, wild
Creeping beggar-weed	Pigweed, prostrate	Virginia creeper
Creeping Jenny (Creeping Charlie)	Pigweed, redroot	Western clematis
Cupid's shaving brush	Pigweed, Russian	Western salsify
Cudweed	Pigweed, smooth	Wild aster
Daisy, English	Pigweed, tumble	Wild buckwheat
Daisy, oxeye	Pineapple weed	Wild carrot
Dandelion, common	Plains coreopsis (tickseed)	Wild four-o'clock
Dandelion, false	Plantain, black-seed	Wild garlic
Day-flower	Plantain, bracted	Wild lettuce
Deadnettle	Plantain, broadleaf	Wild marigold
Dichondra	Plantain, buckhorn	Wild mustard
Dock, broadleaf	Plantain, hoary	Wild onion
Dock, curly	Plantain, narrow-leaf	Wild parsnip
Dogbane	Plantain, slender	Wild radish
Dog-fennel	Plantain, woolly	Wild rape
Dollarweed (Pennywort)	Poison ivy	Wild strawberry
Dove-weed	Poison oak	Wild sweet potato
Elderberry	Pokeweed	Wild violet
Falseflax	Prairie sunflower	Woodsorrel, creeping
False sunflower	Prickly lettuce (compass plant)	Woodsorrel, violet
Fiddleneck	Prickly sida	Woodsorrel, yellow (Oxalis)
Filaree, redstem	Prostrate knotweed	Woolly croton
Filaree, whitestem	Puncturevine	Wormseed
Fleabane, daisy (annual)	Purslane, common	Yarrow, common
	Pusley, Brazilian	Yarrow, western
	Pusley, Florida	Yarrow, yellow
		Yellow Rocket

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in original container in a locked storage area inaccessible to children or pets. Keep from freezing.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

Gordon's® logo and Trimec® are registered trademarks of PBI-Gordon Corporation.

810/10-2022 AP022218
EPA REG. NO. 2217-1045



MANUFACTURED BY
PBI/GORDON CORPORATION
P.O. BOX 860350
SHAWNEE, KANSAS 66286
www.GordonsUSA.com

ATTENTION: This specimen label is provided for informational use only. This product may not yet be available for sale in your state or area. The information found in this label may differ from the information found on the product label you are using. Always follow the instructions for use and precautions on the label of the product you are using.