TOTAL 100.00%



ACTIVE INGREDIENTS:

2,4-D, 2-ethylhexyl ester	4.01%
Mecoprop-p acid	0.49%
Dicamba acid	0.27%
Carfentrazone-ethyl	0.16%
OTHER INGREDIENTS:	95.07%

THIS PRODUCT CONTAINS:

0.226 lb 2,4-dichlorophenoxyacetic acid equivalent per gallon or 2.66% 0.042 lb (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid equivalent per gallon or 0.49%

0.023 lb 3,6-dichloro-o-anisic acid equivalent per gallon or 0.27% 0.014 lb Ethyl α,2-dichloro-5-[4(difluoromethyl)-4,5-dihydro-3-methyl-5-oxo-1H-1,2,4-triazol-1-yl]-4-fluorobenzenepropanoate per gallon or 0.16%

KEEP OUT OF REACH OF CHILDREN CAUTION



PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if absorbed through the skin. Avoid contact with skin, eyes, or clothing. When using this product, wear long-sleeved shirt and long pants, socks, shoes, and gloves. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

First Aid

If on skin:

- · Take off contaminated clothing.
- · Rinse skin immediately with plenty of water for 15 20 minutes.
- · Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment advice.

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

Physical and Chemical Hazards

Combustible. Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

1. Product Description

Trimec® Speed Lawn Weed Killer Ready Spray is a selective herbicide that provides post-emergent control of listed broadleaf weeds in established residential lawns. Trimec Speed Lawn Weed Killer Ready Spray may be applied to lawns containing Kentucky bluegrass, perennial ryegrass, fescues, common and hybrid bermudagrass, bahiagrass, bentgrass, buffalograss, centipedegrass, zoysiagrass, and St. Augustinegrass (except Floratam). Trimec Speed Lawn Weed Killer Ready Spray will control or suppress more than 250 broadleaf weeds, including dandelion, clover, henbit, oxalis, chickweed, knotweed, spurge, plantain, and many others as listed. Results will be visible within hours of application. Symptoms may include lead reddening, leaf and stem curling and twisting, and chlorosis.

2. Use Restrictions

- · For residential use only
- · Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until sprays have dried.
- · Do not apply when windy. Do not apply as a fine mist because of potential for injury to desirable plants. Do not use this product on or near desirable plants, including contact of spray on exposed root systems or adventitious shoots within the drip line of desirable trees and shrubs, since injury may result.
- To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day. Rinse applicator over lawn or garden area only.
- · Do not use on Carpetgrass, Dichondra, food crops (gardens, fruits, and vegetables), forage crops, or ornamental plants (flowers, trees, shrubs, hedges, woody ornamentals, groundcovers, groundcovers established in landscape plantings), 'Floratam' St. Augustinegrass, and turfgrass species that are not listed on this label.
- · Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day. Rinse applicator over lawn area only.

3. How Much To Apply

Over-application or rates above those specified on this label can cause turf injury.

For Kentucky Bluegrass, Bahiagrass, Common Bermudagrass, Fine Fescues, Tall Fescue, Perennial Ryegrass, and Zoysiagrass Entire Lawn Treatments: Apply 6 fl.oz. per 1000 sq.ft. This 32 fl.oz. container covers 5,300 sq.ft. Measure the size of your yard. If the area to be treated is larger than 5,300 sq.ft., you may need to buy more

For Hybrid Bermudagrass, Buffalograss, Centipedegrass, St. Augustinegrass (do not use on 'Floratam' St. Augustinegrass), and Bentgrass,

Entire Lawn Treatments: Apply 3 fl.oz. per 1000 sq.ft. This 32 fl.oz. container covers 10,600 sq.ft. Measure the size of your yard. If the area to be treated is larger than 10,600 sq.ft., you may need to buy more

The maximum rate is 6 fl.oz. of product per 1000 sq.ft per application and the maximum number of broadcast applications is 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 12 fl.oz. of product per 1000 sq.ft. excluding spot treatments.

4. When To Use

Spray when broadleaf weeds are young and actively growing for best results. This product may be applied in the spring, summer, or fall. Spring and fall treatments under adequate soil moisture conditions are preferred to the summer treatments. Generally, summer broadcast applications to older, drought stressed weeds are less effective.

Use broadcast (entire areas) applications or follow-up applications at 30 day intervals for more mature weeds, for dense infestations, and for adverse environmental conditions.

- · For Established Lawns: Apply when daytime temperatures are below 90°F.
- · For Newly Seeded Areas: Apply this product to newly seeded grasses when well-established (approximately 6 weeks after seedling emergence) or after the third mowing.
- For Newly Sodded, Sprigged or Plugged areas: Apply this product 3 to 4 weeks after the sodding, sprigging, or plugging operations.

- Reseeding: Treated areas can be reseeded at 2 weeks after the application of this product.
- Irrigation: Adequate soil moisture and favorable growing conditions enhance the performance of this product. If dry conditions exist, schedule irrigations (watering) before and after the application. Avoid applications of this product when turf is under stress since injury may result.
- Mowing: Avoid mowing within 1 to 2 days before and after the application of this product.

5. How To Apply

- Connect your garden hose to the hose-end sprayer. With your thumb, snap off the safety locking tab that prevents accidental discharge of spray solution. Leave control valve in the OFF position.
- 2. To BEGIN spraying: Turn on water at faucet. Extend hose to the furthest point from the faucet. Point nozzle toward the lawn to be treated and turn the control valve to the ON position. Water will automatically mix with the product. Walk backwards as you apply in order to avoid contact with the treated area. For best results, spray evenly over the measured area.
- To STOP spraying: Turn control valve to the OFF position. Turn off water at faucet. To relieve pressure, turn control valve to ON position and wait until water stops spraying before removing sprayer from hose.

6. Use Precautions

- Delay mowing 2 days before and until 2 days after the application of this product.
- If the lawn is dry, water 24 hours before using this product.
- · For best results, delay watering lawn until 2 hours after treatment.
- For new lawns, apply only when grass is well-established (approximately 6 weeks after seedling emergence) or after the third mowing.
- · Bare spots may be seeded 3 to 4 weeks after application.
- · Spray when air temperatures are cooler than 90°F.
- Avoid applications during the spring transition of the warm season grasses and during periods of extremely high temperatures.

7. Weeds Controlled

Amaranth, purple Hawkweed, orange Clover, rabbit-foot Clover, red Amaranth, slender Hawkweed, yellow Clover, strawberry Aster, heath Healall Heart-leaf drymary Aster, white heath Clover, yellow sweet Aster, purple Clover, white sweet Hemp Henbit Aster, white prairie Artichoke Clover, white Cockle Hoary-cress Horsenettle Austrian fieldcress Cocklebur Bedstraw, catch-weed Cocklebur, spiny Horseweed Bedstraw, smooth Common mullein Jimsonweed Bedstraw, northern Corn spurry Johnny jump-up violet Bedstraw, pineywoods Creeping beggar-weed Knawel Beggar-ticks, hairy Creeping Jenny Knotweed, prostrate Betony, Florida (creeping Charlie) Kochia Bindweed, field Cupid's shaving brush Lambsquarters Bindweed, hedge Cudweed Lespedeza Bitter Wintercress Daisy, English Mallow, alkali Bitter-cress, hairy Daisy, oxeye Mallow, bristly Dandelion, common Bitterweed Mallow, common Mallow, dwarf Mallow, Venice Black-eyed Susan Dandelion, false Black medic Day-flower Blood-flower milkweed Deadnettle Marcela Blue lettuce Dichondra Matchweed Brassbuttons Dock. broadleaf Mexican-weed Bristly oxtongue Dock, curly Moneywort Broomweed Dogbane Morningglory, bigroot Buckhorn Dog-fennel Morningglory, ivyleaf Morningglory, red Bull-nettle Dollarweed Bur-clover (Pennywort) Morningglory, tall Burdock Dove-weed Morningglory, wolly Burweed, lawn Elderberry Mugwort Mustard, ball Buttercup, bulbous Falseflax False sunflower Buttercup, creeping Mustard, black Buttonweed, common Fiddleneck Mustard, blue (poorjoe) Filaree, redstem Mustard, hedge Buttonweed, Virginia Carolina geranium Filaree, whitestem Mustard, Indian Fleabane, daisy Mustard, tansv Mustard, tumble (annual) Carpetweed Fleabane, rough Mustard, white Catnin Catsear, spotted (false Galinsoga, hairy Mustard, wormseed Galinsoga, no smallflower dandelion) Nettle, stinging Chamber bitter Nettle, tall Nettle, wood Chickweed, common Goldenrod Chickweed, mouse-ear Ground ivy Parsley-piert Chickweed, sticky Groundsel Parsnip Chicory Gumweed Pearlwort Cinquefoil Hairy fleabane Pennycress Clover, crimson Hawkweed, meadow Pepperweed, perennial Clover, Alyce Hawkweed, Clover, hop mouse-ear (cont. on next column)

7. Weeds Controlled (cont.)

Pepperweed, Virginia	Smartweed,	Trailing crown-vetch
(Peppergrass)	ladysthumb	Velvetleaf
Pigweed, prostrate	Smartweed, pale	Vervain, blue
Pigweed, redroot	Smartweed,	Vervain, hoary
Pigweed, Russian	Pennsylvania	Vervain, prostrate
Pigweed, smooth	Smooth chaff-flower	Vervain, tall
Pigweed, tumble	Smooth dock	Vetch, bird
Pineapple weed	Sorrel, red (sheep)	Vetch, hairy
Plains coreopsis	Sowthistle, annual	Vetch, milk
(tickseed)	Sowthistle, spiny	Vetch, narrow-leaf
Plantain, black-seed	Spanish needles	Vetch, wild
Plantain, bracted	Speedwell, bi-lobed	Virginia creeper
Plantain, broadleaf	Speedwell, common	Western clematis
Plantain, buckhorn	Speedwell, corn	Western salsify
Plantain, hoary	Speedwell, ivyleaf	Wild aster
Plantain, narrow-leaf	Speedwell, Persian	Wild buckwheat
Plantain, slender	Speedwell, purslane	Wild carrot
Plantain, woolly	Speedwell, slender	Wild four-o-clock
Poison ivy	Speedwell, snow	Wild garlic
Poison oak	Speedwell, thyme-leaf	Wild lettuce
Pokeweed	Spiny amaranth	Wild marigold
Prairie sunflower	Spurge, spotted	Wild mustard
Prickly lettuce	Spurge, prostrate	Wild onion
(compass plant)	Star of Bethlehem	Wild parsnip
Prickly sida	Strawberry,	Wild radish
Prostrate knotweed	Indian mock	Wild rape
Puncturevine	Tansy ragwort	Wild strawberry
Purslane, common	Tanweed	Wild sweet potato
Pusley, Brazilian	Thistle, blessed	Wild violet
Pusley, Florida	Thistle, bull	Woodsorrel, creeping
Pusley, large flower	Thistle, Canada	Woodsorrel, violet
Ragweed, bur	Thistle, Flodman's	Woodsorrel, yellow
Ragweed, common	Thistle, musk	(Oxalis)
Ragweed, lance-leaf	Thistle, plumeless	Woolly croton
Ragweed, western	Thistle, prairie	Wormseed
Redstem filaree	Thistle, Russian	Yarrow, common
Rough cinquefoil	Thistle, Scotch	Yarrow, western
Scarlet pimpernel	Thistle, wavy-leaf	Yarrow, yellow
Shepherd's purse	Thistle, yellow	Yellow Rocket

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in original container in a locked storage area inaccessible to children or pets. Keep from freezing.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTIAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

Gordon's® logo and Trimec® are registered trademarks of PBI-Gordon Corporation.

810/10-2022 AP022218 EPA REG. NO. 2217-1045



Employee-Owned

MANUFACTURED BY PBI/GORDON CORPORATION P.O. BOX 860350 SHAWNEE, KANSAS 66286 www.GordonsUSA.com

ATTENTION: This specimen label is provided for informational use only. This product may not yet be available for sale in your state or area. The information found in this label may differ from the information found on the product label you are using. Always follow the instructions for use and precautions on the label of the product you are using.