

ACTIVE INGREDIENTS:

2,4-D, 2-ethylhexyl ester	0.184%
Mecoprop-p acid	0.022%
Dicamba acid	
Carfentrazone-ethyl	0.007%
OTHER INGREDIENTS:	99.775%
TOTAL 1	100.000%

THIS PRODUCT CONTAINS:

0.010 lb 2,4-Dichlorophenoxyacetic acid equivalent 0.122%

0.002 lb (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid equivalent per gallon or 0.022%

0.001 lb 3,6-dichloro-o-anisic acid equivalent per gallon or 0.012%

0.001 lb Ethyl a,2-dichloro-5-[4(difluoromethyl)-4,5-dihydro-3-methyl-5-oxo-1H-1,2,4-triazol-1-yl]-4-fluorobenzenepropanoate per gallon or 0.007%

KEEP OUT OF REACH OF CHILDREN CAUTION



READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if absorbed through the skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

First Aid	
If on skin:	 Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment advice.

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

1. Product Description

Trimec[®] Speed Lawn Weed Killer Ready-To-Use is a selective herbicide that provides post-emergent control of listed broadleaf weeds in established residential lawns. Trimec Speed Lawn Weed Killer Ready-To-Use may be applied to lawns containing Kentucky bluegrass, perennial ryegrass, fescues, common and hybrid bermudagrass, bahiagrass, bentgrass, bettfalograss, centipedegrass, zoysiagrass, and St. Augustinegrass (except Floratam). Trimec Speed Lawn Weed Killer Ready-To-Use will control or suppress more than 250 broadleaf weeds, dandelion, clover, henbit, oxalis, chickweed, knotweed, spurge, plantain, and many others as listed. Results will be visible within hours of application. Symptoms may include leaf reddening, leaf and stem curling and twisting, and chlorosis.

2. Use Restrictions

- · For home lawns only.
- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until spray has dried.
- Do not apply as a fine mist because of potential for injury to desirable plants. Do not use this product on or near desirable plants, including contact of spray on exposed root systems or adventitious shoots within the drip line of desirable trees and shrubs, since injury may result.
- Do not use on Carpetgrass, Dichondra, food crops (gardens, fruits, and vegetables), forage crops, or ornamental plants (flowers, trees, shrubs, hedges, woody ornamentals, groundcovers, groundcovers established in landscape plantings), 'Floratam' St. Augustinegrass, and turfgrass species that are not listed on this label.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.

3. Where To Use

Trimec Speed Lawn Weed Killer Ready-To-Use may be used on residential lawns without harming lawn grasses when used as directed. Turfgrass species which may be treated include Kentucky Bluegrass, Perennial Ryegrass, Fine Fescues, Tall Fescue, Bentgrass, Bahiagrass, Buffalograss, Centipedegrass, Common and Hybrid Bermudagrasses, St. Augustinegrass (except 'Floratam' St. Augustinegrass), and Zoysiagrass.

4. When To Use

- Spray when weeds are actively growing.
- Spray when air temperatures are below 90°F and are not expected to exceed 90°F for 6 hours.
- · Hard to kill weeds may require a repeat application in 30 days.
- The number of treatments is limited to 2 per year.

5. How To Apply

- Remove wand from side clip.
 Pull connector by slightly twisting from holder and unwrap hose completely.
- 3. Loosen screws of battery cover.
- 4. Remove the insulation strip.
- 5. Tighten screws again.
- Remove cover from cap and insert plug end of hose into knob opening on cap until it clicks.
- 7. Flip open wand until it clicks and locks into position. Extend wand fully.
- 8. Slide trigger lock on wand handle to the unlocked position.
- 9. Twist nozzle toward STREAM until medium-coarse droplets are delivered.

Heavy Droplets Medium-Coarse Droplets

Fine mist (for comparison)







How to use the battery operated sprayer:

- Point nozzle away from body and hold trigger for continuous spray.
 Hold tip of wand approximately 12-18 inches away from the target weed and spray to wet.
- 3. Release the trigger to stop spray.

Important Use Information: Do not submerge in water. When storing sprayers for long periods, remove batteries.

6. Use Precautions

- Rain or watering 2 hours after application will not wash away effectiveness.
- Weeds in newly seeded lawns may be sprayed after the lawn has been mowed 3 times.
- Temporary discoloration may occur on St. Augustinegrass, centipedegrass, and bentgrass lawns.

7. Broadleaf Weeds Controlled

7. Broadleat Weeds	Controlled	
Amaranth, purple	Dog-fennel	Plains coreopsis
Amaranth, slender	Dollarweed (Pennywort)	(tickseed)
Aster, heath	Dove-weed	Plantain, black-seed
Aster, white heath	Elderberry	Plantain, bracted
Aster, purple	Falseflax	Plantain, broadleaf
Aster, white prairie	False sunflower	Plantain, buckhorn
Artichoke	Fiddleneck	Plantain, hoary
Austrian fieldcress	Filaree, redstem	Plantain, narrow-leaf
Bedstraw, catch-weed	Filaree, whitestem	Plantain, slender
Bedstraw, smooth	Fleabane, daisy (annual)	Plantain, woolly
Bedstraw, northern	Fleabane, rough	Poison ivy
Bedstraw, pineywoods	Galinsoga, hairy	Poison oak
Beggar-ticks, hairy	Galinsoga, smallflower	Pokeweed
Betony, Florida Bindweed, field	Goldenrod Ground ivy	Prairie sunflower
Bindweed, hedge	Groundsel	Prickly lettuce (compass plant)
Bitter Wintercress	Gumweed	Prickly sida
Bitter-cress, hairy	Hairy fleabane	Prostrate knotweed
Bitterweed	Hawkweed, meadow	Puncturevine
Black-eyed Susan	Hawkweed, mouse-ear	Purslane, common
Black medic	Hawkweed, orange	Pusley, Brazilian
Blood-flower milkweed	Hawkweed, yellow	Pusley, Florida
Blue lettuce	Healall	Pusley, large flower
Brassbuttons	Heart-leaf drymary	Ragweed, bur
Bristly oxtongue	Hemp	Ragweed, common
Broomweed	Henbit	Ragweed, lance-leaf
Buckhorn	Hoary-cress	Ragweed, western
Bull-nettle	Horsenettle	Redstem filaree
Bur-clover	Horseweed	Rough cinquefoil
Burdock	Jimsonweed	Scarlet pimpernel
Burweed, lawn	Johnny jump-up violet	Shepherd's purse
Buttercup, bulbous	Knawel	Smartweed, ladysthumb
Buttercup, creeping	Knotweed, prostrate	Smartweed, pale
Buttonweed, common	Kochia	Smartweed,
(poorjoe) Buttonweed, Virginia	Lambsquarters Lespedeza	Pennsylvania Smooth chaff-flower
Carolina geranium	Mallow, alkali	Smooth dock
Carpetweed	Mallow, bristly	Sorrel, red (sheep)
Catnip	Mallow, common	Sowthistle, annual
Catsear, spotted (false	Mallow, dwarf	Sowthistle, spiny
dandelion)	Mallow, Venice	Spanish needles
Chamber bitter	Marcela	Speedwell, bi-lobed
Chickweed, common	Matchweed	Speedwell, common
Chickweed, mouse-ear	Mexican-weed	Speedwell, corn
Chickweed, sticky	Moneywort	Speedwell, ivyleaf
Chicory	Morningglory, bigroot	Speedwell, slender
Cinquefoil	Morningglory, ivyleaf	Speedwell, snow
Clover, crimson	Morningglory, red	Speedwell, thyme-leaf
Clover, Alyce	Morningglory, tall	Spiny amaranth
Clover, hop	Morningglory, wolly	Speedwell, Persian
Clover, rabbit-foot	Mugwort	Speedwell, purslane
Clover, red Clover, strawberry	Mustard, ball Mustard, black	Spurge, spotted
Clover, yellow sweet	Mustard, Indian	Spurge, prostrate Star of Bethlehem
Clover, white sweet	Mustard, tansy	Strawberry, Indian mock
Clover, white	Mustard, tumble	Tansy ragwort
Cockle	Mustard, white	Tanweed
Cocklebur	Mustard, blue	Thistle, blessed
Cocklebur, spiny	Mustard, hedge	Thistle, bull
Common mullein	Mustard, wormseed	Thistle, Canada
Corn spurry	Nettle, stinging	Thistle, Flodman's
Creeping beggar-weed	Nettle, tall	Thistle, musk
Creeping Jenny (creeping	Nettle, wood	Thistle, plumeless
Charlie)	Parsley-piert	Thistle, prairie
Cupid's shaving brush	Parsnip	Thistle, Russian
Cudweed	Pearlwort	Thistle, Scotch
Daisy, English	Pennycress Bopperwood perophial	Thistle, wavy-leaf
Daisy, oxeye	Pepperweed, perennial Pepperweed, Virginia	Thistle, yellow
Dandelion, common Dandelion, false	(Peppergrass)	Trailing crown-vetch Velvetleaf
Day-flower	Pigweed, prostrate	Vervain, blue
Deadnettle	Pigweed, redroot	Vervain, hoary
Dichondra	Pigweed, Russian	Vervain, prostrate
Dock, broadleaf	Pigweed, smooth	Vervain, tall
Dock, curly	Pigweed, tumble	Vetch, bird
Dogbane	Pineapple weed	(cont. on next column)
· · · ·		,

Vetch, hairy	Wild garlic	Woodsorrel, creeping
Vetch, milk	Wild lettuce	Woodsorrel, violet
Vetch, narrow-leaf	Wild marigold	Woodsorrel, yellow
Vetch, wild	Wild mustard	(Oxalis)
Virginia creeper	Wild onion	Woolly croton
Western clematis	Wild parsnip	Wormseed
Western salsify	Wild radish	Yarrow, common
Wild aster	Wild rape	Yarrow, western
Wild buckwheat	Wild strawberry	Yarrow, yellow
Wild carrot	Wild sweet potato	Yellow Rocket
Wild four-o-clock	Wild violet	

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Keep from freezing. Keep pesticide in original container. Do not put into food or drink containers. Avoid contamination of feed and foodstuffs. Store in a cool, dry place, preferably in a locked storage area.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call you local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY STATEMENT cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

Gordon's $^{\!\!\!\circ}$ logo and Trimec $^{\!\!\!\circ}$ are registered trademarks of PBI-Gordon Corporation.

885/10-2022 AP040320 EPA REG. NO. 2217-1046



MANUFACTURED BY PBI/GORDON CORPORATION P.O. BOX 860350 SHAWNEE, KANSAS 66286 www.GordonsUSA.com

ATTENTION: This specimen label is provided for informational use only. This product may not yet be available for sale in your state or area. The information found in this label may differ from the information found on the product label you are using. Always follow the instructions for use and precautions on the label of the product you are using.