

# CAUTION

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

## Trio Tebuconazole 430 SC Fungicide

ACTIVE CONSTITUENT:  
430 g/L TEBUCONAZOLE

GROUP	<b>3</b>	FUNGICIDE
-------	----------	-----------

For the control of leaf spot and leaf speckle on bananas; rust, leaf spot and net blotch of peanuts; foliar diseases on cereals crops; and other diseases on beans, peas, onions, pawpaw, pyrethrum and ryegrass and fescue seed crops as specified in the Directions for Use.

**IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE**

**Contents: 5L, 10L, 20L, 110L, 200L, 1000L**

CTS Chemicals Pty Ltd. ABN 22 605 759 644  
Manning Buildings, 135 High Street Mall  
Fremantle WA 6160 Australia  
Phone 1800 749 140

## **STORAGE AND DISPOSAL**

Keep out of reach of children.

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.

Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

### **Refillable Containers:**

Empty contents fully into application equipment. Close all valves and return to point of supply/designated collection point for refill or storage.

## **SAFETY DIRECTIONS**

Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container, preparing the spray and using prepared spray wear cotton overalls buttoned to the neck and wrist, washable hat, elbow-length PVC gloves. After use, and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and contaminated clothing.

## **FIRST AID INSTRUCTIONS**

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766.

## **SAFETY DATA SHEET**

Additional information is listed in the Safety Data Sheet that can be obtained from the supplier.

## **CONDITIONS OF SALE**

CTS Chemicals Pty. Ltd. makes no warranty or guarantee and will not accept any responsibilities whatsoever and howsoever arising and whether for consequential, special or indirect loss or otherwise in connection with the supply, storage or use of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatory imposed by Statutes applicable to the sale or supply of these goods. Always read the product labels on the container for all instructions, recommendations, critical comments and conditions of sale.

APVMA Approval No: 89560 / 125267

Batch No:

Date of Manufacture:

# CAUTION

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

## Trio Tebuconazole 430 SC Fungicide

ACTIVE CONSTITUENT:  
430 g/L TEBUCONAZOLE

GROUP	<b>3</b>	FUNGICIDE
-------	----------	-----------

**For the control of leaf spot and leaf speckle on bananas; rust, leaf spot and net blotch of peanuts; foliar diseases on cereals crops; and other diseases on beans, peas, onions, pawpaw, pyrethrum and ryegrass and fescue seed crops as specified in the Directions for Use.**

**IMPORTANT: READ THIS LEAFLET BEFORE USE**

APVMA Approval No: 89560 / 125267

CTS Chemicals Pty Ltd. ABN 22 605 759 644  
Manning Buildings, 135 High Street Mall  
Fremantle WA 6160 Australia  
Phone 1800 749 140

## DIRECTIONS FOR USE

Crop	Disease	Application	Rate mL/ha	WHP	Critical Comments
Bananas (Qld, NSW, WA, NT only)	Leaf spot (Yellow sigatoka) Leaf speckle Black Sigatoka	<b>Ground Application</b> Apply by misting machine or airblast sprayer in a convenient volume of water <b>Aerial application</b> Apply at least 20L of spray mixture per hectare	230  Add 3-6 L water miscible oil per ha	H 1 day	Maintain good de-leafing practices to reduce disease inoculum. Very old leaves and leaves with advanced lesions should be removed, or infected parts of the leaf removed, prior to the application of Trio Tebuconazole 430 SC Fungicide. <b>Tropical Areas (e.g. North Qld, NT, Ord River):</b> Apply a regular schedule of protectant sprays. When conditions favour disease, apply a minimum of 2 and a maximum of 3 consecutive Trio Tebuconazole 430 SC Fungicide sprays at 14 day intervals. Do not apply more than 6 Trio Tebuconazole 430 SC Fungicide sprays in any 12 month period. Do not apply any Trio Tebuconazole 430 SC Fungicide sprays in the months of July, August and September. <b>Sub-Tropical Areas (e.g. South Qld, NSW):</b> Commence spraying with Trio Tebuconazole 430 SC Fungicide at the onset of warm and humid/wet weather, normally December. Repeat at 21 to 28 day intervals using a maximum of 2 consecutive Trio Tebuconazole 430 SC Fungicide sprays. Do not apply more than 5 Trio Tebuconazole 430 SC Fungicide sprays in any 12 month period. Trio Tebuconazole 430 SC Fungicide is approved for use in banana plantations inter-planted with avocados
Peanuts (South Qld and NSW only)	Early Leaf spot Late leaf spot Rust Net Blotch	<b>Low Disease Ground Application</b> Apply in at least 100L of water per ha <b>Aerial application</b> Apply at least 30L of spray mixture per hectare	175 + Agridex 1L/ha	H 3 weeks G 3 weeks	Regularly check high-risk areas in the crop for disease (e.g. lower leaves, shaded plants). When leaf spot or net blotch or rust can easily be found, then either; <ul style="list-style-type: none"> <li>Spray immediately after last disease conducive weather (e.g. rain or heavy dews)</li> </ul> or <ul style="list-style-type: none"> <li>Spray before the next conducive weather.</li> </ul> Repeat after 14 days if conditions remain favourable to disease development. If not, repeat just before or just after the next disease conducive weather. For resistance strategy, see General Instructions.
		<b>High disease OR Wet weather</b>	290 + Agridex 1L/ha		
		<b>High disease AND Wet weather</b>	440 + Agridex 1L/ha		

Peanuts (North Qld, WA, NT only)	Early Leaf Spot Late leaf spot Rust	<b>Low Disease</b>	230 + Agridex 1L/ha		Begin spraying at 3-4 weeks after planting. If band spraying, apply half the appropriate rate as a 45cm band directly over the row. Sprays should not be banded after 6 weeks from planting. Repeat applications at 14-day intervals. If prolonged wet weather or heavy rains occur, shorten spray interval to 10-12 days. For resistance strategy, see General Instructions.	
		<b>Moderate disease</b>	290 + Agridex 1L/ha			
		<b>Severe disease</b>	440 + Agridex 1L/ha			
	Net blotch	<b>Low Disease</b>	290 + Agridex 1L/ha			Apply at 14 day intervals. If prolonged cool moist weather occurs, shorten spray interval to 10-12 days. For resistance strategy, see General Instructions.
	<b>Moderate - severe disease</b>	440 + Agridex 1L/ha				
Green Beans	Rust	Can be applied by aircraft or ground rig	350 + Agridex 1L/ha	H 3 days G 3 days	Spray when rust infection begins or at budding, whichever is the earlier. Repeat application 10-14 days later. A third application may be necessary when infection occurs early or disease pressure is high	
Peas	Powdery Mildew	<b>Ground Application</b> Apply in at least 50L of water per ha	145		Apply at flowering or at the first sign of disease, whichever occurs first. A second spray 14 days later may be necessary under some conditions	
Wheat	Leaf rust Stripe rust Septoria nodorum blotch Yellow leaf spot	<b>Aerial application</b> Apply in at least 10L of water per hectare	145 or 290	H 5 weeks G 14 days	Use the higher rate when longer disease control is required. <b>Stripe rust:</b> See spray timings under General Instructions.  Other diseases: Apply from full flag leaf emergence to early head emergence. The addition of mineral crop oil (e.g. DC Trate or equivalent) at 1% may improve performance of Trio Tebuconazole 430 SC Fungicide on wheat, oats and barley.	
	Septoria tritici blotch		290			
Oats	Crown rust		145 or 290			
Barley	Scald		145			Apply at late tillering to early jointing.
	Powdery mildew		145 or 290			Apply when 5% of the leaf area is infected. Use higher rate when longer disease control is required.
Wheat Oats	Stem rust					Apply if more than 5% of stems become infected between full flag emergence to late flowering. When stem rust is the major disease, yield responses are usually optimised by delaying application until full head emergence, and using the higher rate. In severe cases, if a majority of stems are infected prior to full head emergence, apply at 145 mL/ha as soon as possible and if necessary, repeat after 3 weeks when heads are fully emerged.

Onions (Tas only)	White root rot	Before sowing apply Trio Tebuconazole 430 SC Fungicide onto lime super. Ensure good coverage of all lime super particles.	1.45 mL/100 m of row mixed with 145-218 g lime super/100 m of row	-	Apply Trio Tebuconazole 430 SC Fungicide treated lime super when sowing onion seed. Seed and lime super can either be mixed in the same box on the drill or placed in different boxes and sown down the same tube. Apply in a bandwidth of 2cm. Ensure that the correct rate of Trio Tebuconazole 430 SC Fungicide is used otherwise some delay in emergence and reduced stands of seedlings may occur.
Pawpaw	Black spot	Ensure thorough coverage of leaves and fruit.	290	H 3 days	Ensure infected plant material is regularly removed and destroyed to reduce inoculum levels. Spray equipment must be properly calibrated to apply the correct amount of Trio Tebuconazole 430 SC Fungicide. Apply Trio Tebuconazole 430 SC Fungicide at 14 day intervals. Alternate Trio Tebuconazole 430 SC Fungicide with sprays of a protectant fungicide (e.g. mancozeb). Do not apply more than 6 sprays of Trio Tebuconazole 430 SC Fungicide (or any DMI fungicide) on any block in any 12 month period.
Pyrethrum	<i>Sclerotinia sclerotiorum</i>		350	-	Apply twice, in rotation with other control measures, at 7-10 day intervals. Commence at 1 to 2% flowering. Use under direction of pyrethrum advisers.
Ryegrass and fescue seed crops	Leaf rust Stem rust	Apply in at least 100L of water per ha	290	G 14 days	Monitor crops closely and spray at the first signs of disease. Continuing disease pressure or re-infection may require a further application 3 – 4 weeks later. Ensure thorough coverage and use higher water volumes in dense or advanced crops.

**NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**WITHHOLDING PERIODS:**

**BANANAS, AVOCADOS: DO NOT HARVEST FOR 1 DAY AFTER APPLICATION.**

**BEANS, PEAS: DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION AND DO NOT GRAZE OR CUT FOR LIVESTOCK FOR 3 DAYS AFTER APPLICATION.**

**CEREALS: DO NOT HARVEST FOR 5 WEEKS AFTER APPLICATION AND DO NOT GRAZE OR CUT FOR LIVESTOCK FOR 14 DAYS AFTER APPLICATION.**

**ONIONS: NOT REQUIRED WHEN USED AS DIRECTED.**

**PAWPAW: DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION.**

**PEANUTS: DO NOT HARVEST FOR 21 DAYS AFTER APPLICATION AND DO NOT GRAZE OR CUT FOR LIVESTOCK FOR 21 DAYS AFTER APPLICATION.**

**RYE GRASS AND FESCUE SEED CROPS: DO NOT GRAZE OR CUT FOR STOCKFOOD FOR 14 DAYS AFTER APPLICATION.**

**H = Harvest, G = Grazing**

## GENERAL INSTRUCTIONS

### RESISTANCE MANAGEMENT RECOMMENDATIONS – PEANUTS

Apply no more than 3 consecutive sprays of DMI fungicides (e.g. tebuconazole) before switching to a non-DMI fungicide. Apply no more than 5 DMI sprays per season.

### SPECIAL WARNING – BANANAS

There are certain conditions where the surface of recently emerged fruit is particularly prone to marking damage from spray applications. In circumstances where application will be made to very rapidly growing fruit in hot conditions with strong direct light, it is recommended that all emerged bunches be bagged prior to spraying to minimise the risk of fruit marking.

**Never include adjuvants** other than water miscible oils with Trio Tebuconazole 430 SC Fungicide sprays. The inclusion of wetting agents is known to cause phototoxicity to young fruit.

### FOLIAR DISEASES ON CEREAL CROPS

Do not apply to cereal crops more than once in a season. Treatment will give approximately three weeks disease suppression. Economic response may not be gained by spraying crops past flowering stage. The effects of fungicide application will not be clearly seen for 7-10 days after application. Yield potential: crops with potential yield under 2 t/ha are unlikely to give economic responses to a fungicide spray except under conditions of very severe disease. Economic responses are most likely with crops with potential yield of over 3 t/ha.

#### Spray Timings for Stripe Rust Control

Obtain advisory literature from Department of Agriculture for classification of resistant and susceptible varieties.

The following spray program is suggested:

**Seedling infections:** When approximately 20 out of 100 leaves show signs of infection during tillering or jointing, apply a spray within a week.

#### Adult infections

**Susceptible varieties:** When approximately 10 out of 100 leaves show first sign of infection apply a spray within a week.

**Moderately Susceptible varieties:** When approximately 15-20 out of 100 leaves show first sign of infection apply a spray within a week. Do not delay.

**Moderately Resistant and Resistant Varieties:** Monitor carefully. If rust appears and spreads, spray within 1 week.

### MIXING

Prior to pouring, shake container vigorously, then add the required amount of Trio Tebuconazole 430 SC Fungicide to water in the spray vat while stirring or with agitators in motion. Add the required amount of Agridex (peanuts, beans) or water miscible oil (bananas) and mix thoroughly.

### APPLICATION

Aircraft should fly as low as possible under the prevailing conditions to minimise drift.

### FUNGICIDE RESISTANCE WARNING

GROUP	3	FUNGICIDE
-------	---	-----------

Trio Tebuconazole 430 SC Fungicide is a member of the DMI group of fungicides. For fungicide resistance management the product is a Group 3 Fungicide.

Some naturally occurring individual fungi resistant to Trio Tebuconazole 430 SC Fungicide and other Group 3 Fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by Trio Tebuconazole 430 SC Fungicide or other Group 3 Fungicides, thus resulting in a reduction in efficacy and possible yield loss.

Since the occurrence of resistant fungi is difficult to detect prior to use, CTS Chemicals Pty Ltd accepts no liability for any losses that may result from the failure of Trio Tebuconazole 430 SC Fungicide to control resistant fungi.

## **PRECAUTIONS**

### **Re-Entry Period**

Do not allow entry into treated areas until the spray has dried.

When prior entry is necessary, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), chemical resistant gloves and footwear. Clothing must be laundered after each day's use.

## **PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

DO NOT contaminate streams, rivers or waterways with chemical or used containers. A spray drift minimisation strategy should be employed at all times when aerially applying sprays. The strategy envisaged is exemplified by the cotton industry's Best Management Practise Manual.

## **STORAGE AND DISPOSAL**

Keep out of reach of children.

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.

Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

### **Refillable Containers:**

Empty contents fully into application equipment. Close all valves and return to point of supply/designated collection point for refill or storage.

## **SAFETY DIRECTIONS**

Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container, preparing the spray and using prepared spray wear cotton overalls buttoned to the neck and wrist, washable hat, elbow-length PVC gloves. After use, and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and contaminated clothing.

## **FIRST AID INSTRUCTIONS**

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766.

## **SAFETY DATA SHEET**

Additional information is listed in the Safety Data Sheet that can be obtained from the supplier.

## **CONDITIONS OF SALE**

CTS Chemicals Pty. Ltd. makes no warranty or guarantee and will not accept any responsibilities whatsoever and howsoever arising and whether for consequential, special or indirect loss or otherwise in



connection with the supply, storage or use of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatory imposed by Statutes applicable to the sale or supply of these goods. Always read the product labels on the container for all instructions, recommendations, critical comments and conditions of sale.

NOTE: Add the following to the back and top label, not the booklet.

GHS STATEMENTS:

Hazard Statements:

**Harmful if swallowed. Suspected of damaging fertility or the unborn child.**

Precautionary Statements - Prevention

Obtain special instructions before use. Do not handle until all safety precautions have been read and understood. Do not eat, drink or smoke when using this product.


Precautionary Statements - Response

IF exposed or concerned: Get medical advice/attention. IF SWALLOWED: Call a POISON CENTER or doctor if you feel unwell. Rinse mouth.

Precautionary Statements - Storage

Store locked up.

DG Classification:



UN3082

ENVIRONMENTALLY HAZARDOUS SUBSTANCES, LIQUID, N.O.S

(Tebuconazole)

In a transport emergency Dial 000 Police or Fire Brigade