



VANGO™

WG



Contains cyprodinil, the active ingredient used in Vanguard® WG Fungicide.

ACTIVE INGREDIENT:	(% by weight)
Cyprodinil: 4-cyclopropyl-6-methyl-N-phenyl-pyrimidinamine*	75.0%
OTHER INGREDIENTS:	25.0%
TOTAL:	100.0%

Vango™ WG is a water-dispersible granule containing 75% cyprodinil.

*CAS No. 121552-61-2

EPA Reg. No.: 91234-78

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

SEE BELOW FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

FIRST AID	
If in eyes:	<ul style="list-style-type: none"> ▪ Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. ▪ Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. ▪ Call a poison control center or doctor for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none"> ▪ Take off contaminated clothing. ▪ Rinse skin immediately with plenty of water for 15 - 20 minutes. ▪ Call a poison control center or doctor for treatment advice.
NOTE TO PHYSICIAN: If ingested, induce emesis or lavage stomach. Treat symptomatically.	
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall at 1-844-685-9173 for emergency medical treatment information.	

**For Chemical Emergency: Spill, Leak, Fire, Exposure, or Accident,
Call CHEMTREC Day or Night Within USA and Canada:
1-800-424-9300 or +1 703-527-3887 (collect calls accepted)**

Vango™ WG is not manufactured, or distributed by Syngenta Crop Protection, seller of Vanguard® WG Fungicide.



Manufactured for:
Atticus, LLC
940 NW Cary Parkway, Suite 200
Cary, NC 27513

PRECAUTIONARY STATEMENTS
Hazards to Humans and Domestic Animals
CAUTION

Causes moderate eye irritation. Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment (PPE):

Some materials that are chemical-resistant to this product are listed below.

Handlers applying this product as a pre-plant dip to strawberry roots and crowns and workers packaging or preparing treated roots and crowns for shipment must wear:

- Chemical-resistant apron made of any waterproof material
- Elbow-length chemical-resistant gloves made of any waterproof material
- Chemical-resistant boots made of any waterproof material

All other applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof materials (polyethylene or polyvinyl chloride)
- Shoes plus socks

In addition, mixers and loaders for aerial and groundboom applications must wear:

- A minimum of a NIOSH-approved particulate filtering facepiece respirator with any N, R, or P filter; OR a NIOSH-approved elastomeric particulate respirator with any N1, R, or P filter; OR a NIOSH-approved powered air purifying respirator with HE filters.

USER SAFETY REQUIREMENTS

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statements:

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards:

This pesticide is toxic to fish and aquatic invertebrates. For terrestrial uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwater or rinsate.

Surface and Ground Water Advisory:

This chemical may contaminate water through runoff. This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This chemical has a potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this chemical. A level, well-maintained vegetative buffer strip between areas to which this chemical is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of cyprodinil from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours. Sound erosion control practices will reduce this product's potential to reach aquatic sediment via runoff.

Physical or Chemical Hazards:

Do not use, pour, spill or store near heat or open flame.

Do not mix or allow coming into contact with oxidizing agents, hazardous chemical reaction may occur.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

PRODUCT INFORMATION

Vango WG is a broad spectrum fungicide which controls certain diseases in fruits, herbs, nuts and vegetables. Failure to follow directions and precautions on this label may result in crop injury, poor disease control, and/or illegal residues.

APPLICATION INFORMATION

Vango WG has preventative and systemic properties and is labeled for the control of many important plant diseases. **Vango WG** provides excellent disease control of Botrytis, several leaf spots and powdery mildews. **Vango WG** is applied as a foliar spray and can be used in block, alternating spray, or tank-mix programs with other crop protection products. Make all applications according to the use directions that follow.

When using **Vango WG** in tank mixtures, add all products in water-soluble packaging to the tank before any other tank-mix partner, including **Vango WG**. Allow the water-soluble packaging to completely dissolve and the product(s) to completely disperse before adding any other tank-mix partner to the tank.

If using **Vango WG** in a tank mixture, it is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Tank mixtures or other applications of products referenced on this label are permitted only in those states in which the referenced products are registered.

Additives:

Vango WG is compatible with most crop protection additives. Do not use X-77® with **Vango WG** for bloom sprays applied to stone fruits, almonds and pistachios.

APPLICATION

For best disease control, apply **Vango WG** in sufficient water to provide thorough and uniform coverage. Use minimum ground spray volumes of 50 gal/A for tree crops, 30 gal/A for vine crops, and 15 gal/A for field and vegetable crops. For aerial application, see spray volume requirements in the specific crop directions for use.

For ground applications, do not apply when wind speeds exceed 15 miles per hour at the application site.

To prevent spray drift, do not apply when conditions favor drift beyond the target area. Spray overlap may cause crop injury.

For air assisted or air blast sprayers, move spray droplets into the canopy using a forced air stream. Set up the fan to deliver only enough air volume to penetrate the canopy and provide good coverage. Adjust deflectors or other aiming devices to direct spray only to the target area.

Equip sprayers with nozzles that provide accurate and uniform application. Check whirl plates and nozzle discs for wear and replace as necessary. Calibrate sprayer before use. Use a pump with capacity to maintain the correct rated pressure for the nozzles selected. Maintain sufficient agitation to keep the spray mixture in suspension. Use a jet agitator, liquid sparge tube, or mechanical paddle for agitation. Do not air sparge. Use screens to prevent nozzles from clogging. Use 50-mesh or coarser screens placed after the tank and before the nozzles. Check nozzle manufacturer's recommendations.

For more information on spray equipment and calibration, consult sprayer manufacturers and state recommendations. For specific local directions and spray schedules, consult the current state agricultural experiment station recommendations.

OBSERVE THE FOLLOWING RESTRICTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES, RESERVOIRS, RIVERS, PERMANENT STREAMS, MARSHES OR NATURAL PONDS, ESTUARIES, AND COMMERCIAL FISH PONDS.

- Do not apply within 75 ft of bodies of water such as lakes, reservoirs, rivers, permanent streams, natural ponds, marshes or estuaries.
- Shut off the sprayer when row ends.
- Do not cultivate within 10 ft of aquatic areas in order to allow a vegetative filter strip.
- Do not apply when weather conditions favor drift to aquatic areas. Do not apply when gusts or sustained winds exceed 10 mph.
- Do not apply during a temperature inversion. Mist or fog may indicate the presence of an inversion in humid areas.
- For perennial crops such as tree crops and grapes:
 - For all plantings within 150 ft of bodies of water as described above, spray crops from outside the planting away from the bodies of water.
 - Spray last three rows windward of aquatic areas using nozzles on one side only, with spray directed away from aquatic areas. Adjust or turn off top nozzles to prevent spray going over the tops of trees. Shut off nozzles on the side away from the grove/orchard when spraying the outside row. Shut off nozzles when turning at ends of row or passing tree gaps in the rows.

AERIAL SPRAY DIRECTIONS

Avoid applications under conditions when uniform coverage cannot be obtained or when excessive drift may occur.

Aerial Spray Restrictions:

Observe the following restrictions when spraying in the vicinity of aquatic areas such as lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial fish ponds.

- Use only on crops where aerial applications are indicated.
- Do not apply by air within 150 ft of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial fish ponds.
- Mount the spray boom on the aircraft so as to minimize the drift caused by wing tip vortices.
- Release spray at the lowest height consistent with pest control and flight safety. Do not make applications more than 10 feet above the crop canopy.
- Do not apply when weather conditions favor drift to aquatic areas.
- Do not apply during a temperature inversion. Mist or fog may indicate the presence of an inversion in humid areas.
- For aerial applications: Do not apply when wind speeds exceed 15 mph at the application site. If the windspeed is greater than 10 mph, the boom length must be 65% or less of the wingspan for fixed wing aircraft and 75% or less of the rotor diameter for helicopters. Otherwise, the boom length must be 75% or less of the wingspan for fixed-wing aircraft and 90% or less of the rotor diameter for helicopters.

Aerial Spray Precautions:

Observe the following precautions when spraying in the vicinity of aquatic areas such as lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial fish ponds.

- Use the largest droplet size consistent with good pest control.
- Formation of very small droplets may be minimized by appropriate nozzle selection, by orientating nozzles away from the air stream as much as possible, and by avoiding excessive spray boom pressure.
- Reduce risk of exposure to aquatic areas by avoiding applications when wind direction is toward the aquatic area.
- Low humidity and high temperatures increase the evaporation rate of spray droplets, and therefore the likelihood of increased spray drift to aquatic area. Avoid spraying during conditions of low humidity and/or high temperatures.

Application Through Irrigation Systems (Chemigation):

- Use only on crops for which chemigation is specified on this label.
- Apply this product only through center pivot, solid set, hand move, or moving wheel irrigation systems. Do not apply this product through any other type of irrigation system.
- Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.
- Apply in 0.125 - 0.25 inches/A of water. Excessive water may reduce efficacy.
- If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers, or other experts.
- Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system, unless the pesticide label-prescribed safety devices for public water systems are in place.
- A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Note: Do not inject **Vango WG** at full strength or deterioration of valves and seals may occur. Use a dilution ratio of at least 10 parts water to 1 part **Vango WG**. **Vango WG** is corrosive to many seal materials. Leather seals are best. EPDM or silicone rubber seals can be used but should be replaced once a year. Do not use Viton®, Buna-N, Neoprene, or PVC seals.

Operating Instructions:

1. The system must contain a functional check-valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water-source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check-valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended.

Center Pivot Irrigation Equipment:

- Determine the size of the area to be treated.
- Determine the time required to apply 1/8 - 1/2 inch of water over the area to be treated when the system and injection equipment are operated at normal pressures as recommended by the equipment manufacturer. When applying **Vango WG** through irrigation equipment use the lowest obtainable water volume while maintaining uniform distribution. Run the system at 80 - 95% of the manufacturer's rated capacity.
- Using water, determine the injection pump output when operated at normal line pressure.
- Determine the amount of **Vango WG** required to treat the area covered by the irrigation system.
- Add the required amount of **Vango WG** and sufficient water to meet the injection time requirements to the solution tank.
- Make sure the system is fully charged with water before starting injection of the **Vango WG** solution. Time the injection to last at least as long as it takes to bring the system to full pressure.
- Maintain constant solution tank agitation during the injection period.
- Continue to operate the system until the **Vango WG** solution has cleared the sprinkler head.

Restrictions:

1. Use only with drive systems which provide uniform water distribution.
2. Do not use end guns when chemigating **Vango WG** through center pivot systems because of non-uniform application.

Solid Set, Hand Move, and Moving Wheel Irrigation Equipment:

- Determine the acreage covered by the sprinklers.
- Fill injector solution tank with water and adjust flow rate to use the contents over a 20 to 30-minute interval. When applying **Vango WG** through irrigation equipment use the lowest obtainable water volume while maintaining uniform distribution.
- Determine the amount of **Vango WG** required to treat the area covered by the irrigation system.
- Add the required amount of **Vango WG** into the same quantity of water used to calibrate the injection period.
- Operate the system at the same pressure and time interval established during the calibration.
- Stop injection equipment after treatment is completed. Continue to operate the system until the **Vango WG** solution has cleared the last sprinkler head.

SPECIFIC INSTRUCTIONS FOR PUBLIC WATER SYSTEMS

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, back-flow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
4. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.

CROP USE DIRECTIONS

TREE, NUTS AND VINES

Crop	Disease	Product Rate oz/Acre	Remarks
Almonds	Brown Rot Blossom Blight (<i>Monilinia</i> spp.) Suppression: Green Fruit Rot (Jacket Rot) (<i>Botrytis cinerea</i>) Shot Hole (<i>Wilsonomyces carpophilus</i>)	5 - 10	Apply Vango WG at 5 - 10% bloom. Additional applications at 50 - 100% bloom and petal fall may be necessary. When used for control of brown rot blossom blight, Vango WG will provide suppression of shot hole. For broad spectrum disease control in tank mixture, apply Vango WG at a minimum rate of 5 oz in tank mixtures with other fungicides registered for use on almonds. For suppression of green fruit rot, apply Vango WG at full bloom. Disease suppression for almond diseases refers to erratic control from fair to good, or consistent control at a level below that obtained with products registered for control.

Application Instructions: Application may be made by ground or air. Good coverage is essential for good disease control. Use a minimum of 20 gallons/A spray volume by air. Make no more than two applications by air.

Specific Use Restrictions:

1. Make no more than two applications by air.
2. Do not apply more than 30 oz/A of **Vango WG** (1.4 lb ai/A of cyprodinil) per year.
3. Do not apply within 60 days of harvest (60-day PHI).

Crop	Disease	Product Rate oz/Acre	Remarks
Citrus* Lemon Lime	Alternaria Stem End Rot (<i>A. citri</i>) Anthracnose (<i>Colletotrichum gloeosporioides</i>) Blue Mold (<i>Penicillium italicum</i>) Green Mold (<i>Penicillium digitatum</i>)	5.5 - 7	Make one application near harvest to prevent post-harvest fruit rot. The application may be made up to and including the day of harvest.

Application Instructions: Good coverage is essential for good disease control.

Specific Use Restrictions:

1. Application may be made by ground only.
2. Do not apply more than 7 oz/A of **Vango WG** per year.
3. Do not apply more than 0.33 lb ai/A of cyprodinil-containing products per year.
4. May be applied on the day of harvest (0-day PHI).
5. Do not exceed one application per year.

*Not registered for use in California.

Crop	Disease	Product Rate oz/Acre	Remarks
Small Fruit Vine Climbing Subgroup 13-07F (except fuzzy kiwifruit) Grapes Amur River Grape Hardy Kiwifruit Maypop Schisandra Berry And cultivars and/or hybrids of these	Botrytis Bunch Rot <i>(Botrytis spp.)</i> Suppression: Powdery Mildew <i>(Uncinula necator)</i>	Vango WG Alone 10	Begin applications of Vango WG at early bloom. Make an additional application at berry touch, veraison, or preharvest using at least a 7-day spray interval. Botrytis bunch rot is most effectively controlled by ground application, using sufficient water volume to provide thorough coverage. Thorough coverage of bunches is essential. When used at 10 oz/A, Vango WG will provide significant suppression (approximately 60% control) of powdery mildew.
		Vango WG Tank Mixtures 5 - 10	Apply Vango WG in tank mixture with the label rate of another fungicide registered on grapes for control of Botrytis bunch rot.

Application Instructions: Application may be made by ground or air. Good coverage is essential for good disease control. Use a minimum of 20 gallons/A spray volume by air. Make no more than two applications by air.

Specific Use Restrictions:

1. Make no more than two applications by air.
2. Do not apply more than 30 oz/A of **Vango WG** (1.4 lb ai/A of cyprodinil) per year.
3. Do not apply within 7 days of harvest (7-day PHI).

Crop	Disease	Product Rate oz/Acre	Remarks
Kiwi	Botrytis Fruit Rot <i>(Botrytis spp.)</i>	10	Make 1 - 2 applications on a 7 - 10 day interval prior to harvest to prevent post-harvest fruit rot.

Application Instructions: Application may be made by ground or air. Good coverage is essential for good disease control. Use a minimum of 20 gallons/A spray volume by air.

Specific Use Restrictions:

1. Make no more than two applications by air.
2. Do not apply more than 20 oz/A of **Vango WG** (0.94 lb ai/A of cyprodinil) per year.
3. May be applied on the day of harvest (0-day PHI).

Crop	Disease	Product Rate oz/Acre	Remarks
Pistachios	Alternaria <i>(Alternaria alternata)</i> Botrytis <i>(Botrytis spp.)</i>	5.5 - 7	Make the first application during early bloom and repeat applications at 14-day intervals if conditions remain favorable for disease development.

Application Instructions: Application may be made by ground or air. Good coverage is essential for good disease control. Use a minimum of 20 gallons/A spray volume by air. Make no more than two applications by air.

Specific Use Restrictions:

1. Make no more than two applications by air.
2. Do not apply more than 28 oz/A of **Vango WG** (1.3 lb ai/A of cyprodinil) per year.
3. Do not apply within 7 days of harvest (7-day PHI).

Crop	Disease	Product Rate oz/Acre	Remarks
Pome Fruits¹ Apples Crabapples Loquat Mayhaw Pears (see tank mixture instructions) Quince And cultivars and/or hybrids of these	Prebloom Scab <i>(Venturia spp.)</i>	Vango WG Alone 5	For pome fruits except pear, begin application at green tip and continue on a 7 - 10 day interval. Under severe disease pressure, use the shorter interval.
		Vango WG Tank Mixtures 3 - 5	For scab control utilizing multiple modes of action, apply Vango WG in tank mixture with the label rate of a protectant or systemic fungicide registered on pome fruit. Make applications on a 7 - 10 day interval.
	Pink, bloom, post-bloom Scab <i>(Venturia spp.)</i>		Apply Vango WG in tank mix combination with the label rate of a protectant fungicide. Use of the label rate of an EBDC fungicide will broaden the disease control spectrum to include cedar apple rust. The addition of the label rate of sulfur or a sterol inhibitor (SI) fungicide to this tank mix will broaden the disease control spectrum to include powdery mildew.

¹ **Complete List of Pome Fruit Crops:** Apple; Azarole; Crabapple; Loquat; Mayhaw; Medlar; Pear; Pear, Asian; Quince; Quince, Chinese; Quince, Japanese; Tejocote and cultivars, varieties and/or hybrids of these.

Application Instructions: Application may be made by ground or air. Good coverage is essential for good disease control. Use a minimum of 20 gallons/A spray volume by air. Make no more than two applications by air.

Specific Use Restrictions:

1. When applying to pears, apply **Vango WG** in tank mix only.
2. Make no more than two applications by air.
3. Do not apply more than 30 oz/A of **Vango WG** (1.4 lb ai/A of cyprodinil) per year.
4. Do not apply within 0 days of harvest (0-day PHI).

Crop	Disease	Product Rate oz/Acre	Remarks
Stone Fruits Crop Group 12-12¹ Apricots Nectarines Peaches Plums Prunes Tart Cherries And cultivars and/or hybrids of these	Brown Rot Blossom Blight <i>(Monilinia spp.)</i>	5	Begin applications at bloom stage (apricots at red bud, cherries at popcorn, peaches and nectarines at pink bud, plums and prunes at green tip). Make a second application at full bloom. Use Vango WG alone or in tank mixture with the label rate of another fungicide registered for stone fruit.
		Vango WG Alone 10	Apply a maximum of 2 applications of Vango WG during the preharvest period up through 2 days prior to harvest as needed.
		Vango WG Tank Mixtures 5	Apply Vango WG in tank mixture with the label rate of another fungicide registered on stone fruit for control of fruit brown rot.

¹ **Complete List of Stone Fruit:** Apricot; Apricot, Japanese; Cherry, tart; Jujube, Chinese; Nectarine; Peach; Plum; Plum, American; Plum, beach; Plum, Canada; Plum, cherry; Plum, Chickasaw; Plum, Damson; Plum, Japanese; Plum, Klamath; Plum, prune; Plumcot; Prune; Sloe; as well as other cultivars and hybrids of these.

Application Instructions: Application may be made by ground or air. Good coverage is essential for good disease control. Use a minimum of 20 gallons/A spray volume by air. Make no more than two applications by air.

Specific Use Restrictions:

1. Do not apply **Vango WG** to sweet cherries.
2. Make no more than two applications by air.
3. Do not apply more than 30 oz/A of **Vango WG** (1.4 lb ai/A of cyprodinil) per year.
4. Do not apply more than a maximum total of 4 applications (air plus ground) per year.
5. Do not apply within 2 days of harvest (2-day PHI).

Crop			Disease	Product Rate oz/Acre	Remarks
Tropical Fruits			Alternaria Fruit Rot (<i>Alternaria</i> spp.) Anthracnose (<i>Colletotrichum</i> spp.) Botrytis Fruit Rot (<i>Botrytis</i> spp.)	5.5 - 7	Make the first application during early bloom and repeat on 7 - 10 day intervals if conditions remain favorable for disease development. Resistance Management: After 2 applications of Vango WG , alternate with another fungicide with a different mode of action for 2 applications.
Acerola	Jaboticaba	Pulasan			
Avocado	Longan	Rambutan			
Black Sapote	Lychee	Sapodilla			
Canistel	Mamey Sapote	Spanish Lime			
Dragon Fruit	Mango	Star Apple			
Feijoa	Papaya	Starfruit			
Guava	Passionfruit	Wax Jambu			

Application Instructions: Application may be made by ground or air. Good coverage is essential for good disease control. Use a minimum of 20 gallons/A spray volume by air. Make no more than two applications by air.

Specific Use Restrictions:

1. Make no more than two applications by air.
2. Do not apply more than 28 oz/A of **Vango WG** (1.3 lb ai/A of cyprodinil) per year.
3. Do not apply more than a maximum total of 4 applications (air plus ground) per year.
4. May be applied on the day of harvest (0-day PHI).

CROP USE DIRECTIONS

BERRIES, FRUITS, HERBS AND VEGETABLES

Crop	Disease	Product Rate oz/Acre	Remarks
Artichoke, Globe*	Ramularia Bud Spot (<i>R. cynarae</i>) Ramularia Leaf Spot	5.5 - 7	Begin applications prior to disease onset when conditions are conducive for disease. Apply Vango WG on a 14-day schedule, making no more than 2 sequential applications before alternating to another fungicide with a different mode of action.

Application Instructions: For best results, sufficient water volume must be used to provide thorough coverage. **Vango WG** can be applied by ground, chemigation, or aerial application. For ground applications, apply in 50 - 200 gallons of water per acre to obtain coverage without excessive runoff. For aerial applications, use a minimum of 10 gallons/A of water. For chemigation, apply in 0.1 - 0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.

Specific Use Restrictions:

1. Do not apply more than 28 oz/A of **Vango WG** (1.3 lb ai/A of cyprodinil) per year.
2. Do not apply more than 1.3 lb ai/A/year of a cyprodinil-containing product.
3. Do not apply more than a maximum total of 4 applications (air plus ground plus chemigation) per year.
4. Do not apply within 3 days of harvest (3-day PHI).

*Not registered for use in California.

Crop	Disease	Product Rate oz/Acre	Remarks
Beans (Dried and Succulent except Cowpeas)* Bean (<i>Lupinus</i> spp.) (grain lupin, sweet lupin, white lupin, white sweet lupin) Bean (<i>Phaseolus</i> spp.) (kidney, lima, mung, navy, pinto, snap, wax) Bean (<i>Vigna</i> spp.) (asparagus, blackeyed pea) Broad Bean (fava bean) Chickpea (garbanzo bean)	Gray Mold (<i>Botrytis cinerea</i>)	5.5 - 7	Begin applications prior to or at the onset of disease and repeat applications on a 7-day interval if conditions remain favorable for disease development. Resistance Management: After 2 applications of Vango WG , alternate with another fungicide with a different mode of action for 2 applications.

Application Instructions: Application may be made by ground, air, or chemigation. Good coverage is essential for good disease control. Use a minimum of 5 gallons/A spray volume by air. Make no more than two applications by air. For chemigation, apply in 0.1 - 0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.

Specific Use Restrictions:

1. Make no more than two applications by air.
2. Do not apply more than 28 oz/A of **Vango WG** (1.3 lb ai/A of cyprodinil) per year.
3. Do not apply within 7 days of harvest (7-day PHI).

*Not registered for use in California.

Crop	Disease	Product Rate oz/Acre	Remarks
Berries - Bushberry Subgroup 13-07B* Aronia Berry Highbush Cranberry Black Currant Huckleberry Blueberry Jostaberry High And Low Bush Juneberry Buffalo Currant (saskatoon berry) Chilean Guava Lingonberry Edible Honeysuckle Native Currant Elderberry Red Currant European Barberry Salal Gooseberry Sea Buckthorn Berries - Caneberry Subgroup 13-07A* Blackberry Wild Raspberry Loganberry And cultivars and/or Red and Black hybrids of these Raspberry	Alternaria Fruit Rot (<i>Alternaria tenuissima</i>) Anthracnose (<i>Colletotrichum</i> spp.) Botrytis Fruit Rot (<i>Botrytis cinerea</i>) Mummy Berry (<i>Monilinia vaccinicorymbosi</i>)	5.5 - 7	Begin applications prior to or at the onset of disease and repeat applications on a 7 - 10 day interval if conditions remain favorable for disease development. Resistance Management: After 2 applications of Vango WG , alternate with another fungicide with a different mode of action for 2 applications.

Application Instructions: Application may be made by ground or air. Good coverage is essential for good disease control. Use a minimum of 5 gallons/A spray volume by air. Make no more than two applications by air.

Specific Use Restrictions:

1. Make no more than two applications by air.
2. Do not apply more than 28 oz/A of **Vango WG** (1.3 lb ai/A of cyprodinil) per year.
3. May be applied on the day of harvest (0-day PHI).

*Not registered for use in California.

Crop	Disease	Product Rate oz/Acre	Remarks
Brassica (Cole) Leafy Vegetables¹* Broccoli Kale Brussels Sprouts Mustard Greens Cabbage Turnip Greens Cabbage, Chinese And cultivars and/or Cauliflower hybrids of these Collards	Powdery Mildew (<i>Erysiphe polygoni</i>)	5.5 - 7	Begin applications prior to or at the onset of disease and repeat applications on a 7 - 10 day interval if conditions remain favorable for disease development. Resistance Management: After 2 applications of Vango WG , alternate with another fungicide with a different mode of action for 2 applications.

¹ **Additional Cole Vegetables:** Broccoli, Chinese; Broccoli raab; Cavolo broccolo; Kohlrabi; Mizuna; Mustard spinach; and Rape greens.

Application Instructions: Application may be made by ground, air, or chemigation. Good coverage is essential for good disease control. Use a minimum of 10 gallons/A spray volume by air. Make no more than two applications by air. For chemigation, apply in 0.1 - 0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.

Specific Use Restrictions:

1. Make no more than two applications by air.
2. Do not use roots of treated turnips for food or feed. Only turnip varieties harvested for their leaves may be treated.
3. Do not apply more than 28 oz/A of **Vango WG** (1.3 lb ai/A of cyprodinil) per year.
4. Do not apply within 7 days of harvest (7-day PHI).

*Not registered for use in California.

Crop	Disease	Product Rate oz/Acre	Remarks
Cucurbits* Cantaloupe Muskmelon Chayote Pumpkin Chinese Waxgourd Squash Cucumber Watermelon Gourds Zucchini Honeydew And cultivars and/or <i>Momordica</i> spp. hybrids of these (bitter melon, balsam apple)	Alternaria Leaf Blight (<i>A. cucumerina</i>) Alternaria Leaf Spot (<i>A. alternate</i>) Gummy Stem Blight (<i>Didymella bryoniae</i>) Powdery Mildew (<i>Erysiphe cichoracearum</i> , <i>Sphaerotheca fuliginea</i>)	5.5 - 7	Begin applications prior to or at the onset of disease and repeat applications on a 7 - 10 day interval if conditions remain favorable for disease development. Resistance Management: After 2 applications of Vango WG , alternate with another fungicide with a different mode of action for 2 applications.

Application Instructions: Application may be made by ground, air, or chemigation. Good coverage is essential for good disease control. Use a minimum of 10 gallons/A spray volume by air. Make no more than two applications by air. For chemigation, apply in 0.1 - 0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.

Specific Use Restrictions:

1. Make no more than two applications by air.
2. Do not apply more than 28 oz/A of **Vango WG** per year.
3. Do not apply more than 1.3 lb ai/A of cyprodinil-containing products per year.
4. May be applied up to 1 day before harvest (1-day PHI).

*Not registered for use in California.

Crop	Disease	Product Rate oz/Acre	Remarks
Herbs^{1*} (Dried and Fresh) Basil Parsley Chive Rosemary Coriander, Leaves (cilantro) Sage Dillweed Tarragon Lemongrass Thyme	Alternaria Leaf Spot <i>(Alternaria spp.)</i> Botrytis Leaf Blight <i>(Botrytis spp.)</i>	5.5 - 7	Begin applications prior to or at the onset of disease and repeat applications on a 7 - 10 day interval if conditions remain favorable for disease development. Resistance Management: After 2 applications of Vango WG , alternate with another fungicide with a different mode of action for 2 applications.
¹ Additional Herbs (dried and fresh): Angelica, Balm, Borage, Burnet, Camomile, Catnip, Chervil (dried leaves), Clary, Costmary, Culantro (leaves), Curry (leaves), Horehound, Hyssop, Lavender, Lovage (leaves), Marigold, Marjoram, Nasturtium, Pennyroyal, Rue, Savory (summer and winter), Sweet bay, Tansy, Wintergreen, Woodruff, and Wormwood.			
Application Instructions: Application may be made by ground, air, or chemigation. Good coverage is essential for good disease control. Use a minimum of 10 gallons/A spray volume by air. Make no more than two applications by air. For chemigation, apply in 0.1 - 0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.			
Specific Use Restrictions: 1. Make no more than two applications by air. 2. Do not apply more than 28 oz/A of Vango WG (1.3 lb ai/A of cyprodinil) per year. 3. Do not apply within 7 days of harvest (7-day PHI). *Not registered for use in California.			

Crop	Disease	Product Rate oz/Acre	Remarks
Leafy Greens Subgroup 4A (except Brassica)* and Leaf Petioles Subgroup 4B* Amaranth Cress Purslane Arugula Dandelion Radicchio Cardoon Dock Rhubarb Celery Endive (escarole) Spinach Celery, Chinese Fennel, Florence Spinach Vine Celtuce Lettuce, Head and Leaf Swiss Chard Chervil New Zealand Spinach And cultivars and/ or hybrids of these Chrysanthemum, Edible Orach Corn salad Parsley	Alternaria Leaf Spot <i>(Alternaria spp.)</i> Gray Mold <i>(Botrytis cinerea)</i> Suppression: Powdery Mildew <i>(Erysiphe cichoracearum)</i>	5.5 - 7	Begin applications prior to or at the onset of disease and repeat applications on a 7 - 10 day interval if conditions remain favorable for disease development. Resistance Management: After 2 applications of Vango WG , alternate with another fungicide with a different mode of action for 2 applications.
Application Instructions: Application may be made by ground, air, or chemigation. Good coverage is essential for good disease control. Use a minimum of 10 gallons/A spray volume by air. Make no more than two applications by air. For chemigation, apply in 0.1 - 0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.			
Specific Use Restrictions: 1. Make no more than two applications by air. 2. Do not apply more than 28 oz/A of Vango WG (1.3 lb ai/A of cyprodinil) per year. 3. May be applied on the day of harvest (0-day PHI). *Not registered for use in California.			

Crop		Disease	Product Rate oz/Acre	Remarks
Leaves of Root and Tuber Vegetables^{1*}		Alternaria Leaf Blight (<i>Alternaria dauci</i>) Powdery Mildew (<i>Erysiphe</i> spp.)	5.5 - 7	Begin applications prior to or at the onset of disease and repeat applications on a 7 - 10 day interval if conditions remain favorable for disease development. Resistance Management: After 2 applications of Vango WG , alternate with another fungicide with a different mode of action for 2 applications.
Carrot	Radish (oriental)			
Beet, Garden	Rutabaga			
Beet, Sugar	Sweet Potato			
Parsnip	Turnip			
Radish	Yam (true)			

¹ **Additional Leaves of Root and Tuber Vegetables:** Burdock (edible), Cassava, Celeriac, Chicory, Dasheen, Salsify (including black and Spanish), Tanier, and Turnip-rooted chervil.

Application Instructions: Application may be made by ground, air, or chemigation. Good coverage is essential for good disease control. Use a minimum of 10 gallons/A spray volume by air. Make no more than two applications by air. For chemigation, apply in 0.1 - 0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.

Specific Use Restrictions:

1. Make no more than two applications by air.
2. Radish ONLY - Make no more than two applications per year.
3. Radish ONLY - Do not apply more than 14 oz of **Vango WG** per crop per year or more than 0.66 lb ai/A of cyprodinil-containing products per year.
4. Do not apply more than 28 oz/A of **Vango WG** per year for specified leaves of root and tuber vegetables except radish.
5. Do not apply more than 1.3 lb ai/A of cyprodinil-containing products per year for specified leaves of root and tuber vegetables except radish.
6. Do not apply within 7 days of harvest (7-day PHI).
7. Do not allow cattle or other livestock to feed upon the leaves of root vegetables.

*Not registered for use in California.

Crop			Disease	Product Rate oz/Acre	Remarks
Onions - Bulb Vegetables Crop Group 3-07A and 3-07B			Botrytis Leaf Blight or Blast (<i>Botrytis</i> spp.) Purple Blotch (<i>Alternaria porri</i>) Suppression: Neck Rot (<i>Botrytis</i> spp.)	10	Begin applications prior to or at the onset of disease and repeat applications on a 7 - 10 day interval if conditions remain favorable for disease development. For optimal effect on neck rot, apply on a 7-day schedule at the 10 oz rate. Resistance Management: After 2 applications of Vango WG , alternate with another fungicide with a different mode of action for 2 applications.
Bulb Onion	Green Onion	Green Onion (continued)			
Chinese Onion	Beltsville Bunching Onion	Shallot Fresh Leaves			
Daylily Bulb	Chinese Chive Fresh Leaves	Tree Tops Onion			
Dry Bulb Onion	Fresh Chive Leaves	Welsh Onion Tops			
Fritillaria Bulb	Fresh Onion	Wild Leek			
Garlic	Fritillaria Leaves	Wild Onion			
Great-headed Garlic	Green Onion	Onions Grown for Seed			
Lily Bulb	Hosta Elegans	And cultivars and/ or			
Pearl Onion	Kurrat	hybrids of these			
Potato Onion	Lady's Leek				
Serpent Garlic	Leek				
Shallot	Macrostem Onion				

Application Instructions: Application may be made by ground, air, or chemigation. Good coverage is essential for good disease control. Use a minimum of 5 gallons/A spray volume by air. Make no more than two applications by air. For chemigation, apply in 0.1 - 0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.

Specific Use Restrictions:

1. Make no more than two applications by air.
2. Do not apply more than 28 oz/A of **Vango WG** (1.3 lb ai/A of cyprodinil) per year.
3. Do not apply within 7 days of harvest (7-day PHI).

Crop	Disease	Product Rate oz/Acre	Remarks
Tuberous and Corm Vegetables Crop Subgroup 1C^{1*} Potatoes* Sweet Potatoes*	Brown Spot (<i>Alternaria alternata</i>) Early Blight (<i>A. solani</i>) Powdery Mildew (<i>Erysiphe cichoracearum</i>) Septoria Leaf Spot (<i>Septoria lycopersici</i>) Tan Spot (<i>Botrytis cinerea</i>)	5.5 - 7	Begin applications prior to or at the onset of disease and repeat applications on a 7 - 10 day interval if conditions remain favorable for disease development. Resistance Management: After 2 applications of Vango WG , alternate with another fungicide with a different mode of action for 2 applications.

¹ **Additional Vegetables, Tuberous and Corm Subgroup 1C:** Arracacha, Arrowroot, Artichoke (Chinese and Jerusalem), Canna, Cassava (bitter and sweet), Chayote (root), Chufa, Dasheen (Taro), Ginger, Leren, Tanier, Turmeric, Yam (bean and true), and cultivars and/or hybrids of these

Application Instructions: Application may be made by ground, air, or chemigation. Good coverage is essential for good disease control. Use a minimum of 5 gallons/A spray volume by air. Make no more than two applications by air. For chemigation, apply in 0.1 - 0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.

Specific Use Restrictions:

1. Make no more than two applications by air.
2. Do not apply more than 28 oz/A of **Vango WG** per year.
3. Do not apply more than 1.3 lb ai/A of cyprodinil-containing products per year.
4. Do not apply within 14 days of harvest (14-day PHI).

*Not registered for use in California.

Crop	Disease	Product Rate oz/Acre	Remarks
Root and Tuber Vegetables except Sugar Beet^{1*} Carrot Beet, Garden Ginseng Horseradish Parsnip	Alternaria Leaf Blight (<i>Alternaria dauci</i>) Powdery Mildew (<i>Erysiphe spp.</i>)	5.5 - 7	Begin applications prior to or at the onset of disease and repeat applications on a 7 - 10 day interval if conditions remain favorable for disease development. Resistance Management: After 2 applications of Vango WG , alternate with another fungicide with a different mode of action for 2 applications.

¹ **Additional Root and Tuber Vegetables:** Burdock, edible; Celeriac; Chicory; Salsify (including black and Spanish); Skirret; Turnip-rooted parsley; and Turnip-rooted chervil.

Application Instructions: Application may be made by ground, air, or chemigation. Good coverage is essential for good disease control. Use a minimum of 5 gallons/A spray volume by air. Make no more than two applications by air. For chemigation, apply in 0.1 - 0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.

Specific Use Restrictions:

1. Make no more than two applications by air.
2. Radish ONLY - Make no more than two applications per year.
3. Radish ONLY - Do not apply more than 14 oz of **Vango WG** per crop per year or more than 0.66 lb ai/A of cyprodinil-containing products per year.
4. Do not apply more than 28 oz/A of **Vango WG** per year for specified leaves of root and tuber vegetables, except radish.
5. Do not apply more than 1.3 lb ai/A of cyprodinil-containing products per year for specified leaves of root and tuber vegetables, except radish.
6. Do not apply within 7 days of harvest (7-day PHI).
7. Do not allow cattle or other livestock to feed upon the leaves of root vegetables.

*Not registered for use in California.

Crop	Disease	Product Rate oz/Acre	Remarks
Berry, Low Growing Subgroup 13- 07G (except Cranberry)¹* Strawberry*	Anthracnose (<i>Colletotrichum</i> spp.) Gray Mold (<i>Botrytis cinerea</i>) Powdery Mildew (<i>Sphaerotheca macularis</i>)	5.5 - 10	Begin application at or before bloom and continue on a 7 - 10 day interval. Resistance Management: After 2 applications of Vango WG , alternate with another fungicide with a different mode of action for 2 applications.
	Root and Crown Anthracnose at-planting (<i>Colletotrichum</i> spp.)	2.5 - 4 oz per 100 gal water	Apply as a pre-plant dip to strawberry roots and crowns at the rate of 2.5 to 4 oz per 100 gal of water for suppression of root and crown rot caused by anthracnose. Wash transplants to remove excess soil prior to dipping. This helps to remove adhering spores from the external plant parts. Completely immerse planting stock in dip solution. Dip or expose plants for a minimum of 2 to 5 minutes. DO NOT reuse solution. Dispose of dip solution according to local regulations. Plant treated plants as quickly as possible. For continued anthracnose control, follow with foliar applications beginning 2 - 3 weeks after transplant.
¹ Additional Low Growing Berries: Bearberry; bilberry; cloudberry; muntries; partridgeberry and cultivars and/or hybrids of these.			
Application Instructions: Application may be made by ground, air, or chemigation. Good coverage is essential for good disease control. Use a minimum of 5 gallons/A spray volume by air. Make no more than two applications by air. For chemigation, apply in 0.1 - 0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.			
Specific Use Restrictions: 1. Make no more than two applications by air. 2. Make only one pre-plant dip application per crop. 3. Do not apply more than 28 oz/A of Vango WG (1.3 lb ai/A of cyprodinil) per year. 4. May be applied on the day of harvest (0-day PHI). *Not registered for use in California.			

SPECIMEN

Crop	Disease	Product Rate oz/Acre	Remarks
Fruiting Vegetable Crop Group 8-10^{1*} Eggplant Groundcherry Pepino Pepper (includes bell pepper, chili pepper, cooking pepper, pimento, sweet pepper) Tomatillo Tomatoes	Early Blight (<i>Alternaria solani</i>) Grey Mold (<i>Botrytis cinerea</i>) Powdery Mildew (<i>Leveillula taurica</i>)	5.5 - 7	Begin applications prior to or at the onset of disease and repeat applications on a 7 - 10 day interval if conditions remain favorable for disease development. Resistance Management: After 2 applications of Vango WG , alternate with another fungicide with a different mode of action for 2 applications.

¹ **Additional Fruiting Vegetables:** African eggplant; bush tomato; cocona; currant tomato; garden huckleberry; goji berry; martynia; naranjilla; okra; pea eggplant; pepino; pepper, nonbell; roselle; scarlet eggplant; sunberry; tree tomato; cultivars, varieties, and/or hybrids of these.

Application Instructions: Application may be made by ground, air, or chemigation. Good coverage is essential for good disease control. Use a minimum of 10 gallons/A spray volume by air. Make no more than two applications by air. For chemigation, apply in 0.1 - 0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.

Specific Use Restrictions:

1. Make no more than two applications by air.
2. Do not apply more than 28 oz/A of **Vango WG** per year.
3. Do not apply more than 1.3 lb ai/A of cyprodinil-containing products per year.
4. Do not apply more than a maximum total of 4 applications (air plus ground plus chemigation) per year.
5. May be applied on the day of harvest (0-day PHI).

*Not registered for use in California.

Crop	Disease	Product Rate oz/Acre	Remarks
Watercress*	Cercospora Leaf Spot (<i>Cercospora</i> spp.)	5.5 - 7	Begin applications prior to or at the onset of disease and repeat applications on a 7 - 10 day interval if conditions remain favorable for disease development. Resistance Management: After 2 applications of Vango WG , alternate with another fungicide with a different mode of action for 2 applications.

Application Instructions: Application may be made by ground or chemigation. Good coverage is essential for good disease control. For chemigation, apply in 0.1 - 0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.

Specific Use Restrictions:

1. Do not apply more than 28 oz/A of **Vango WG** (1.3 lb ai/A of cyprodinil) per year.
2. May be applied on the day of harvest (0-day PHI).
3. Applications can be made to a dry bed only. Do not apply directly to water.

*Not registered for use in California.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Keep this product in its tightly closed original container, when not in use. Store in a cool, dry (preferably locked) area that is inaccessible to children and animals.

PESTICIDE DISPOSAL: Pesticide wastes may be toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING: Nonrefillable Container (≤ 50 pounds): Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

LIMITATION OF WARRANTY AND LIABILITY

IMPORTANT: READ BEFORE USE. Read the entire Directions for Use, Conditions of Warranties and Limitations of Liability before using this product. If these terms and conditions are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Disclaimer of Warranties and Limitations of Liability. **CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Ineffectiveness, injury, and other unintended consequences may result because of such factors as manner of use or application (including misuse), the presence of other materials, weather conditions, and other unknown factors, all of which are beyond the control of ATTICUS, LLC. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, ATTICUS, LLC makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond statements on this label. **LIMITATIONS OF LIABILITY:** To the extent consistent with applicable law, neither ATTICUS, LLC the manufacturer, nor the Seller shall be liable for any indirect, special, incidental or consequential damages resulting from the use, handling, application, storage, or disposal of this product. To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use, handling, application, or storage of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid.

Vango™ is a trademark of Atticus, LLC

Vanguard® is a registered trademark of Syngenta Group Company.

F20190930c

SPECIMEN