

**HERBICIDE**

WILBUR-ELLIS®

Ideas to Grow With™

VENGEANCE® PLUS

SELECTIVE HERBICIDE

FOR SELECTIVE BROADLEAF WEED CONTROL IN ORNAMENTAL LAWNS AND TURF GRASSES, SOD FARMS, NON-TURF AREAS, INDUSTRIAL VEGETATION MANAGEMENT AND NON-CROPLAND USE.

CONTAINS MCPA, TRICLOPYR AND DICHLORPROP-p

CONTROLS: Dandelion, Chickweed, Black medic, Knotweed, Plantain, Oxalis, Clover, Cocklebur, Thistle and many other species of broadleaf weeds; some of which are listed on this label.

ACTIVE INGREDIENTS:	% By Wt.
Dimethylamine Salt of 2-Methyl-4-Chlorophenoxyacetic Acid*	46.87%
Triethylamine Salt of 3,5,6-Trichloro-2-Pyridinyloxyacetic Acid**	10.68%
Dimethylamine Salt of (+)-R-2-(2,4-Dichlorophenoxy) propionic Acid***‡	9.12%
OTHER INGREDIENTS:	33.33%
	100.00%

By Isomer specific AOAC Method Equivalent to:

*2-Methyl-4-Chlorophenoxyacetic Acid	38.27%, 3.72 lbs./gal.
**3,5,6-Trichloro-2-Pyridinyloxyacetic Acid	7.65%, 0.75 lbs./gal.
***(+)-R-2-(2,4-Dichlorophenoxy) propionic Acid	7.65%, 0.75 lbs./gal.

‡CONTAINS THE SINGLE ISOMER FORM OF DICHLORPROP-p

For Use By Turf Maintenance Personnel, Landscaping or Commercial Applicators Only

EPA REG. NO. 228-411-2935

EPA EST. NO. 228-IL-001

KEEP OUT OF REACH OF CHILDREN
DANGER - PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
 (If you do not understand the label, find someone to explain it to you in detail.)

SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS.

For Chemical Spill, Leak, Fire or Exposure, Call CHEMTREC (800) 424-9300

For Medical Emergencies Only, Call (877) 325-1840

NET CONTENTS: 2.5 GALLONS (9.46 Liters)
Nonrefillable Container

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS DANGER – PELIGRO

Corrosive. Causes irreversible eye damage. May be fatal if absorbed through skin. Harmful if swallowed. Do not get in eyes, on skin or on clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are made out of any waterproof material. If you want more options, follow the instructions for category A on an EPA chemical-resistance selection chart.

Mixers, loaders, applicators and other handlers must wear:

- Protective eyewear (goggles or face shield)
- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant gloves
- Chemical-resistant footwear and socks
- Chemical-resistant headgear for overhead exposure, and
- Chemical-resistant apron when cleaning equipment

See engineering controls statements for further instructions.

Follow manufacturer's instructions for cleaning/maintaining Personal Protective Equipment (PPE). If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENTS

If the mechanical system is used in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Discard clothing and other absorbent material that have been drenched or heavily contaminated with the product's concentrate. Do not reuse them.

FIRST AID

If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor for treatment advice. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give any liquid to the person. Do not give anything by mouth to an unconscious person.
<p>HOT LINE NUMBER: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.</p>	
<p>NOTE TO PHYSICIAN: For eye irritation, examination by an ophthalmologist may be indicated. If swallowed, probable mucosal damage may contraindicate the use of gastric lavage.</p>	

ENVIRONMENTAL HAZARDS

This pesticide may be toxic to fish, aquatic invertebrates, aquatic plants and may adversely affect other non-target plants. Drift or runoff may adversely affect non-target plants and may be hazardous to aquatic organisms in water adjacent to treated areas. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Aerial application of this product is prohibited.

Do not use this product on or near desirable plants, including within the drip line of the roots of desirable trees and shrubs, since injury may result.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: Coveralls, chemical-resistant gloves made of any waterproof material, shoes plus socks, and protective eyewear.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter or allow others to enter the treated area until sprays have dried.

This product is for use on Ornamental Turf Lawns (Residential, Industrial and Institutional), Parks, Cemeteries, Athletic Fields and Golf Courses (Fairways, Aprons, Tees* and Roughs), Non-crop Areas and Sod Farms.

*Excluding Bentgrass Tees

USE PRECAUTIONS

Avoid drift of spray mist to vegetables, grapes, flowers, ornamental plants, shrubs, trees and other desirable plants. Do not pour spray solutions near desirable plants. Do not use on carpetgrass, Dichondra, nor on lawns or turf where desirable clovers are present. Avoid fine mists. Use lawn type sprayer with coarse spray as wind drift is less likely. Avoid contact with exposed feeder roots of ornamentals and trees. Maximum control of weeds will be obtained from spring or early fall applications when weeds are actively growing. Use the higher rate for hard-to-control weeds. Do not exceed specified dosages for any area; be particularly careful within the dripline of tree and other ornamental species. Do not apply to newly seeded grasses until well established. Do not spray when air temperature exceeds 90°F.

Avoid application to Centipede or St. Augustine grasses unless turf injury can be tolerated. The suitable use of this product on non-recommended turf species may be determined by treating a small area at any rate per acre which does not exceed 3 pints per acre. Prior to treatment of larger areas, the treated area should be observed for any sign of turf injury for a period of 30 days of normal growing conditions to determine the phytotoxicity and efficacy to the treated area. The user assumes the responsibility for any plant damage or other liability resulting from the use of this product on species not recommended on this label.

For optimum results: (1) avoid applying during excessively dry or hot periods unless irrigation is used; (2) turf should not be mowed 1 to 2 days before and following application; (3) reseed no sooner than 3 to 4 weeks after application of this product. Adding oil, wetting agent, or other surfactant to the spray may be used to increase effectiveness on weeds but doing so may reduce selectivity to turf resulting in turf damage. Clean and rinse spray equipment using soap or detergent and water, and rinse thoroughly before reuse for other sprays.

WEEDS CONTROLLED

This product is designed to control many broadleaf weeds such as:

Alder	Burdock	Dead nettle	Henbit	Narrowleaf
Annual yellow	Burning nettle	Dock	Hoary cress	plantain
sweet clover	Bur ragweed	Dogbane	Hoary plantain	Narrowleaf vetch
Artichoke	Burweed	Dogfennel	Hoary vervain	Nettle
Aster	Buttercup	Elderberry	Honeysuckle	Orange hawkweed
Austrian fieldcress	Canada thistle	English daisy	Hop clover	Oxalis
Bedstraw	Carolina geranium	Fall dandelion	Horsenettle	Oxeye daisy
Beggartick	Carpetweed	False dandelion	Horsetail	Parsley-piert
Biden	Catchweed	False flax	Indiana mallow	Parsnip
Bindweed	bedstraw	False sunflower	Ironweed	Pearlwort
Bird vetch	Catsear	Fiddleneck	Jewelweed	Pennygrass
Bitterweed	Catnip	Field bindweed	Jimsonweed	Pennywort
Bitter wintercress	Chickweed	Field pansy	Kochia	Peppergrass
Black-eyed Susan	Chicory	Flea bane (daisy)	Knawel	Pepperweed
Black medic	Cinquefoil	Fixweed	Knotweed	Pigweed
Black mustard	Clover	Florida betony	Lambsquarter	Pineywoods
Black-seed	Cockle	Florida pusley	Lespedeza	bedstraw
plantain	Cocklebur	Frenchweed	Locoweed	Plains coreopsis
Blessed thistle	Coffebean	Galinsoga	Lupine	Plantain
Blue lettuce	Coffeeweed	Garlic mustard	Mallow	Poison hemlock
Blue vervain	Common	Goathead	Marshelder	Poison ivy
Box elder	chickweed	Goatsbeard	Matchweed	Poison oak
Bracted plantain	Common mullein	Goldenrod	Mexicanweed	Pokeweed
Brassbuttons	Common	Ground ivy	Milk vetch	Poorjoe
Bristly oxtongue	sowthistle	Gumweed	Milkweed	Poverty weed
Broadleaf dock	Corn chamomile	Hairy bittercress	bloodflower	Prickly lettuce
Broadleaf plantain	Creeping jenny	Hairy fleabane	Mugwort	Prickly sida
Broomweed	Crimson clover	Hawkweed	Morningglory	Primrose
Buckhorn	Croton	Healall	Mousear	Prostrate
Buckhorn plantain	Cudweed	Heartleaf drymary	knotweed	knotweed
Bulbous buttercup	Curly dock	Hedge bindweed	Musk thistle	Prostrate pigweed
Bull nettle	Curly indigo	Hedge mustard	Mustard	Prostrate spurge
Bull thistle	Dandelion	Hemp		Prostrate vervain

(continued)

WEEDS CONTROLLED (cont.)

Puncture vine	Smartweed	Sumac	Virginia	Wild radish
Purslane	Smooth dock	Sunflower	buttonweed	Wild rape
Ragweed	Smooth pigweed	Sweet clover	Virginia creeper	Wild strawberry
Red clover	Sneezeweed	Tall nettle	Virginia	Wild sweet potato
Redroot pigweed	Southern wild	Tall vervain	pepperweed	Wild vetch
Red sorrel	rose	Tansy mustard	Wavyleaf	Willow
Redstem filaree	Sowthistle	Tansy ragwort	bulbthistle	Witchweed
Rough cinquefoil	Spanish needle	Tanweed	Western clematis	Woolly
Rough fleabane	Spatterdock	Tarweed	Western salsify	morningglory
Roundleafed	Speedwell	Thistle	White clover	Woodsorrel
marigold	Spiny amaranth	Tick trefoil	White mustard	Woolly croton
Rush	Spiny cocklebur	Toadflax	Wild mustard	Woolly plantain
Russian pigweed	Spotted catsear	Trailing	Wild aster	Wormseed
Russian thistle	Spotted	crownvetch	Wild buckwheat	Yarrow
St. Johnswort	knapweed	Tumble mustard	Wild carrot	Yellow rocket
Scarlet pimpernel	Spotted spurge	Tumble pigweed	Wild four-o'-clock	Yellowflower
Scotch thistle	Spurge	Tumbleweed	Wild garlic	pepperweed and
Sheep sorrel	Spurweed	Velvet leaf	Wild geranium	other broadleaf
Shepherd's purse	Stinging nettle	Venice mallow	Wild lettuce	weeds
Slender plantain	Stinkweed	Veronica	Wild marigold	
Smallflower	Stichwort	Vervain	Wild onion	
galinsoga	Strawberry clover	Vetch	Wild parsnip	

ORNAMENTAL LAWNS AND TURF

Apply this product at the rate of 2 to 3 pints in 20 to 240 gallons of water per acre (0.73 to 1.10 fluid ounces in 0.5 to 5.5 gallons of water per 1,000 square feet) to control weeds growing in turf grasses planted to Bluegrass, Fescue, Rye, Bent (excluding golf course greens and tees), Bahia, Bermuda and Zoysia.

For residential turf and golf course uses, do not apply more than 1.5 lbs ae MCPA (3 pints) per acre per year. Do not apply more than 2 applications per year with a minimum retreatment interval of 30 days.

Controlled Droplet Applicators - (CDA), Atomizers, and Spinning Disk Applicators: Use this product at the rate of 2 to 3 pints per acre (0.73 to 1.10 fluid ounces per 1,000 square feet) in sufficient water to assure coverage (1 to 4 gallons of water per acre is normal for this type of equipment).

NOTE: For all grasses: (1) Do not overlap spray patterns; (2) Use reduced rates if grass is stressed from heat, drought, etc.; and (3) Follow CDA equipment spray instructions.

SOD FARMS

For sod farms, do not apply more than 1.5 lb of MCPA (3 pints) per acre per year. Do not apply more than 2 applications per year with a minimum retreatment interval of 30 days.

This product is intended for use on sod farms. This herbicide provides selective broadleaf control in warm season and cool season turfgrass established for commercial sod production.

Timing

Apply this herbicide to broadleaf weeds that are actively growing. Follow-up applications may be required for dense infestations of perennial and biennial weeds. Do not apply this product to Floratam St. Augustinegrass.

The application of this herbicide to grass seedlings is recommended after the second mowing.

The application of this product to newly sodded, sprigged, or plugged grasses should be delayed until 3 or 4 weeks after the sodding, sprigging, or plugging operations. Also, applications to dormant Bermudagrass, dormant Zoysiagrass and dormant Bahiagrass are suggested.

Rates for Sod Farms

For Bluegrass, Ryegrass and Fescue, apply 1-1/2 to 3 pints per acre. For Creeping Bentgrass use .9 pints per acre. For Common and Hybrid Bermuda, Bahia, Zoysia and St. Augustine grass apply 1 to 2 pints per acre. Spray volume for Sod Farm application is 5 to 175 gallons per acre.

Generally, the lower application rates will provide satisfactory control of sensitive weeds. The higher application rates will be required for dense infestations of perennial weeds, for adverse/extreme environmental conditions, or for weeds beyond the appropriate growth stages.

NON-CROP AREAS

(Rights of Way, Roadsides, Fence Rows, Hedge Rows, Railroads, Industrial Management Sites, Non-irrigation Ditch banks, Airports)

For Broadcast Treatment:

DO NOT apply more than 1.5 lbs ae MCPA (3.2 pints of this product) per acre per year.

DO NOT apply more than 2 applications per year with a minimum retreatment interval of 30 days.

For Spot Treatments:

DO NOT apply more than 3.0 lbs ae MCPA (6.4 pints of product) per acre per year. Spot treatment is defined as a treatment area no greater than 1,000 ft. sq. per acre.

APPLICATION DIRECTIONS

Herbaceous Broadleaf Weed Control

Apply 1.6 to 3.2 pints of this product in sufficient volume of water to allow for thorough coverage of weeds. Do not apply more than 3.2 pints of this product per acre per year.

Woody Brush and Vine Control/ Suppression

High Volume Foliar Spot Applications: Mix 4.8 to 6.4 pints of this product in sufficient volume of water to allow for thorough coverage. Spray volume applied will depend on the size and density of the brush to be treated, but do not apply more than 6.4 pints of this product (3.0 lbs ae MCPA) per acre per year. Direct the spray to treat all foliage, stems, and root collars to wet.

Broadcast Applications with Ground Equipment: Apply up to 3.2 pints of this product in sufficient volume of water to allow for thorough coverage per treated acre. Do not apply more than 3.2 pints of product (1.5 lbs ae MCPA) per acre per year. Spray all foliage, stems and root collars to wet. For optimum control of woody species, a tank mix may be necessary.

BRUSH, BROADLEAF, INVASIVE, AND NOXIOUS WEED SPECIES CONTROLLED

SPECIES CONTROLLED

When used as directed, this product will control or suppress many herbaceous broadleaf weeds (annual, biennial, and perennial) as well as suppress many unwanted woody plant and vine species.

Species controlled include:

ANNUALS

Buckwheat	Knapweed	Purslane
Carpetweed	Kochia	Ragweed
Chickweed	Lambsquarter	Russian thistle
Clover	Morningglory	Smartweed
Cocklebur	Mustard	Velvetleaf
Daisy, English	Pigweed	
Henbit	Puncturevine	

BIENNIALS

Ragwort, Tansy	Thistle, Musk
----------------	---------------

PERENNIALS

Bindweed, Field	Dogfennel	Sorrel, Sheep
Carrot, Wild (Queen Anne's Lace)	Knapweed, Russian	Spurge, Leafy
Dock, Curly	Milkweed	Thistle, Canada
	Ragweed, Perennial	Toadflax, Dalmatian

WOODY BRUSH AND VINES*

Alder	Cucumbertree	Locust	Redcedar, Eastern	Snowberry
Ash	Dogwood	Maple	Redvine	Spruce
Aspen	Elderberry	Oak	Rose, Multiflora	Sumac
Basswood	Elm	Olive, Russian	Sagebrush	Sycamore
Beech	Gum	Persimmon	Sassafras	Trumpet creeper
Birch	Hawthorn	Pine	Schinus	Waxmyrtle
Blackberry	Hemlock	Plum, Wild	(Florida Holly)	Willow
Cherry	Honeysuckle	Poplar	Brazil Peppertree,	Witchhazel
Creeper, Virginia	Ivy, Poison	Puncturevine	Christmas-berry)	Yaupon
Creosotebush	Kudzu	Raspberry	Serviceberry	

*Suppression

TANK MIX TREATMENTS

TANK MIX RECOMMENDATIONS ARE FOR USE ONLY IN STATES WHERE THE TANK MIX PRODUCT AND APPLICATION SITE ARE REGISTERED. READ AND FOLLOW THE LABEL OF EACH TANK MIX PRODUCT USED FOR PRECAUTIONARY STATEMENTS, DIRECTIONS FOR USE, AND OTHER RESTRICTIONS. For broader spectrum control, this product may be tank mixed with other herbicides registered for non-cropland uses (e.g. railroad, highway, pipeline, forest brush control, etc.), including but not limited to the following herbicides, in accordance with the more restrictive of label limitations and precautions. No label dosage rates should be exceeded. This product cannot be mixed with any product containing a label prohibition against such mixing. Add water to the spray tank prior to the addition of the tank mix products. Do not premix concentrates.

COMPATIBILITY

A compatibility test is recommended prior to mixing with liquid fertilizers and liquid iron. All regulations, either state or Federal, relating to the application of liquid fertilizers or liquid iron and this product must be strictly followed.

HERBICIDE

Asulam (Asulox [®])	Metsulfuron Methyl (Patriot [®])**
Bromacil (Hyvar [®])	MSMA
Chlorsulfuron (Telar [®])	Picloram (Tordon [®])*
Dicamba (Diablo [®])	Simazine (Princep [®])
Diuron (Karmex [®])	Sulfometuron methyl (Spyder [®])
Fosamine ammonium (Krenite [®])	Tebuthiuron (Spike [®])
Glyphosate (Razor [®] Pro)	Triclopyr (Tahoe [®])
Hexazinone (Velpar [®])	2,4-D
Imazapyr (Arsenal [®])	2,4-DP

Due to variations that may occur in formulated products and specific use ingredients (e.g. water supplies) a COMPATIBILITY TEST is recommended prior to actual tank mixing.

*Restricted use pesticides limited to certified applicators.

** Using Vengeance[®] Plus and Patriot[®] to reduce the development and spread of resistant biotypes and problem weeds such as Kochia and Russian thistle: some commonly resistant weeds typically require multiple spray applications to obtain adequate control. To reduce the number of applications required, applicators can utilize a mixture of Patriot[®] and Vengeance[®] Plus. Apply as a tank mix for post-emergence as follows: Add .5 to 1 ounce of Patriot[®] into 1.6 to 4.8 pints of Vengeance[®] Plus per acre. A CPDA-certified nonionic or silicone surfactant may be used for wetting, penetration or drift control.

NOTE: All intended tank mix combinations should be used only in recommended areas on the same broadleaf weed species found on both labels. For application methods and other use specifications, use the most restricted limitations from labeling of both products.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Always use original container to store pesticides in a secured warehouse or storage building. Store at temperatures above 32°F. If allowed to freeze, remix before using. This does not alter this product. Containers should be opened in well ventilated areas. Keep container tightly sealed when not in use. Do not stack cardboard cases more than two pallets high. Do not store near open containers of fertilizer, seed or other pesticides.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. If container is damaged or if pesticide has leaked, contain all spillage. Absorb and clean up all spilled material with granules or sand. Place in a closed labeled container for proper disposal. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Nonrefillable Containers 5 Gallons or Less: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

Nonrefillable containers larger than 5 gallons: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. If recycling or reconditioning is not available, puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds.

Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Conditions of Sale and Limitation of Warranty and Liability:

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using the product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

ALL STATEMENTS MADE HEREIN ARE SUBJECT TO APPLICABLE LAW, AND TO THE EXTENT THERE IS ANY INCONSISTENCY OR CONTENTION, APPLICABLE LAW SHALL GOVERN.

The Directions for Use of the product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of many different factors including, without limitation, manner of use or application, weather, combination with other products, or crop conditions. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Manufacturer and Seller harmless from any claims relating to such factors.

Seller warrants that this product conforms to the chemical description on the label. EXCEPT FOR THIS WARRANTY, THE PRODUCT IS FURNISHED "AS-IS", AND NEITHER SELLER NOR MANUFACTURER MAKES ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, WITH RESPECT TO THE SELECTION, PURCHASE OR USE OF THIS PRODUCT; SELLER AND MANUFACTURER SPECIFICALLY DISCLAIM ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE BEYOND WHAT IS STATED ON THE LABEL. Buyer and User accept all risks arising from any use of this product, including without limitation, uses contrary to label instructions, or under conditions not reasonably foreseeable to (or beyond the control of) Seller or Manufacturer.

Neither Manufacturer nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. THE EXCLUSIVE REMEDY OF THE BUYER OR USER, AND THE EXCLUSIVE LIABILITY OF MANUFACTURER AND SELLER, FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THIS PRODUCT, OR, AT THE ELECTION OF MANUFACTURER OR SELLER, THE REPLACEMENT OF THE PRODUCT.

These Conditions of Sale and Limitation of Warranty and Liability shall be interpreted, unless otherwise required by the law of the state of purchase, in accordance with the laws of the State of California, excluding its conflicts of laws rules, and may not be amended by any oral or written agreement.

WILBUR-ELLIS® logo and Ideas to Grow With® are registered trademarks of Wilbur-Ellis Company LLC. HERBICIDE™ logo is a trademark of Wilbur-Ellis Company LLC.

Vengeance® is a registered Trademark of Bayer CropScience.
All other trademarks are the property of their respective owners.

**For Chemical Spill, Leak, Fire or Exposure, Call CHEMTREC (800) 424-9300
For Medical Emergencies Only, Call (877) 325-1840**

Distributed by:
WILBUR-ELLIS COMPANY LLC
PO BOX 16458 • FRESNO, CA 93755

Specimen



VENGEANCE® PLUS

SELECTIVE HERBICIDE

ACTIVE INGREDIENTS:	% By Wt.
Dimethylamine Salt of 2-Methyl-4-Chlorophenoxyacetic Acid	46.87%
Triethylamine Salt of 3,5,6-Trichloro-2-Pyridinyloxyacetic Acid**	10.68%
Dimethylamine Salt of (+)-R-2-(2,4-Dichlorophenoxy) propionic Acid***‡	9.12%
OTHER INGREDIENTS:	33.33%
	100.00%

By Isomer specific AOAC Method Equivalent to:

*2-Methyl-4-Chlorophenoxyacetic Acid	38.27%	3,72 lbs./gal.
**3,5,6-Trichloro-2-Pyridinyloxyacetic Acid	7.65%	0,75 lbs./gal.
***(+)-R-2-(2,4-Dichlorophenoxy) propionic Acid	7.65%	0,75 lbs./gal.

‡CONTAINS THE SINGLE ISOMER FORM OF DICHLORPROP-p

For Use By Turf Maintenance Personnel, Landscaping or Commercial Applicators Only

EPA REG. NO. 228-411-2935

EPA EST. NO. 228-IL-001

KEEP OUT OF REACH OF CHILDREN DANGER - PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

For Chemical Spill, Leak, Fire or Exposure, Call CHEMTREC (800) 424-9300

For Medical Emergencies Only, Call (877) 325-1840

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Corrosive. Causes irreversible eye damage. May be fatal if absorbed through skin.
Harmful if swallowed. Do not get in eyes, on skin or on clothing.

FIRST AID	
If in eyes:	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> Call a poison control center or doctor for treatment advice. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person.
HOT LINE NUMBER: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.	
NOTE TO PHYSICIAN: For eye irritation, examination by an ophthalmologist may be indicated. If swallowed, probable mucosal damage may contraindicate the use of gastric lavage.	

ENVIRONMENTAL HAZARDS

This pesticide may be toxic to fish, aquatic invertebrates, aquatic plants and may adversely affect other non-target plants. Drift or runoff may adversely affect non-target plants and may be hazardous to aquatic organisms in water adjacent to treated areas. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Always use original container to store pesticides in a secured warehouse or storage building. Store at temperatures above 32°F. If allowed to freeze, remix before using. This does not alter this product. Containers should be opened in well ventilated areas. Keep container tightly sealed when not in use. Do not stack cardboard cases more than two pallets high. Do not store near open containers of fertilizer, seed or other pesticides. **PESTICIDE DISPOSAL:** Pesticide wastes are acutely hazardous. If container is damaged or if pesticide has leaked, contain all spillage. Absorb and clean up all spilled material with granules or sand. Place in a closed labeled container for proper disposal. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. **CONTAINER DISPOSAL: Nonrefillable Containers 5 Gallons or Less:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke. **Nonrefillable containers larger than 5 gallons:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. If recycling or reconditioning is not available, puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

SEE INSIDE BOOKLET FOR ADDITIONAL PRECAUTIONARY AND STORAGE AND DISPOSAL STATEMENTS.

Distributed by: WILBUR-ELLIS COMPANY LLC
PO BOX 16458 • FRESNO CA 93755

PULL HERE TO OPEN
PRESS TO RESEAL

13152000

F-031918(3)