

MEFENOXAM	GROUP	4	FUNGICIDE
SEDAXANE	GROUP	7	FUNGICIDE

AZOXYSTROBIN	GROUP	11	FUNGICIDE
FLUDIOXONIL	GROUP	12	FUNGICIDE

PULL HERE TO OPEN ►



## Fungicide

A seed treatment product for protection against damage from listed diseases on cotton.

*Active Ingredients:*

Azoxystrobin <sup>1</sup> .....	6.71%
Fludioxonil <sup>2</sup> .....	1.12%
Mefenoxam <sup>3</sup> .....	6.71%
Sedaxane <sup>4</sup> .....	3.13%

<i>Other Ingredients:</i>	82.33%
---------------------------	--------

<i>Total:</i>	100.00%
---------------	---------

<sup>1</sup>CAS No. 131860-33-8

<sup>2</sup>CAS No. 131341-86-1

<sup>3</sup>CAS No. 70630-17-0 and 69516-34-3

<sup>4</sup>CAS No. 874967-67-6

Vibrance CST is a flowable concentrate for seed treatment containing 0.63 lb azoxystrobin, 0.10 lb fludioxonil, 0.63 lb mefenoxam and 0.29 lb sedaxane per gallon.

**KEEP OUT OF REACH OF CHILDREN**

## CAUTION

See additional precautionary statements and directions for use in booklet.

EPA Reg. No. 100-1589

EPA Est. 100-NE-001

SCP 1589A-L1A 0222

4155897

**14.25 Gallons**

Net Contents



<b>FIRST AID</b>	
<b>If swallowed</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to by a poison control center or doctor.</li> <li>• Do not give anything to an unconscious person.</li> </ul>
<b>If on skin</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
<b>HOTLINE NUMBER</b> For 24-Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident), Call <b>1-800-888-8372</b>	

## PRECAUTIONARY STATEMENTS

### Hazards to Humans and Domestic Animals

#### CAUTION

Harmful if swallowed or absorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

#### PERSONAL PROTECTIVE EQUIPMENT (PPE)

##### Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of barrier laminate, butyl rubber  $\geq 14$  mils, neoprene rubber  $\geq 14$  mils, nitrile rubber  $\geq 14$  mils, polyvinyl chloride (PVC)  $\geq 14$  mils, or Viton™  $\geq 14$  mils
- Shoes plus socks

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### ENGINEERING CONTROLS

When handlers use closed systems in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240(d)(4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS.

*continued...*

## **PRECAUTIONARY STATEMENTS (continued)**

### **USER SAFETY RECOMMENDATIONS**

#### **Users should:**

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### **ENVIRONMENTAL HAZARDS**

This pesticide is toxic to fish, aquatic invertebrates, oysters and shrimp. Do not contaminate water when disposing of equipment washwater or rinsate.

#### **Groundwater Advisory**

Azoxystrobin, Mefenoxam and a degradate of Azoxystrobin are known to leach through soil into groundwater under certain conditions as a result of label use. Fludioxonil has properties and characteristics associated with chemicals detected in groundwater. These chemicals may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

#### **Physical or Chemical Hazards**

Do not mix or allow coming in contact with oxidizing agent. Hazardous chemical reaction may occur.

## **CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY**

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, LLC or Seller. To the extent permitted by applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of the product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and (2) Buyer and User assume the risk of any such use. TO THE EXTENT PERMITTED BY APPLICABLE LAW, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.

To the extent permitted by applicable law, in no event shall SYNGENTA be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

## **DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Use is permitted in commercial seed treatment facilities. Use is also permitted on-farm, except in the state of California. In the state of California, use is restricted to commercial seed treatment facilities only. Do not use for at-plant applications (e.g. hopper box, planter box, etc.). This product is to be applied as a water-based slurry through standard slurry- or mist-type commercial seed treatment equipment.

Store between a range of 32°F to 100°F.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Treatment of highly mechanically scarred or damaged seed or seed known to be of low vigor and poor quality may result in reduced germination and/or reduction of seed and seedling vigor. Treat a quantity of seed using equipment similar to that planned for treating the total seed lot. Prior to treatment, conduct germination tests on a portion of seed before committing the total seed lot to a selected seed treatment.

Due to seed quality, crop or variety sensitivity, and seed storage conditions beyond the control of Syngenta, no claims are made to guarantee the germination of carry-over seed or propagating material for all crop seed when treated with Vibrance CST.

**Maximum usage when applying both metalaxyl- and mefenoxam-containing products to the same crop within the same season:** Do not apply more than the maximum yearly total application rate for the active ingredient as stated on the label of the product containing the lowest yearly total on that crop.

**FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR DISEASE CONTROL, AND/OR ILLEGAL RESIDUES.**

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.** Exception: If the seed is treated with the product and the treated seed is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves made of barrier laminate, butyl rubber  $\geq 14$  mils, neoprene rubber  $\geq 14$  mils, nitrile rubber  $\geq 14$  mils, polyvinyl chloride (PVC)  $\geq 14$  mils, or Viton  $\geq 14$  mils
- Shoes plus socks

### PRODUCT INFORMATION

Vibrance CST is a broad-spectrum seed treatment fungicide containing four active ingredients: Azoxystrobin, Fludioxonil, Mefenoxam, and Sedaxane. All four of these active ingredients have a different mode of action against fungal pathogens. Vibrance CST protects cotton against damage from the Seedling Disease Complex: Damping-off, Seedling Blight, Seed Rot, and Root Rot caused by *Fusarium* species, *Pythium* species, and *Rhizoctonia* species.

Azoxystrobin is active against *Rhizoctonia* species causing seedling damping-off and *Pythium* species causing seedling blight in cotton.

Fludioxonil is active against *Fusarium* and *Rhizoctonia* species in cotton.

Mefenoxam is active against *Pythium* species causing damping-off and seed rot in cotton.

Sedaxane is active against *Rhizoctonia* species causing seed decay, seedling blight and damping-off in cotton.

## RESISTANCE MANAGEMENT

MEFENOXAM	GROUP	4	FUNGICIDE
SEDAXANE	GROUP	7	FUNGICIDE
AZOXYSTROBIN	GROUP	11	FUNGICIDE
FLUDIOXONIL	GROUP	12	FUNGICIDE

For resistance management, please note that Vibrance CST contains Group 4/mefenoxam, Group 7/sedaxane, Group 11/azoxystrobin and Group 12/fludioxonil fungicides. Any fungal population may contain individuals naturally resistant to Vibrance CST and other Group 4, Group 7, Group 11 or Group 12 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

Mefenoxam belongs to the phenylamide class of chemistry which interferes with fungal RNA synthesis. Sedaxane is a succinate dehydrogenase inhibitor (SDHI) and belongs to the carboxamide class of chemistry which disrupts cellular respiration and energy generation. Azoxystrobin belongs to the strobilurin class of chemistry which disrupts cellular respiration and energy generation. Fludioxonil belongs to the phenylpyrrole class of chemistry which interferes with osmotic signal transduction.

To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of Vibrance CST or other Group 4, Group 7, Group 11, or Group 12 fungicides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicides from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
- For further information or to report suspected resistance contact Syngenta at 1-866-SYNGENT(A) (866-796-4368). You can also contact your pesticide distributor or university extension specialist to report resistance.

Syngenta encourages responsible product stewardship to ensure effective long term control of the fungal diseases on this label.

### For additional information on Fungicide Resistance Management:

- Contact Syngenta representatives at 1-800-334-9481.
- Contact your local extension specialist or certified crop advisor.
- Visit the Fungicide Resistance Action Committee (FRAC) on the web at: <http://www.frac.info>.

## MIXING PROCEDURES

**Important:** Always re-circulate Vibrance CST thoroughly before using.

Follow the manufacturer's application instructions for the seed treatment equipment being used.

Apply Vibrance CST as a water-based slurry utilizing standard slurry seed treatment equipment which provides uniform seed coverage. Uneven or incomplete seed coverage may not give the desired level of disease control. Thoroughly mix the specified amount of Vibrance CST into the required amount of water or liquid inoculant for the slurry treater and dilution rate to be used.

Certain crops require addition of inoculants when the seed is treated or planted. Vibrance CST is compatible with several liquid inoculant products. Consult the maker of the inoculant product and a Syngenta representative for directions before applying Vibrance CST with inoculants.

The total application volume must be sufficient to provide desired level of coverage. Dilution is typically done with water or liquid inoculants.

Continuous agitation or mixing of the slurry mixture is necessary to prevent settling out of the solution. Clean out any unused product from the treater after treating or maintain constant agitation if the leftover slurry will be maintained overnight.

- Use an EPA-approved dye or colorant that imparts an unnatural color to the seed as required in 40 CFR 153.155(c).
- Allow seed to dry before bagging.

## SEED CONTAINER LABEL REQUIREMENTS

The Federal Seed Act requires that containers of treated seeds shall be labeled with the following statements:

- This seed has been treated with Azoxystrobin, Fludioxonil, Mefenoxam, and Sedaxane fungicides.
- Do not use for feed, food, or oil purposes.

In addition, the U.S. Environmental Protection Agency requires the following statements on containers of seeds treated with Vibrance CST:

- **Groundwater Advisory:** Azoxystrobin, Mefenoxam and a degradate of Azoxystrobin are known to leach through soil into groundwater under certain conditions as a result of label use. Fludioxonil has properties and characteristics associated with chemicals detected in groundwater. These chemicals may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.
- Store treated seed away from feeds and foodstuffs.
- Do not allow children, pets, or livestock to have access to treated seed.
- Wear long-sleeved shirt, long pants and chemical-resistant gloves when handling treated seed.
- Treated seed must be planted into the soil at a depth greater than 1 inch.
- Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading and planting.

- Dispose of all excess treated seed. Leftover treated seed may be double sown around the headland or buried away from water sources in accordance with local requirements.
- Do not contaminate water bodies when disposing of planting equipment washwaters.
- Dispose of seed packaging in accordance with local requirements.
- Excess treated seed may be used for ethanol production only if:
  - (1) By-products are not used for livestock feed, and
  - (2) No measurable residues of pesticide remain in the ethanol by-products that are used in agronomic practice.
- In the event of crop failure or harvest of a crop grown from seed treated with Vibrance CST, crops may be replanted according to the following schedule:

#### Plantback Interval Table

Immediate Plantback
Cereals: Barley, Corn (Field, Pop, Sweet), Oats, Rye, Sorghum, Triticale, and Wheat
Cotton
Dried Shelled Pea and Bean Crop Subgroup 6C
Potato
Rapeseed (Including Canola) Crop Subgroup 20A
Soybean
Sugarbeet

- For any other crops the minimum plantback interval is 30 days from the date seeds treated with Vibrance CST were planted.
- The maximum application rate is not to exceed 0.00018 lb ai/seed of azoxystrobin.
- **DO NOT** apply more than 0.01 lb ai/Acre/calendar year (4.7 g ai/Acre/calendar year) of mefenoxam as a seed treatment on Cotton.
- **DO NOT** apply more than 0.125 lb ai/Acre/calendar year (56.7 g ai/Acre/calendar year) of mefenoxam regardless of type of application (seed treatment, soil or foliar) on Cotton.



## CROP USE DIRECTIONS

When applied according to the **Vibrance CST Rate Table**, Vibrance CST provides early season protection against the diseases listed in the table below.

**Vibrance CST Rate Table**

Crop	Protection Against the Following Diseases <sup>1</sup>	Use Rate (fl oz/100 lb)	Use Rate (Total mg ai / seed) <sup>2</sup>	Use Rate (g ai/100 kg)
Cotton	Seedling Disease Complex: Damping-off, Seedling Blight, Seed Rot, and Root Rot caused by <i>Fusarium</i> species, <i>Pythium</i> species, and/or <i>Rhizoctonia</i> species	3.06 – 4.08	0.040 – 0.053	Azoxystrobin: 15 – 20 Fludioxonil: 2.5 – 3.34 Mefenoxam: 15 – 20 Sedaxane: 7 – 9.33

<sup>1</sup>For protection against damage from Black Root Rot, *Thielaviopsis basicola*, use Vibrance CST in combination with products containing the active ingredient myclobutanil. Use labeled rates for application as a seed treatment.

<sup>2</sup>Based on an average seed weight of 4,500 cotton seeds/lb.

### RESTRICTIONS:

- The maximum application rate for cotton is not to exceed 0.00018 lb ai/seed
- **DO NOT** apply more than 0.01 lb ai/Acre/calendar year (4.7 g ai/Acre/calendar year) of mefenoxam as a seed treatment on Cotton.
- **DO NOT** apply more than 0.125 lb ai/Acre/calendar year (56.7 g ai/Acre/calendar year) of mefenoxam regardless of type of application (seed treatment, soil or foliar) on Cotton.

## **STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

### **Pesticide Storage**

Store in the original container and only in a cool, dry, secure place. Store between a range of 32°F to 100°F.

### **Pesticide Disposal**

Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste Representative at the nearest EPA regional office for guidance.

### **Container Handling (less than or equal to 5 gallons)**

**Non-refillable container.** Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

### **Container Handling (greater than 5 gallons)**



**Non-refillable container.** Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

### **Container Handling (greater than 5 gallons)**

**Refillable container.** Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes.

**CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER.**

Vibrance®, the ALLIANCE FRAME   
the SYNGENTA Logo and the PURPOSE ICON   
are Trademarks of a Syngenta Group Company

Viton™ is a trademark of The Chemours Company FC, LLC

©2022 Syngenta

For non-emergency (e.g., current product information), call  
Syngenta Crop Protection at 1-800-334-9481

Manufactured for:  
Syngenta Crop Protection, LLC  
P.O. Box 18300  
Greensboro, North Carolina 27419-8300

**SCP 1589A-L1A 0222**  
**4155897**

MEFENOXAM	GROUP	4	FUNGICIDE
SEDAXANE	GROUP	7	FUNGICIDE
AZOXYSTROBIN	GROUP	11	FUNGICIDE
FLUDIOXONIL	GROUP	12	FUNGICIDE



## Fungicide

A seed treatment product for protection against damage from listed diseases on cotton.

Active Ingredients:	
Azoxystrobin <sup>1</sup> .....	6.71%
Fludioxonil <sup>2</sup> .....	1.12%
Mefenoxam <sup>3</sup> .....	6.71%
Sedaxane <sup>4</sup> .....	3.13%
Other Ingredients:	82.33%
Total:	100.00%

<sup>1</sup>CAS No. 131860-33-8

<sup>2</sup>CAS No. 131341-86-1

<sup>3</sup>CAS No. 70630-17-0 and 69516-34-3

<sup>4</sup>CAS No. 874967-67-6

Vibrance CST is a flowable concentrate for seed treatment containing 0.63 lb azoxystrobin, 0.10 lb fludioxonil, 0.63 lb mefenoxam and 0.29 lb sedaxane per gallon.

EPA Reg. No. 100-1589

EPA Est. 100-NE-001

Vibrance® and the SYNGENTA Logo are Trademarks of a Syngenta Group Company

©2022 Syngenta

Manufactured for:  
Syngenta Crop Protection, LLC  
P.O. Box 18300  
Greensboro, North Carolina 27419-8300

SCP 1589A-L1A 0222  
4155897

**14.25 Gallons**  
Net Contents

## KEEP OUT OF REACH OF CHILDREN CAUTION

See additional precautionary statements and directions for use in booklet.

### FIRST AID

**If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything to an unconscious person. **If on skin:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

**HOTLINE NUMBER:** For 24-Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident), Call 1-800-888-8372.

### PRECAUTIONARY STATEMENTS

#### Hazards to Humans and Domestic Animals

#### CAUTION

Harmful if swallowed or absorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

**ENVIRONMENTAL HAZARDS:** This pesticide is toxic to fish, aquatic invertebrates, oysters and shrimp. Do not contaminate water when disposing of equipment washwater or rinsate.

**Groundwater Advisory:** Azoxystrobin, Mefenoxam and a degradate of Azoxystrobin are known to leach through soil into groundwater under certain conditions as a result of label use. Fludioxonil has properties and characteristics associated with chemicals detected in groundwater. These chemicals may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

**Physical or Chemical Hazards:** Do not mix or allow coming in contact with oxidizing agent. Hazardous chemical reaction may occur.

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Store in the original container and only in a cool, dry, secure place. Store between a range of 32°F to 100°F.

**Pesticide Disposal:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste Representative at the nearest EPA regional office for guidance.

#### Container Handling

**Non-refillable container.** Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes.

**CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER.**

