

GROUP 3 FUNGICIDE

PEEL HERE  
TO OPEN

# Viticure®

*fungicide*

Agricultural Fungicide

Active Ingredient:	(% by weight)
Triflumizole [1-[1-((4-chloro-2-(trifluoromethyl) phenyl) imino)-2-propoxyethyl]-1H-imidazole]. . . . .	42.14%
Other Ingredients: . . . . .	57.86%
Total: . . . . .	100.00%

Contains 4 lb triflumizole per gallon.

## KEEP OUT OF REACH OF CHILDREN CAUTION

See inside booklet for additional Precautionary Statements, and Directions for Use  
For Product Use Information Call 1-866-761-9397

EPA Reg. No. 400-518

(BD) EPA Est. No. 037429-GA-002

(CJ) EPA Est. No. 070815-GA-001

The EPA Establishment Number is identified by the circled letters that match the first two letters in the batch number.  
004/031014a  
522297-B

Manufactured by:  
MacDermid Agricultural Solutions, Inc.  
245 Freight Street  
Waterbury, CT 06702

NET CONTENTS: 1 GALLON

#### FIRST AID

**IF SWALLOWED**

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by the poison control center or doctor.
- Do not give anything to an unconscious person.

**EMERGENCY ASSISTANCE (Human health):** Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

**24-HOUR EMERGENCY PHONE 1-866-928-0789 or +1-215-207-0061**

#### PRECAUTIONARY STATEMENTS

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION:** Harmful if swallowed. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or the toilet.

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category A on an EPA chemical resistance category selection chart.

**Applicators and Other Handlers Must Wear:** A long-sleeved shirt & long pants; chemical-resistant gloves made of any waterproof material; shoes plus socks.

**USER SAFETY REQUIREMENTS**

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

**ENGINEERING CONTROLS**

When handlers use closed systems or enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

**USER SAFETY RECOMMENDATIONS**

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling the product. Wash thoroughly and change into clean clothing.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. DO NOT contaminate water when cleaning equipment or disposing of equipment washwater or rinsate. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas.

**PHYSICAL or CHEMICAL HAZARDS**

Do not use or store near heat or open flame.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read entire label directions before using.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls; chemical-resistant gloves made of any waterproof material; and shoes plus socks.

#### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in a cool, dry, secure location.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL** - Nonrefillable container. Do not reuse or refill this container.

**For containers equal to or less than 5 gallons in size:** Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

**If pressure rinsing:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

**Then** offer container for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, by incineration or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**Recycling:** Once cleaned, some agricultural plastic pesticide containers can be taken to a container collection site or picked up for recycling. To find the nearest site, contact your chemical dealer or manufacturer or contact the Ag Container Recycling Council (ACRC) at 1-877-952-2272 (toll free) or [www.acrecycle.org](http://www.acrecycle.org).

#### PRODUCT INFORMATION

VITICURE® is a broad-spectrum foliar fungicide for control of certain diseases of grapes. VITICURE is active as a protectant and as an eradicant by preventing disease symptoms after infection has occurred. Its anti-sporulant activity reduces spores after lesions become visible.

Read the entire label and directions before using VITICURE.

### APPLICATION INSTRUCTIONS

VITICURE can be applied with dilute or concentrate ground equipment. Equipment should be frequently checked for calibration. Apply VITICURE in sufficient water to ensure complete and thorough coverage of foliage, bloom and fruit.

Use higher specified application rates on more susceptible varieties and/or under higher disease pressure.

Consult your State Agricultural Experimental Station or Extension Service Specialist for guidance on selection of specific dosage rates and timing of application appropriate to your specific crop and area conditions. State regulations may contain additional restrictions or requirements.

### PRODUCT RESTRICTIONS

Do not allow livestock to graze on treated orchard ground cover.

Do not apply this product through any type of irrigation system.

**RESISTANCE MANAGEMENT:** Populations of fungal pathogens are known to develop resistance to fungicides with the same mode of action when used repeatedly. Thus, MacDermid Agricultural Solutions encourages responsible resistance management to ensure an effective long-term control of diseases. Resistance management strategies include alternating and/or tank mixing with other labeled fungicides having different modes of action or limiting the total number of applications per season.

VITICURE belongs to the demethylation inhibitor (DMI) group of fungicides classified as Group 3. Do not make more than two sequential applications of VITICURE before alternating to a registered fungicide with a different mode of action. Consult your local or state agricultural authorities for resistance management strategies that are appropriate for your disease management program.

**RESTRICTIONS ON ROTATIONAL CROPS:** Do not plant leafy or fruiting vegetables within 30 days after application. Do not plant root and tuber vegetables within 60 days after application. Do not plant cotton, small cereal grains and all other crops not registered on the label for VITICURE, within one year after application.

**GRAPES including FRUIT, SMALL FRUIT VINE CLIMBING SUBGROUP 13-7F, EXCEPT FUZZY KIWIFRUIT**

Use VITICURE as a protectant for control of powdery mildew. Start applications before bloom and continue on a 14 to 21 day interval at 4 to 8 fl oz/acre. For the most susceptible varieties, use 6 to 8 fl oz/acre and/or shorter spray intervals. When disease pressure is low, use 4 to 6 fl oz/acre and/or the spray interval can be extended up to 21 days. Select spray volume, nozzle adjustments and ground speed to ensure complete coverage of fruit clusters and foliage. Use a minimum of 50 gallons of water per acre for early season applications. Use enough volume to ensure thorough coverage as the amount of foliage increases. Do not apply within 7 days of harvest. Do not exceed 32 fl oz VITICURE per acre per season. Do not apply by air. Sulfur at specified use rates can be applied before, with, and/or between VITICURE applications.

<b>CROP</b>	<b>DISEASE</b>	<b>Rate of VITICURE fl oz/acre</b>	<b>COMMENTS</b>
<b>Amur River Grape Grape Gooseberry Maypop Schisandra berry cultivars, varieties, and/or hybrids of these</b>	<b>Powdery Mildew</b> <i>(Uncinula necator)</i>	4 to 8	Start applications before bloom and continue on a 14 to 21 day interval. Use higher rates and/or shorter intervals on susceptible varieties or under heavy disease pressure. VITICURE applied in a powdery mildew spray schedule, will enhance the activity of registered fungicides used for the control of Botrytis bunch rot. Applications must be made on a 14-day spray schedule for suppression of Botrytis bunch rot. Restricted Entry Interval (REI): The REI for girdling grapes is 24 hours (1-day).

**IMPORTANT NOTICE**—To the extent consistent with applicable law, MacDermid Agricultural Solutions, Inc. warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under conditions not reasonably foreseeable to MacDermid Agricultural Solutions, Inc., and to the extent consistent with applicable law, the buyer assumes the risk of any such use.

VITICURE® is a registered trademark of MacDermid Agricultural Solutions, Inc.

©2016 MacDermid Agricultural Solutions, Inc.

**This page was intentionally left blank.**

**This page was intentionally left blank.**

GROUP 3 FUNGICIDE

Viticulture  
fungicide

Agricultural Fungicide

Active Ingredient:	(% by weight)
Triflumizole [1-[1-(4-chloro-2-(trifluoromethyl) phenyl) imino]-2 propoxyethyl]-1H-imidazole]	42.14%
Other Ingredients:	57.86%
Total:	100.00%

Contains 4 lb triflumizole per gallon.

## KEEP OUT OF REACH OF CHILDREN CAUTION

See inside booklet for additional Precautionary Statements, and Directions for Use  
For Product Use Information Call 1-866-761-9397

FIRST AID	
<b>IF SWALLOWED</b>	<ul style="list-style-type: none"> <li>Call a poison control center or doctor immediately for treatment advice.</li> <li>Have person sip a glass of water if able to swallow.</li> <li>Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>Do not give anything to an unconscious person.</li> </ul>
<b>EMERGENCY ASSISTANCE (Human health):</b> Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
<b>24-HOUR EMERGENCY PHONE 1-866-928-0789 or +1-215-207-0061</b>	

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Harmful if swallowed. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or the toilet.

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in a cool, dry, secure location.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL** - Nonrefillable container. Do not reuse or refill this container.

**For containers equal to or less than 5 gallons in size:** Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

**If pressure rinsing:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

**Then** offer container for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, by incineration or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**Recycling:** Once cleaned, some agricultural plastic pesticide containers can be taken to a container collection site or picked up for recycling. To find the nearest site, contact your chemical dealer or manufacturer or contact the Ag Container Recycling Council (ACRC) at 1-877-952-2272 (toll free) or [www.acrcycle.org](http://www.acrcycle.org).

VITICURE® is a registered trademark of MacDermid Agricultural Solutions, Inc.

Manufactured for:  
MacDermid Agricultural Solutions, Inc.  
245 Freight Street  
Waterbury, CT 06702

EPA Reg. No. 400-518

(B0) EPA Est. No. 037429-GA-002

(C1) EPA Est. No. 070815-GA-001

The EPA Establishment Number is identified by the circled letters that match the first two letters in the batch number.

004/031014

522297-B

**NET CONTENTS: 1 GALLON**