

(Booklet/base – 2.5 Gallon)
 (Booklets - 30/265 Gallon)

THIAMETHOXAM	GROUP	4A	INSECTICIDE
DIFENOCONAZOLE	GROUP	3	FUNGICIDE
MEFENOXAM	GROUP	4	FUNGICIDE
SEDAXANE	GROUP	7	FUNGICIDE
FLUDIOXONIL	GROUP	12	FUNGICIDE

Warden® Cereals WRII

A seed treatment product for protection against damage from certain insects and diseases of small grain cereals.

Active Ingredient:

Thiamethoxam ¹	5.75%
Difenoconazole ² :	3.45%
Mefenoxam ³ :	0.86%
Fludioxonil ⁴ :.....	0.72%
Sedaxane ⁵	1.44%
Other Ingredients:	87.78%
Total:	100.00%

¹CAS No. 153719-23-4

²CAS No. 119446-68-3

³CAS Nos. 70630-17-0 and 69516-34-3

⁴CAS No. 131341-86-1

⁵CAS No. 874967-67-6

Warden Cereals WRII contains 0.51 lbs. thiamethoxam; 0.31 lbs. difenoconazole; 0.08 lbs. mefenoxam; 0.06 lbs. fludioxonil and 0.13 lbs. sedaxane per gallon.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

See additional precautionary statements and directions for use in booklet [on label].

EPA Reg. No. 100-1527 -1381

EPA Est. No. 100-NE-001

SCPSR-WIN-1527A-L2B 0919
 SCPSR-WIN-1527A-L2B 0919
 SCPSR-WIN-1527A-L5B 0919

[Product ID 57208 - 30 gal Bulk]
 [Product ID 57471- 265 gal Bulk]
 [Product ID 58305 - 2x2.5GL]

_____ [Bulk]
Net Contents

1/0920/9

Distributed by:
Winfield Solutions, LLC
P.O. Box 64589
St. Paul, MN 55164-0589

FIRST AID
Have the product container with you when calling a poison control center or doctor, or going for treatment.
HOT LINE NUMBER
For 24 Hour Medical Emergency Assistance (Human or Animal) Call 1-877-424-7452 or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident), Call 1-800-424-9300

PRECAUTIONARY STATEMENTS

Personal Protective Equipment (PPE)

Mixers, loaders, applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves made of polyvinyl chloride [PVC] ≥14 mils, nitrile rubber ≥14 mils, butyl rubber ≥14 mils, barrier laminate, neoprene rubber ≥14 mils, natural rubber ≥14 mils, polyethylene, or Viton® ≥14 mils

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statements

When handlers use closed systems in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticides get inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

This product is toxic to wildlife, freshwater and estuarine/marine fish and highly toxic to aquatic invertebrates. Runoff may be hazardous to aquatic organisms in neighboring areas. Exposed treated seed may be hazardous to wildlife. Do not contaminate water when disposing of equipment washwater or rinsate.

Pollinator Precautions

Thiamethoxam is highly toxic to bees, and effects are possible as a result of exposure to translocated residues in blooming crops.

If treated seed is spilled outdoors or in areas accessible to birds, promptly clean up or bury to prevent ingestion.

Groundwater Advisory

Mefenoxam is known to leach through soil into ground water under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

Thiamethoxam has properties and characteristics associated with chemicals detected in ground water. This chemical may leach into the ground water if used in areas where soils are permeable, particularly where the water table is shallow.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER AND USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE

PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USER OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THE NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the seller, and purchase price will be refunded.

DIRECTIONS FOR USE

For commercial or on-farm use.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

FAILURE TO FOLLOW DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR DISEASE CONTROL, AND/OR ILLEGAL RESIDUES.

Treatment of highly mechanically scarred or damaged seed, or seed known to be of low vigor and poor quality, except for the purpose of curative control of existing disease pests, may result in reduced germination and/or reduction of seed and seedling vigor. Treat using equipment similar to that planned for treating the total seed lot. Conduct germination tests on a small portion of seed before committing the total seed lot to a selected seed treatment. Due to seed quality and seed storage conditions beyond the control of WINFIELD SOLUTIONS, LLC, no claims are made to guarantee the germination of carry-over treated seed.

Maximum usage when applying both metalaxyl- and mefenoxam-containing products to the same crop within the same season: Do not apply more than the maximum yearly total application rate for the active ingredient as stated on the label of the product containing the lowest yearly total on that crop.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notifications to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours. Exception: If the seed is treated with the product and the treated seed is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves made of polyvinyl chloride [PVC] ≥14 mils, nitrile rubber ≥14 mils, butyl rubber ≥14 mils, barrier laminate, neoprene rubber ≥14 mils, natural rubber ≥14 mils, polyethylene, or Viton® ≥14 mils
- Shoes plus socks

USE RESTRICTIONS

- Do not use for feed, food, or oil purposes.
- Store away from feeds and foodstuffs.
- Wear long-sleeved shirt, long pants and chemical resistant gloves when handling treated seed.
- Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading.
- Do not allow children, pets, or livestock to have access to treated seed.
- Dispose of all excess treated seed. Leftover treated seed may be doublesown around the headland or buried away from water sources in accordance with local requirements.
- Do not contaminate water bodies when disposing of planting equipment wash waters.
- Dispose of seed packaging in accordance with local requirements.
- Treated seed must be planted into the soil at a depth greater than 0.5 inch.
- For seed treated with Warden Cereals WR II, do not graze or feed livestock on treated areas for 45 days after planting.
- In the event of a crop failure or harvest of a crop grown from Warden Cereals WR II treated seed, the field may be replanted immediately to canola, soybean, barley, oat, rye triticale and wheat, sweet corn, and chickpea.
- Alfalfa, *Brassica* (cole) leafy vegetables, buckwheat, corn (except sweet corn), pearl millet, proso millet, popcorn, rice (dry-seeded), sorghum, teosinte, wild rice, cotton, cucurbit vegetables, dry bulb onions, fruiting vegetables, leafy vegetables, legume vegetables (except chickpea), mint (peppermint and spearmint), oil seed crops (black mustard seed, borage seed, crambe seed, field mustard seed, flax seed, Indian mustard seed, Indian rapeseed seed, peanuts, rapeseed seed, and safflower seed), root vegetables, strawberry, sunflowers, tobacco, and tuberous and corm vegetables may be planted 30 days from the date the Warden Cereals WR II treated seed was planted.
- For any other crop, the minimum plant back interval is 120 days from the date the Warden Cereals WR II treated seed was planted. A cover crop other than the crops listed above that is planted for erosion control or soil improvement may be planted sooner than the 120 day interval; however, the crop may not be grazed or harvested for food or feed.

- Do not make any soil or foliar application of products containing thiamethoxam to crops grown from seed treated with Warden Cereals WRII.
- Do not use at a rate that will result in more than 0.05 lb thiamethoxam per acre (22.7 grams ai/A) per year as a seed treatment application.
- Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in the ethanol by-products that are used in agronomic practice.

PRODUCT INFORMATION

Warden Cereals WRII is a seed treatment product containing one insecticide active ingredient (thiamethoxam) and four fungicide active ingredients (difenoconazole, mefenoxam, fludioxonil and sedaxane). Warden Cereals WRII seed treatment protects against damage from certain early-season insects and certain seed- and soil-borne diseases of cereal crops. Difenoconazole provides protection against several seed, seedling and certain foliar disease including bunts and smuts. Mefenoxam protects against damping-off caused by *Pythium*. Fludioxonil provides protection against certain soil-borne and seed-borne diseases and is active against *Fusarium*, *Rhizoctonia*, and weakly pathogenic fungi such as *Aspergillus* and *Penicillium*. Sedaxane protects against *Rhizoctonia* and loose smuts (*Ustilago*).

Thiamethoxam is a systemic seed treatment insecticide belonging to the neoticotinoid class of chemistry. Thiamethoxam in Warden Cereals WRII protects against aphids, wireworm and European chafer.

Warden Cereals WRII may be applied in tank mixes, or sequentially with other EPA-registered seed treatment pesticide products. If using Warden Cereals WRII in a tank mixture with other seed treatment products, observe all directions for use, crops/sites, use rates, dilution ratios, precautions, and limitations which appear on the tank-mix partner label(s). Do not exceed any label dosage and follow the most restrictive label precautions and limitations. This product must not be mixed with any product which prohibits such mixing.

MIXING INSTRUCTIONS

Important: Always re-circulate Warden Cereals WRII thoroughly before using.

Apply Warden Cereals WRII as a water-based slurry utilizing standard slurry seed treatment equipment which provides uniform seed coverage. Uneven or incomplete seed coverage may not give the desired level of insect and disease control.

Consult the manufacturer of the application equipment you plan to employ for suitability for this application and for instructions on operation and calibration of the equipment. Follow the manufacturer application instructions for the seed treatment equipment being used.

Continuous agitation or mixing of the slurry mixture is necessary to prevent settling out of the solution.

Allow seed to dry before bagging.

Follow planter manufacturer specifications for use of talc or other hopper box additives at planting. Seed must be completely dry before adding to planter.

Warden Cereals WRII contains an EPA approved dye/colorant that imparts an unnatural color to the seed as required by the Federal Seed Act.

SEED BAG LABEL REQUIREMENTS

The Federal Seed Act requires that bags containing treated seeds shall be labeled with the following statements:

- This seed has been treated with difenoconazole, mefenoxam, fludioxonil and sedaxane fungicides and thiamethoxam insecticide.
- Do not use for feed, food, or oil purposes.
- User is responsible for ensuring that the seed bag meets all requirements under the Federal Seed Act.

In addition, the U.S. Environmental Protection Agency requires the following statements on bags containing seeds treated with Warden Cereals WRII:

- Pollinator Precaution:
Thiamethoxam is highly toxic to bees, and effects are possible as a result of exposure to translocated residues in blooming crops.
- Ground Water Advisory:
Mefenoxam is known to leach through soil into ground water under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.
Thiamethoxam has properties and characteristics associated with chemicals detected in ground water. This chemical may leach into the ground water if used in areas where soils are permeable, particularly where the water table is shallow.
- Store away from feeds and foodstuffs.
- Wear long-sleeved shirt, long pants and chemical resistant gloves when handling treated seed.
- Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading.
- Do not allow children, pets, or livestock to have access to treated seed.
- Dispose of all excess treated seed. Leftover treated seed may be doublesown around the headland or buried away from water sources in accordance with local requirements.

- Do not contaminate water bodies when disposing of planting equipment wash waters.
- Dispose of seed packaging in accordance with local requirements.
- Treated seed must be planted into the soil at a depth greater than 0.5 inch.
- For seed treated with Warden Cereals WRII, do not graze or feed livestock on treated areas for 45 days after planting.
- In the event of a crop failure or harvest of a crop grown from Warden Cereals WRII treated seed, the field may be replanted immediately to canola, soybean, barley, oat, rye triticale and wheat, sweet corn, and chickpea.
- Alfalfa, *Brassica* (cole) leafy vegetables, buckwheat, corn (except sweet corn), pearl millet, proso millet, popcorn, rice (dry-seeded), sorghum, teosinte, wild rice, cotton, cucurbit vegetables, dry bulb onions, fruiting vegetables, leafy vegetables, legume vegetables (except chickpea), mint (peppermint and spearmint), oil seed crops (black mustard seed, borage seed, crambe seed, field mustard seed, flax seed, Indian mustard seed, Indian rapeseed seed, peanuts, rapeseed seed, and safflower seed), root vegetables, strawberry, sunflowers, tobacco, and tuberous and corm vegetables may be planted 30 days from the date the Warden Cereals WRII treated seed was planted.
- For any other crop, the minimum plant back interval is 120 days from the date the Warden Cereals WRII treated seed was planted. A cover crop other than the crops listed above that is planted for erosion control or soil improvement may be planted sooner than the 120 day interval; however, the crop may not be grazed or harvested for food or feed.
- Do not make any soil or foliar application of products containing thiamethoxam to crops grown from seed treated with Warden Cereals WRII.
- Do not use at a rate that will result in more than 0.05 lb thiamethoxam per acre (22.7 grams ai/A) per year as a seed treatment application.
- This seed has been treated with 0.008 mg ai thiamethoxam per seed assuming an average seed size of 12,000 seeds per pound.
- Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in the ethanol by-products that are used in agronomic practice.

CROP USE PRECAUTIONS

Resistance Management

DIFENOCONAZOLE	GROUP	3	FUNGICIDE
MEFENOXAM	GROUP	4	FUNGICIDE
SEDAXANE	GROUP	7	FUNGICIDE
FLUDIOXONIL	GROUP	12	FUNGICIDE

For resistance management, please note that Warden Cereals WRII contains Group 3/difenoconazole, Group 4/mefenoxam, Group 7/sedaxane, and Group 12/fludioxonil fungicides. Any fungal population may contain individuals naturally resistant to Warden Cereals WRII and other Group 3, Group 4, Group 7, or Group 12 fungicides. A gradual

or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

Difenoconazole belongs to the triazole class of chemistry and is a demethylation inhibitor of sterol biosynthesis (DMI) which disrupts membrane synthesis of the fungal cell. Mefenoxam belongs to the phenylamide class of chemistry which interferes with fungal RNA synthesis. Sedaxane is a succinate dehydrogenase inhibitor (SDHI) and belongs to the carboxamide class of chemistry which disrupts cellular respiration and energy generation. Fludioxonil belongs to the phenylpyrrole class of chemistry which interferes with osmotic signal transduction.

To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of Warden Cereals WR11 or other Group 3, Group 4, Group 7, or Group 12 fungicides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicides from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide application. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
- For further information or to report suspected resistance contact Winfield Solutions, LLC at 1-855-494-6343. You can also contact your pesticide distributor or university extension specialist to report resistance.

Winfield Solutions, LLC encourages responsible product stewardship to ensure effective long term control of the fungal diseases on this label.

THIAMETHOXAM	GROUP	4A	INSECTICIDE
---------------------	--------------	-----------	--------------------

Warden Cereals WR^{II} contains a Group 4A insecticide (thiamethoxam, belonging to the neonicotinoid class of chemistry). Insect biotypes with acquired or inherent resistance to Group 4A insecticides may eventually dominate the insect population if Group 4A insecticides are used repeatedly as the predominant method of control for targeted species. This may result in partial or total loss of control of those species by thiamethoxam or other Group 4A insecticides.

In order to maintain susceptibility to this class of chemistry:

- Avoid using Group 4A insecticides exclusively for season long control of insect species with more than one generation per crop season.
- For insect species with successive or overlapping generations, apply Warden Cereals WR^{II} or other Group 4A insecticides using a “treatment window” approach. A treatment window is a period of time as defined by the stage of crop development and/or the biology of the pests of concern. Within the treatment window, depending on the length of residual activity, there may either be single or consecutive applications (seed treatment, soil, foliar, unless otherwise stated in the Directions for Use) of the Group 4A insecticides. When using Warden Cereals WR^{II} seed treatment in the first treatment window rotate to effective products with a different mode of action in the subsequent treatment window before making additional applications of Group 4A insecticides.
- A treatment window rotation, along with other IPM practices for the crop and use area, is considered an effective strategy for preventing or delaying a pest’s ability to develop resistance to this class of chemistry.
- If resistance is suspected, do not reapply Warden Cereals WR^{II} or any other Group 4A insecticides.

Other Insect Resistance Management (IRM) practices include:

- Incorporating IPM techniques into your insect control program.
- Monitoring treated insect populations for loss of field efficacy.
- Using tank-mixtures or premixes with insecticides from a different target site of action group as long as the involved products are all registered for the same crop outlet and effective rates are applied.

For additional information on Insect Resistance Management:

- Contact your local extension specialist, certified crop advisor and/or product manufacturer for additional insect resistance management recommendations.
- Visit the Insecticide Resistance Action Committee (IRAC) on the web at: <http://www.irac-online.org/>.

Consult your local pest control advisor or extension office for additional methods for preventing resistance development. Winfield Solutions, LLC encourages responsible product stewardship to ensure effective long term control of the fungal and insect pests on this label.

CROP USE DIRECTIONS

Crop	Protection Against the following Insects	Use Rate Fluid Ounces per 100 lb seed
Barley	Aphids ¹ European Chafer Wireworms ¹	5 fluid oz

Crop	Protection Against the Following Diseases	Use Rate Fluid Ounces per 100 lb seed
Barley	General Seed Rots ² Seedling Blight, Root Rot, and Damping-Off caused by seed- and soil-borne <i>Fusarium</i> . or <i>Rhizoctonia</i> Seedling Blight, Root Rot, and Damping-Off caused by soil-borne <i>Pythium</i> Seed-borne <i>Septoria</i> Covered Smut False Loose Smut True Loose Smut	5 fluid oz

Crop	Diseases Suppressed³	Use Rate Fluid Ounces per 100 lb seed
Barley	Common Root Rot (<i>Cochliobolus</i>) <i>Fusarium</i> Crown and Foot Rot Take-All	5 fluid oz

Crop	Protection Against the following Insects	Use Rate Fluid Ounces per 100 lb seed
Oats	Aphids ¹ European Chafer Wireworms ¹	5 fluid oz

Crop	Protection Against the Following Diseases	Use Rate Fluid Ounces per 100 lb seed
Oats	General Seed Rots ² Seedling Blight, Root Rot, and Damping-Off caused by seed- and soil-borne <i>Fusarium</i> or <i>Rhizoctonia</i> Seedling Blight, Root Rot, and Damping-Off caused by soil-borne <i>Pythium</i> Seed-borne <i>Septoria</i> Covered Smut Loose Smut	5 fluid oz

Crop	Diseases Suppressed³	Use Rate Fluid Ounces per 100 lb seed
Oats	Common Root Rot (<i>Cochliobolus</i>)	5 fluid oz

Crop	Protection Against the following Insects	Use Rate Fluid Ounces per 100 lb seed
Rye	Aphids ¹ European Chafer Wireworms ¹	5 fluid oz

Crop	Protection Against the Following Diseases	Use Rate Fluid Ounces per 100 lb seed
Rye	General Seed Rots ² Seedling Blight, Root Rot, and Damping-Off caused by seed- and soil-borne <i>Fusarium</i> or <i>Rhizoctonia</i> Seedling Blight, Root Rot, and Damping-Off caused by soil-borne <i>Pythium</i> Seed-borne <i>Septoria</i> Common Bunt ⁴ Dwarf Bunt ⁴	5 fluid oz

Crop	Diseases Suppressed³	Use Rate Fluid Ounces per 100 lb seed
Rye	Common Root Rot (<i>Cochliobolus</i>) <i>Fusarium</i> Crown and Foot Rot Take-All	5 fluid oz

Crop	Protection Against the following Insects	Use Rate Fluid Ounces per 100 lb seed
Winter Wheat	Aphids ¹ European Chafer Wireworms ¹	5 fluid oz

Crop	Protection Against the Following Diseases	Use Rate Fluid Ounces per 100 lb seed
Winter Wheat and Triticale	General Seed Rots ² Seedling Blight, Root Rot, and Damping-Off caused by seed- and soil-borne <i>Fusarium</i> or <i>Rhizoctonia</i> Seedling Blight, Root Rot, and Damping-Off caused by soil-borne <i>Pythium</i> Seed-borne <i>Septoria</i> <i>Septoria</i> Leaf Blotch ⁵ Common Bunt ⁴ Flag Smut <i>Fusarium</i> Seed Scab Dwarf Bunt ⁴ Karnal Bunt Loose Smut <i>Pythium</i> Damping Off	5 fluid oz

Crop	Diseases Supressed ³	Use Rate Fluid Ounces per 100 lb seed
Winter Wheat and Triticale	Common Root Rot (<i>Cochliobolus</i>) <i>Fusarium</i> Crown and Foot and Root Rot Take-All	5 fluid oz

Crop	Protection Against the following Insects	Use Rate Fluid Ounces per 100 lb seed
Spring Wheat	Aphids ¹ European Chafer Wireworms ¹	5 fluid oz

Crop	Protection Against the Following Diseases	Use Rate Fluid Ounces per 100 lb seed
Spring Wheat	General Seed Rots ² Seedling Blight, Root Rot, and Damping-Off caused by seed- and soil-borne <i>Fusarium</i> or <i>Rhizoctonia</i> Seedling Blight, Root Rot, and Damping-Off caused by soil-borne <i>Pythium</i> Seed-borne <i>Septoria</i> Fusarium Seed Scab Common Bunt ⁴ Karnal Bunt Loose Smut <i>Pythium</i> Damping Off	5 fluid oz

Crop	Diseases Supressed³	Use Rate Fluid Ounces per 100 lb seed
Spring Wheat	Common Root Rot (<i>Cochliobolus</i>) <i>Fusarium</i> Crown and Foot and Root Rot Take-All	5 fluid oz

¹The 5 fluid oz/100 lbs rate of Warden Cereals WR II provides suppression of wireworm activity, aphids and aphid vectored barley yellow dwarf virus. If pressure is moderate or high or control is required, apply up to 0.8 fl. oz. additional Cruiser 5FS.

²Protection against general seed rots. This includes rots caused by *Fusarium*, *Pythium*, *Rhizoctonia*, *Penicillium* and *Aspergillus*.

³Suppression means consistent protection at a level which is not optimal but is still of commercial benefit.

⁴Protects against both seed- and soil-borne bunts (common, dwarf).

⁵Early season foliar disease protection for first 4 weeks after planting. For full season protection, apply a foliar fungicide according to label directions.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

Pesticide Storage

Store in original containers only. Keep container closed when not in use. Do not store near food or feed. In case of spill on floor or paved surfaces, mop and remove to chemical waste storage area until proper disposal can be made if product cannot be used according to the label.

Pesticide Disposal

Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative of the nearest EPA Regional Office for guidance.

Container Handling [less than 5 gallons]

Non-refillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use and disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Container Handling [greater than 5 gallons]

Non-refillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container $\frac{1}{4}$ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Container Handling [greater than 5 gallons]

Refillable container. Refill this container with pesticide only. Do not use this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the

refiller. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER.

FOR CHEMICAL EMERGENCY: Spill, leak, fire, or accident, call CHEMTREC 1-800-424-9300

Warden® is a Trademark of Winfield Solutions, LLC

Viton® trademark of E. I. du Pont de Nemours and Company

©2019 Winfield

Distributed by:
Winfield Solutions, LLC
P.O. Box 64589
St. Paul, MN 55164-0589

[Base Label – 2.5 gal]

THIAMETHOXAM	GROUP	4A	INSECTICIDE
DIFENOCONAZOLE	GROUP	3	FUNGICIDE
MEFENOXAM	GROUP	4	FUNGICIDE
SEDAXANE	GROUP	7	FUNGICIDE
FLUDIOXONIL	GROUP	12	FUNGICIDE

Warden® Cereals WRII

A seed treatment product for protection against damage from certain insects and diseases of small grain cereals.

Active Ingredient:

Thiamethoxam ¹	5.75%
Difenoconazole ² :	3.45%
Mefenoxam ³ :	0.86%
Fludioxonil ⁴ :	0.72%
Sedaxane ⁵	1.44%
Other Ingredients:	87.78%
Total:	100.00%

¹CAS No. 153719-23-4

²CAS No. 119446-68-3

³CAS Nos. 70630-17-0 and 69516-34-3

⁴CAS No. 131341-86-1

⁵CAS No. 874967-67-6

Warden Cereals WRII contains 0.51 lbs. thiamethoxam; 0.31 lbs. difenoconazole; 0.08 lbs. mefenoxam; 0.06 lbs. fludioxonil and 0.13 lbs. sedaxane per gallon.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

See additional precautionary statements and directions for use in booklet [on label].

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling under “Agricultural Use Requirements” in the Directions for Use section for information about this standard.

EPA Reg. No. 100-1527 -1381

EPA Est. No. 100-NE-001

SCPSR-WIN-1527A-L5A 0919

[Product ID 58305 - 2x2.5GL]

PRECAUTIONARY STATEMENTS

CAUTION

FIRST AID
Have the product container with you when calling a poison control center or doctor, or going for treatment.
HOT LINE NUMBER
For 24 Hour Medical Emergency Assistance (Human or Animal) Call 1-877-424-7452 or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident), Call 1-800-424-9300

Environmental Hazards

This product is toxic to wildlife, freshwater and estuarine/marine fish and highly toxic to aquatic invertebrates. Runoff may be hazardous to aquatic organisms in neighboring areas. Exposed treated seed may be hazardous to wildlife. Do not contaminate water when disposing of equipment washwater or rinsate.

Pollinator Precautions

Thiamethoxam is highly toxic to bees, and effects are possible as a result of exposure to translocated residues in blooming crops.

If treated seed is spilled outdoors or in areas accessible to birds, promptly clean up or bury to prevent ingestion.

Groundwater Advisory

Mefenoxam is known to leach through soil into ground water under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

Thiamethoxam has properties and characteristics associated with chemicals detected in ground water. This chemical may leach into the ground water if used in areas where soils are permeable, particularly where the water table is shallow.

STORAGE AND DISPOSAL
Do not contaminate water, food or feed by storage and disposal.
Pesticide Storage
Store in original containers only. Keep container closed when not in use. Do not store near food or feed. In case of spill on floor or paved surfaces, mop and remove to chemical

waste storage area until proper disposal can be made if product cannot be used according to the label.

Pesticide Disposal

Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative of the nearest EPA Regional Office for guidance.

Container Handling [less than 5 gallons]

Non-refillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use and disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER.

FOR CHEMICAL EMERGENCY: Spill, leak, fire, or accident, call CHEMTREC 1-800-424-9300

Warden® is a Trademark of Winfield Solutions, LLC

Viton® trademark of E. I. du Pont de Nemours and Company

©2019 Winfield

Distributed by:
Winfield Solutions, LLC
P.O. Box 64589
St. Paul, MN 55164-0589

SCPSR-WIN-1527A-L5B0919

[Product ID 58305 - 2x2.5GL]

2.5 gallons
Net Contents