Center of Butt Print

APPLICABLE FREIGHT CLASSIFICATION

IN CASE OF SPILL: Sweep up

WARNING!

Burning sulfur emits highly toxic fumes. Sulfur dust

Keep away from heat, sparks and flames. Avoid

creating dust in handling. Keep container closed.

Use with adequate ventilation. Keep from contact

IN CASE OF FIRE: Smother with

fog-like spray, sand, or CO2. Wear NIOSH approved

suspended in air ignites easily.

— Slit Edge

Print Edge

self-contained breathing apparatus. spillage. Place in an appropriate chemical waste container. Flush spill area with water. Observe all local, state, and federal laws and regulations

regarding chemical spill, cleanup, removal and

SULFUR, SOLID

Proper Shipping Name

CAS # 7704-34-9

55FG-HEL-000000

MANUFACTURED FOR HELENA CHEMICAL COMPANY 225 SCHILLING BOULEVARD SUITE 300

AD 041411

Genter of Butt Print

Center of Butt Print

The second of the second of

WETTABLE SULPHUR For Use in Water As An Agricultural Fungicide

INERT INGREDIENTS:

TOTAL: 100%

ACTIVE INGREDIENT:

KEEP OUT OF REACH OF CHILDREN CAUTION

IF SWALLOWED: • Call a poison control center or doctor immediately for treatment advise.

IF ON SKIN OR

EPA REG. No. 5905-289

MANUFACTURED FOR

EPA EST. No. 51036-GA-1

HELENA CHEMICAL COMPANY

PRECAUTIONARY STATEMENTS

Long-sleeved shirt and long pants

USER SAFETY RECOMMENDATIONS

AGRICULTURAL USE REQUIREMENTS

Shoes plus socks

DIRECTIONS FOR USE

Shoes plus socks

waste disposal facility

2.25" Slit

CHEMIGATION PROHIBITION

STORAGE AND DISPOSAL

Do not apply this product through any type of irrigation system.

COLLIERVILLE, TENNESSEE 38017

PERSONAL PROTECTIVE EQUIPMENT (PPE)

225 SCHILLING BOULEVARD, SUITE 300

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

PPE requirements may be reduced or modified as specified in the WPS.

Have person sip a glass of water if able to swallow.

Do not give anything to an unconscious or convulsing person.

Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

Causes moderate eye irritation. Harmful if absorbed through skin. Avoid contact with eyes, skin or clothing

• Chemical resistant gloves made of a waterproof material (such as Nitrile, Neoprene, and/or Barrier Laminate)

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

high water mark. Do not contaminate water when disposing of rinsate or equipment washwater.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed

• Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this

product in a way that will contact workers or other persons, either directly or through drift. Only protected

handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Use this product only in accordance with its labeling and with the Workers Protection Standard, 40 CFR part

170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries

and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination

notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The

requirements in this box only apply to uses of this product that are covered by the Workers Protection Standard.

• Chemical resistant gloves made of a waterproof material (such as Nitrile, Neoprene, and/or Barrier Laminate)

Do not enter or allow worker entry into treated areas during the restricted interval (REI) of 24 hours.

involves contact with anything that has been treated, such as plants, soil, or water, is:

local authorities, by burning. If burned stay out of smoke. Offer for recycling, if available.

PPE required for early entry to treated areas is permitted under the Worker Protection Standard and that

PESTICIDE STORAGE: Store in original container only. In case of leak or spill, contain material and dispose as

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved

CONTAINER DISPOSAL: Non-refillable container. Do not reuse or refill this container. Completely empty bag into

application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and

Hold eyelid open and rinse slowly and gently with water 15-20 minutes.

Call a poison control center or doctor immediately for treatment advice.

 Remove contact lenses, if present, after the first 5 minutes, then continue rinsing • Call a poison control center or doctor immediately for treatment advice.

• Do not induce vomiting unless told to do so by a poison control center or doctor.

AD 041411

NET WEIGHT:

COLLIERVILLE, TENNESSEE 38017

WHEN GROWING CROPS FOR PROCESSING, CONSULT PROCESSOR BEFORE APPLYING SULPHUR. DO NOT USE WITHIN FOUR (4) WEEKS OF AN OIL SPRAYING. APPLICATIONS RATES Pour needed amount into tank under agitation. If necessary, add a few drops of oil to avoid foaming. **DIRECTIONS FOR DILUTION: DILUTE APPLICATION:** Ground application- Apply specified rate in 20 to 100 gallons of water per acre. Orchard Applicaton - Apply specified rate in 100 to 1500 gallons of water per acre. CONCENTRATE APPLICATION: Ground Application- Apply specified rate in not less than 5 gallons of water per acre. Orchard Application - Apply specified rate in 20 to 100 gallons of water per acre. Special concentrate equipment is necessary for these areas. **AIR APPLICATION:** Apply specified rate in 5 to 20 gallons of water per acre. Orchard rates should not be applied in less than 10 gallons of water per acre. **FOR BEST RESULTS** use sufficient water per acre to give thorough coverage of all plant parts. **ALMONDS, CHERRIES, PLUMS, PRUNES:** Brown Rot, Leaf Spot, Powdery Mildew, Silver Mite, Red Spider Mite - Apply 10 to 30 lbs. per acre. Apply at bloom or early petal fall and repeat at 10 - 14 day intervals as necessary or after period of wet weather. APPLE, PEARS (Except D'Anjou):
Powdery Mildew, Scab, Red Spider Mite - Apply 10 to 30 lbs. per acre at pre bloom and petal fall and Brown Spot Blossom Blight, Powdery Mildew, Silver Mite, Red Spider Mite - Apply 10 to 30 lbs. per acre at bloom be a light and repeat at 10 - 14 day intervals as necessary or after period of wet weather. CITRUS (GRAPEFRUITS, KUMQUATS, LEMONS, LIMES, ORANGES, TANGERINES, TANGELOS): Flat Mite, Red Spider Mite - Apply 8 to 20 lbs. per acre. Apply in any spray NOT CONTAINING OIL or European Red Mite, Almond Mite, Two-Spotted Mite, Red Spider Mite - Apply 10 to 20 lbs. per acres. Apply with mite sprays to increase effectiveness. CAUTION: Do no apply unless variety is known to be tolerant to sulphur. **Powdery Mildew, Red Spider Mite** - Apply 5 lbs. per acre. Apply in late dormancy, with cutworm spray, early shoot period and every 14 days through season. **BLACKBERRY** (Rubus spp. **Powdery Mildew** - Apply 6 to 15 lbs. per acre. Apply when new shoots are 6" long and before blossoms open. RASPBERRY (Rubus spp.): Powdery Mildew (Sphaerotheca humuli) - Apply 6 to 15 lbs. per acre. Apply each week starting at first bloom color, until fruit is set, then repeat as weather requires. **Powdery Mildew, Red Spider Mite** - Apply 5 to 10 lbs. per acre. Apply at early leaf stage. For powdery mildew, at first signs of disease and repeat as needed. BEANS, PEAS, BROCCOLI, BRUSSELS SPROUT, CABBAGE, CAULIFLOWER, COLLARD, KALE, Powdery Mildew, Red Spider Mite - Apply 5 lbs. per acre. Apply at early leaf stage and repeat every 10 to 14 days. TOMATOES:

Powdery Mildew, Leaf Spot, Red Spider Mite - Apply 8 lbs. per acre. Start at early leaf stage and repeat as

AND LIMITATIONS OF LIABILITY AND REMEDIES

Read the Conditions of Sale - Warranty and Limitations of Liability and Remedies before using this product.

If the terms are ot acceptable, return the product, unopened, and the full purchase price will be refunded.

control of pests and/or injury to the crop to which the product is applied may result from the occurrence of

extraordinary or unusual weather conditions or the failure to follow the label directions or good application practices, all of which are beyond the control of Helena Chemical Company (the "Company") or seller. In

addition, failure to follow label directions may cause injury to crops, animals, man or the environment. The

concerning the product, including no implied warranty of merchantability or fitness for any particular

The exclusive remedy against the Company for any cause of action relating to the handling or use of this product shall be limited to, at Helena Chemical Company's election, one of the following:

Refund of the purchase price paid by buyer or user for the product bought, or

Company warrants that this product conforms to the chemical description on the label and is reasonable fit for the purpose referred to in the directions for use subject to the factors noted above which are beyond the

control of the Company. The Company makes no other warranties or representations of any kind, express or implied,

The directions on this label are believed to be reliable and should be followed carefully. Insufficient

Some fruits and other plants are susceptible to injury from sulphur under certain climatic conditions. The user is advised not to use sulphur on any crop unless local use has proved that sulphur does not damage crops in

that locality. Sulphur may burn foliage when the temperature is high (90° or above). Do not make sulphur applications on Eastern American type grapes, spinach, apricots, or other sulphur susceptible plants. Do not

For light to moderate infestations use lower amount recommended. For moderate to heavy infestations use

make applications within two weeks before or after an oil spray.

higher of recommended amounts. Use enough water to insure adequate coverage.

3.5" Match Line

4.5" Short Seam

Glue

Seam

SUGAR BEETS: Red Spider Mite - Apply 5 lbs. per acre as needed. Black Spot, Powdery Mildew - Apply 6 to 8 lbs. per 100 gallons of water. Apply at first sign of disease and CONDITIONS OF SALE – LIMITED WARRANTY

purpose, and no such warranty shall be implied by law.

Red Spider Mite - Apply 5 lbs. per acre as needed.

COTTON, SEED ALFALFA:

necessary, usually 10-24 days.

Russet Mite - Apply 5 lbs. per care as needed. Thorough coverage required.

© Copyright Helena Chemical Company, 2010

To the extent allowed by law, the Company shall not be liable and any and all claims against the Company are

waived for special, indirect, incidental, or consequential damages or expense of any nature, including, but not

limited to, loss of profits or income. The Company and the seller offer this product and the buyer and user accept it, subject to the foregoing conditions of sale and limitation of warranty, liability and remedies.

3.25" Flap

NET CONTENTS: 30 LBS. (13.61 kg)





