



FOR CONTROL OF CERTAIN BROADLEAF WEEDS IN SMALL GRAINS (WHEAT, BARLEY, OATS AND RYE), CONSERVATION RESERVE PROGRAM (CRP) AREAS, GRASSES GROWN FOR SEED PRODUCTION AND FLAX

ACTIVE INGREDIENT:

Octanoic acid of ester of bromoxynil (3,5-dibromo-4-Hydroxybenzotrile).....	31.7%*
Isooctyl ester of 2-methyl-chlorophenoxyacetic acid**.....	34.0%**
INERT INGREDIENTS	34.3%

TOTAL..... 100.0%

Contains Petroleum Distillate. Patent No. 6,232,272

*Bromoxynil octanoate equivalent to 21.8% of bromoxynil or not less than 2.0 pounds of bromoxynil per gallon.

**Equivalent to 21.8% 2-methyl-chlorophenoxyacetic acid or not less than 2.0 pounds MCPA acid per gallon.

KEEP OUT OF REACH OF CHILDREN CAUTION/PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Avoid breathing spray mist.

**For Chemical Spill, Leak, Fire, or Exposure,
Call CHEMTREC 1-800-424-9300
For Medical Emergencies Only, Call 1-877-325-1840**

FIRST AID

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment information.

Note to Physician: May pose an aspiration pneumonia hazard. Contains petroleum distillate.

NET CONTENTS:

STORAGE AND DISPOSAL

STORAGE: Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Store at temperatures above 32°F. If allowed to freeze, remix before using.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: NONREFILLABLE CONTAINER (EQUAL TO OR LESS THAN 5 GALLONS): Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds.

Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke. **NONREFILLABLE CONTAINER (GREATER THAN 5 GALLONS):** Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke. **REFILLABLE CONTAINERS (LARGER THAN 5 GALLONS):** Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. If unable to refill, offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. Prior to refilling, inspect thoroughly for damage such as cracks, punctures, abrasions and damaged or worn-out threads on closure devices. Do not refill or transport damaged or leaking containers. Check for leaks after refilling and before transportation. Do not refill or transport damaged or leaking containers. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

SEE INSIDE LEAFLET FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND COMPLETE DIRECTIONS FOR USE.

EPA Reg. No. 5905-550

AD 022712

EPA Est. No. 228-IL-1

Product of United Kingdom

Manufactured For
HELENA CHEMICAL COMPANY
225 SCHILLING BOULEVARD, SUITE 300
COLLIERVILLE, TENNESSEE 38017

PEEL BACK BOOK HERE AND RESEAL AFTER OPENING

13187000

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Avoid breathing spray mist.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are Barrier laminate, Butyl rubber, Nitrile rubber, or Viton gloves. If you want more options, follow the instructions for category *F* on an EPA chemical resistance category selection chart.

Mixers, Loaders, Applicators, Flaggers, and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves for cleaning equipment, mixing, and loading
- Chemical-resistant apron for cleaning equipment, mixing, and loading

Additional PPE Requirements for mixers and loaders supporting aerial application to rangelands, pasture lands, or non-cropland. The mixers/loaders also must wear:

- Chemical-resistant apron, and
- NIOSH-approved particulate filtering respirator equipped with N, R, or P class filter media. The respirator should have a NIOSH approval number prefix TC-84A. It is recommended that you require the respirator wearer be fit tested, and trained in the use, maintenance, and limitations of the respirator.

USER SAFETY REQUIREMENTS

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS FOR AERIAL APPLICATIONS

Enclosed Cockpits Engineering Controls: Pilots must use an enclosed cockpit that meets the requirements listed in the WPS for agricultural pesticides [40 CFR 170.240(d)(6)].

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to wildlife and fish. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollution Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

APPLICATION BY CHEMIGATION must be done by fixed pipe, overhead sprinkler systems or hand-moved pipe. If hand moved pipe is used for chemigation, the pipe must not be handled in any way until 24 hours after chemigation has been completed and residues have been flushed from the system. When applying by chemigation, no person may enter the application site unless in an enclosed vehicle.

DURING AERIAL APPLICATIONS, human flaggers are prohibited unless in enclosed vehicles. Aerial application is prohibited within 300 feet of residential areas (e.g., homes, schools, hospitals, shopping areas, etc.).

PHYSICAL AND CHEMICAL HAZARDS

Do not use or store near heat or open flame. Do not cut

WILDCARD® XTRA Herbicide contains low volatile isocyanate. This product may cause injury to susceptible plants. The

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in or on any area where it is prohibited by Federal, State, or local

Do not apply this product in a way that will contact workers or other persons, either directly or by drift, during application, or until after the application is complete. For any requirements or restrictions, see the pesticide label. For any required state or local regulations, see the pesticide label.

AGRICULTURAL USE RESTRICTIONS

Use this product only in accordance with its labeling and the requirements for the protection of agricultural crops from agricultural pesticides. It contains requirements for treatment of specific crops and exceptions pertaining to restricted-entry interval. The requirements in this section are in addition to the requirements in the Protection Standard.

Do not enter or allow worker entry into treated areas until the restricted-entry interval (REI) has expired.

The restricted-entry interval (REI) for cotton is 4 days. For other crops, see the Protection Standard for certified crop advisors does not apply to the treated area during the entire 4-day REI for broadcast applications of this requirement.

PPE required for early entry to treated areas that is in addition to the PPE required for entry to treated areas that has been treated, such as plants, animals, or structures, includes: nitrile, viton or barrier gloves; eye protection; and protective eyewear.

NON-AGRICULTURAL USE RESTRICTIONS

The requirements in this box apply to uses of this product that are not agricultural pesticides (40 CFR Part 170). The WPS applies to uses in forests, nurseries, or greenhouses.

Do not enter or allow others to enter until sprays have dried.

STORAGE AND DISPOSAL

STORAGE: Do not contaminate water, food, or feed with storage or disposal. Store above 32°F. If allowed to freeze, remix before using.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on a pesticide disposal facility.

CONTAINER DISPOSAL: **NONREFILLABLE CONTAINER (EQUAL TO OR LARGER THAN THE ORIGINAL CONTAINER):** Triple rinse container (or equivalent) promptly into application equipment or a mix tank and drain for 10 seconds. Shake for 10 seconds. Pour rinsate into a container and recap. Shake for 10 seconds. Pour rinsate into a container and recap. Shake for 10 seconds. Pour rinsate into a container and recap. Shake for 10 seconds after the flow begins to drip. Refill and dispose of in a sanitary landfill, or incinerator, if allowed. Do not burn.

PHYSICAL AND CHEMICAL HAZARDS

Do not use or store near heat or open flame. Do not cut or weld container.

WARNING

WILDCARD® XTRA Herbicide contains low volatile isooctyl ester of MCPA. At high air or ground surface temperatures, vapors from this product may cause injury to susceptible plants. This fact should be considered when applying **WILDCARD® XTRA**.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read entire label before using this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

The restricted-entry interval (REI) for cotton is 4 days and includes scouts and crop advisors. The exemption in the Worker Protection Standard for certified crop advisors does not apply to bromoxynil. Scouts and crop advisors are prohibited from entering the treated area during the entire 4-day REI for bromoxynil. Applicators and other users must inform crop advisors and scouts of this requirement.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is coveralls over a long-sleeved shirt and long pants, chemical-resistant gloves such as nitrile, viton or barrier laminate, chemical-resistant footwear plus socks, chemical-resistant headgear for overhead exposure and protective eyewear.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter or allow others to enter until sprays have dried.

STORAGE AND DISPOSAL

STORAGE: Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Store at temperatures above 32°F. If allowed to freeze, remix before using.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL:

NONREFILLABLE CONTAINER (EQUAL TO OR LESS THAN 5 GALLONS): Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

(continued)

STORAGE

CONTAINER DISPOSAL: (cont.)

NONREFILLABLE CONTAINER (GREATER THAN 5 GALLONS): Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

REFILLABLE CONTAINERS (LARGER THAN 5 GALLONS): Do not reuse this container for any other purpose. If used for a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke. If used for a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke. If used for a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke. If used for a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

GENERAL INFORMATION

WILDCARD® XTRA is formulated as an emulsifiable concentrate (EC) containing 1.5 pounds per gallon of MCPA.

WILDCARD® XTRA is a selective postemergence herbicide for use on corn, soybeans, and sorghum. It is applied to actively growing weed seedlings. The use of the weed seedlings is essential for optimum control.

WILDCARD® XTRA has little residual activity. Therefore, crops that form a good canopy will help suppress weeds. For example Yaccora Rojo wheat, may not develop tillers. Occasional transitory leaf burn may occur. The temporary leaf burn on **WILDCARD® XTRA** is mainly contact, recovery of the crop may be greater when crops are stressed by abrasion caused by hail, sleet or insect feeding. To reduce the risk of leaf burn, the recommended spray volumes per acre when weeding are:

MIXING, LOADING

2.5 Gallon Containers

It is strongly recommended that special care be taken when mixing and loading to avoid possible drip or splash.

30 Gallon and Bulk Containers

If you will handle a total of 60 gallons or more of this herbicide during operations. If this product is packaged in a 30 gallon container, a drip-free hard coupling which may be used only with a hose should be used to avoid possible drip or splash. Do not presently own or have access to a mechanical system on how to obtain such a system or to modify your present system. Connect the pump or probe from the container until it is used to rinse the empty container and to transfer the rinsate to the application equipment.

WILDCARD® XTRA ALONE: Fill the spray tank 1/2 full with water. Add water to the spray tank to the recommended volume for application.

STORAGE AND DISPOSAL (cont.)

CONTAINER DISPOSAL: (cont.)

NONREFILLABLE CONTAINER (GREATER THAN 5 GALLONS): Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

REFILLABLE CONTAINERS (LARGER THAN 5 GALLONS): Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. If unable to refill, offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. Prior to refilling, inspect thoroughly for damage such as cracks, punctures, abrasions and damaged or worn-out threads on closure devices. Do not refill or transport damaged or leaking containers. Check for leaks after refilling and before transportation. Do not refill or transport damaged or leaking containers. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

GENERAL INFORMATION

WILDCARD® XTRA is formulated as an emulsifiable concentrate containing the equivalent of 2 lbs. per gallon of bromoxynil and 2 pounds per gallon of MCPA.

WILDCARD® XTRA is a selective postemergence herbicide for control of important broadleaf weeds infesting small grains (wheat, barley, oats, rye), conservation reserve program areas, and grass grown for seed. Optimum weed control is obtained when **WILDCARD® XTRA** is applied to actively growing weed seedlings. **WILDCARD® XTRA** is primarily a contact herbicide; therefore thorough coverage of the weed seedlings is essential for optimum control.

WILDCARD® XTRA has little residual activity. Therefore subsequent flushes of weeds will not be controlled by the initial treatment.

Generally, crops that form a good canopy will help shade subsequent weed flushes. However, certain crops or short-straw varieties, for example Yaccora Rojo wheat, may not develop the crop canopy fast enough to shade the subsequent flushes of weeds.

Occasional transitory leaf burn may occur. The temporary leaf burn is similar to that seen with liquid fertilizer. Because the activity of **WILDCARD® XTRA** is mainly contact, recovery of the crop is generally rapid with no lasting effect. Frequency and amount of leaf burn may be greater when crops are stressed by abrasive winds, cool to cold evening temperatures or mechanical injury, such as that caused by hail, sleet or insect feeding. To reduce the potential for temporary leaf burn, applications should be made to dry foliage in the recommended spray volumes per acre when weather conditions are not extreme.

MIXING, LOADING AND HANDLING INSTRUCTIONS

2.5 Gallon Containers

It is strongly recommended that special care be taken in mixing and loading this product. Hands should be placed on the container in such a way as to avoid possible drip or splash.

30 Gallon and Bulk Containers

If you will handle a total of 60 gallons or more of this product per day, you must use a mechanical transfer system for all mixing and loading operations. If this product is packaged in a 30 gallon drum, you must use a mechanical transfer system which terminates in a drip-free hard coupling which may be used only with a spray or mix tank which has been fitted with a compatible coupling. If you do not presently own or have access to a mechanical transfer system with this type of coupling, contact your dealer for information on how to obtain such a system or to modify your present system. When using a mechanical transfer system, do not remove or disconnect the pump or probe from the container until the container has been emptied and rinsed. The pump or probe system must be used to rinse the empty container and to transfer the rinsate directly to the mixing or spray tank.

WILDCARD® XTRA ALONE: Fill the spray tank 1/2 to 3/4 full with clean water. Begin agitation and add the recommended amount of **WILDCARD® XTRA**. Add water to the spray tank to the desired level. Maintain sufficient agitation to ensure a uniform spray mixture during application.

TANK MIXTURES: WILDCARD® XTRA can be applied on approved crops. Refer to the specific crop section in mixture with another product, fill the spray tank powder, soluble powder, flowable or dry flowable product thoroughly mixed with water add the recommended amount. If tank mixing with other product types, add the **W** water thoroughly before adding another product or mixing in water.

Maintain sufficient agitation while mixing and duration remain without agitation for short periods of time,

If tank mixing with products other than those listed on the label, test for compatibility, use the same ratio as the anticipated use. Indications of incompatibility usually will be seen during spraying. Indications of incompatibility usually will be seen during spraying. Indications of incompatibility usually will be seen during spraying. Indications of incompatibility usually will be seen during spraying.

SPRAYABLE LIQUID

WILDCARD® XTRA can be applied in combination with liquid fertilizer. When tank mixing with liquid fertilizer always use **WILDCARD® XTRA**. Always predetermine the correct ratio. Agitation must be maintained during filling and application. Leaf burn may occur when **WILDCARD® XTRA** is applied.

CAUTION: Fertilizers and spray additives can interact with **WILDCARD® XTRA** if they are applied to **WILDCARD® XTRA**.

APPL

WILDCARD® XTRA can be applied to registered uses.

GROUND APPLICATION

Use a standard herbicide boom sprayer that provides a nozzle finer than 50 mesh in the nozzle tips and in-line strainer. Select a spray volume and delivery system that will provide thorough coverage, use of flat fan nozzles (maximum spray pressure and lower spray pressures that produce optimum control. **Raindrop®** nozzles and flood nozzles are not recommended.

In general, a spray volume of 10 to 20 gallons per acre with a minimum spray pressure of 50 psi and a ground application if ground terrain, crop and weed conditions are favorable. A maximum ground speed of 10 mph is suggested. Ground application in wheel track areas. Applications using less than 10 gallons per acre may result in poor spray coverage.

When weed infestations are heavy, use of higher spray volumes may be necessary. When corn or grain sorghum are large enough to prevent weed coverage. If you are unsure of the infestation level, do not apply when winds are gusty or when other conditions are unfavorable.

Additional requirements for ground boom application: Do not apply with a nozzle height greater than 4 feet.

AERIAL APPLICATION

Use orifice discs, cores and nozzle types and arrangements. In general, a minimum spray volume of 5 GPA and 3 gallons per acre may be used if crop canopy and ground conditions are favorable. 5 gallons of spray volume per acre may result in poor spray coverage.

temperatures, vapors from **WILDCARD® XTRA**.

Read the label before using this product.

Only protected handlers are responsible for pesticide application.

Part 170. This standard covers the use of protective equipment (PPE), including coveralls, gloves, and eye protection, by workers in agricultural settings.

Exemption in the Worker Protection Standard for agricultural settings.

and long pants, chemical-resistant headgear, and gloves.

Protection Standard for agricultural plants on farms, and other agricultural settings.

d. Store at temperatures between 40°F and 80°F.

Do not reuse or refill this container. The remaining contents may be used for later use or disposal. If burned, stay out of smoke.

(continued)



TANK MIXTURES: **WILDCARD® XTRA** can be applied in tank mixture with many other herbicides and insecticides registered for use on approved crops. Refer to the specific crop section for rate recommendations and other restrictions. To apply **WILDCARD® XTRA** in mixture with another product, fill the spray tank 1/2 to 3/4 full with clean water and begin agitation. If tank mixing with wettable powder, soluble powder, flowable or dry flowable products, add the powder or flowable product first. After the other herbicide is thoroughly mixed with water add the recommended amount of **WILDCARD® XTRA** and add water to the spray tank to the desired level. If tank mixing with other product types, add the **WILDCARD® XTRA** first before adding the other product. Always mix one product in water thoroughly before adding another product or compatibility problems may occur. Never mix two products together without first mixing in water.

Maintain sufficient agitation while mixing and during application to ensure a uniform spray mixture. If spray mixture is allowed to remain without agitation for short periods of time, be sure to agitate until uniformly mixed before application.

If tank mixing with products other than those listed within each crop section, a compatibility test is recommended to ensure satisfactory spray preparation. To test for compatibility, use a small container and mix a small amount (0.5 to 1 quart) of spray, combining all ingredients in the same ratio as the anticipated use. If any indications of physical incompatibility develop, do not use this mixture for spraying. Indications of incompatibility usually will appear within 5 to 15 minutes after mixing. To ensure maximum crop safety and weed control, follow all cautions and limitations on this label and the labels of products used in the tank mixture with **WILDCARD® XTRA**.

SPRAYABLE LIQUID FERTILIZERS AND SPRAY ADDITIVES

WILDCARD® XTRA can be applied in combination with sprayable liquid fertilizer or spray additives such as surfactants or crop oil concentrate. When tank mixing with liquid fertilizer always add the fertilizer to the spray tank first and agitate thoroughly before adding **WILDCARD® XTRA**. Always predetermine the compatibility with liquid fertilizer by mixing small proportional quantities in advance. Agitation must be maintained during filling and application operations to ensure that **WILDCARD® XTRA** is evenly mixed with the fertilizer. Leaf burn may occur when **WILDCARD® XTRA** is applied with liquid fertilizer, but new leaves are not adversely affected.

CAUTION: Fertilizers and spray additives can increase foliage leaf burn when applied with **WILDCARD® XTRA**. Do not apply fertilizers or spray additives with **WILDCARD® XTRA** if leaf burn is a major concern due to environmental conditions, crop or variety sensitivity to **WILDCARD® XTRA**.

APPLICATION PROCEDURES

WILDCARD® XTRA can be applied to registered use areas by ground, aerial and sprinkler irrigation equipment.

GROUND APPLICATION

Use a standard herbicide boom sprayer that provides uniform and accurate application. Sprayer should be equipped with screens no finer than 50 mesh in the nozzle tips and in-line strainers.

Select a spray volume and delivery system that will ensure thorough and uniform spray coverage. For optimum spray distribution and thorough coverage, use of flat fan nozzles (maximum tip size 8008) with a spray pressure of 40–60 psi are recommended. Other nozzle types and lower spray pressures that produce coarse spray droplets may not provide adequate coverage of the weeds to ensure optimum control. **Raindrop®** nozzles and flood nozzles are not recommended as weed control with **WILDCARD® XTRA** may be reduced.

In general, a spray volume of 10 to 20 gallons per acre (GPA) is recommended for optimum spray coverage. A minimum of 5 GPA with a minimum spray pressure of 50 psi and a maximum ground speed of 10 mph may be used with higher speed, low volume ground application if ground terrain, crop and weed density allow effective spray distribution. When using higher speed equipment, a maximum ground speed of 10 mph is suggested if field conditions cause excessive boom movement during application which results in poor spray coverage. Ground applications made when dry, dusty field conditions exist may provide reduced weed control in wheel track areas. Applications using less than 10 gallons per acre may result in reduced weed control.

When weed infestations are heavy, use of higher spray volumes and spray pressure will be helpful in obtaining uniform weed coverage. When corn or grain sorghum are large enough to interfere with the spray pattern, drop nozzles should be used to obtain uniform weed coverage. If you are unsure of the infestation level or size of crop, consult your local extension service.

Do not apply when winds are gusty or when other conditions favor poor spray coverage and/or off-target spray movement.

Additional requirements for ground boom application:

Do not apply with a nozzle height greater than 4 feet above the crop canopy.

AERIAL APPLICATION

Use orifice discs, cores and nozzle types and arrangements that will provide for optimum spray distribution and maximum coverage. In general, a minimum spray volume of 5 GPA and a maximum pressure of 40 psi are recommended. A minimum spray volume of 3 gallons per acre may be used if crop canopy and weed density allow adequate spray coverage. Aerial applications using less than 5 gallons of spray volume per acre may result in reduced weed control.

Do not apply during inversion conditions, when you get spray movement. Off-target spray movement winds exceed 10 mph.

SPRAY DRIFT MANAGEMENT

Avoiding spray drift at the application site is the related factors determine the potential for spray when making decisions.

Apply only as a medium or coarser spray (ASAE atomizer nozzles).

Apply only when the wind speed is 2–10 mph at

ADDITIONAL REQUIREMENTS FOR AERIAL

The boom length must not exceed 75% of the width of the release spray.

Release spray at the lowest height consistent with the above the crop canopy.

When applications are made with a crosswind, it is recommended to adjust the path of the aircraft upwind.

Do not make applications into temperature inversions.

SPRINKLER IRRIGATION APPLICATION:

WILDCARD® XTRA Herbicide can be applied through

Apply **WILDCARD® XTRA** Herbicide through sprinkler irrigation systems only. If hand-moved pipette after chemigation has been completed and residue may enter the application site unless in an enclosure.

SPECIFIC REQUIREMENTS FOR APPLICATION:

- The system must contain a functional check gate pipeline to prevent water source contamination.
- The pesticide injection pipeline must contain a check gate toward the injection pump.
- The pesticide injection pipeline must also contain a check gate to the side of the injection pump and connected to the irrigation system is either automatic or manual.
- The system must contain functional interlock pump motor stops.
- The irrigation line or water pump must include a pressure relief valve to the point where pesticide is injected.
- Systems must use a metering pump, such as a diaphragm pump, and constructed of materials that are compatible with the pesticide.
- Do not apply when wind speed favors drift.
- Agitation is recommended in the pesticide side of the injection pump.
- WILDCARD® XTRA** Herbicide should be applied through a lateral move system. Application of irrigation set with other overhead sprinkler systems.
- For best performance, set the sprinkler system to apply 1.5 inches of water per inch of wheel track.
- Remove scale, pesticide residues and other foreign matter from the system.
- If **WILDCARD® XTRA** Herbicide is diluted in water, it should be applied through a lateral move system. Start the sprinklers and then inject 1 inch of water. Refer to the **WILDCARD® XTRA** Herbicide label for more information.

CHEMIGATION

Application of more than 0.5 inch/acre of irrigator spray is not recommended. Do not apply when conditions favor drift, when wind speed is greater than 10 mph.



insecticides registered for use. To apply **WILDCARD® XTRA**, if tank mixing with wettable powder the other herbicide is thor-spray tank to the desired level. Always mix one product in products together without first

If spray mixture is allowed to settle.

Recommended to ensure satisfactory (1 quart) of spray, combining all products, do not use this mixture for ure maximum crop safety and mixture with **WILDCARD® XTRA**.

DIRECTIONS

As surfactants or crop oil concentrate thoroughly before adding additional quantities in advance. Always mix evenly with the fertilizer not adversely affected.

WILDCARD® XTRA. Do not apply fertilizers, crop or variety sensi-

equipment.

It should be equipped with screens no

for optimum spray distribution and are recommended. Other nozzles coverage of the weeds to ensure that **WILDCARD® XTRA** may be

coverage. A minimum of 5 GPA with higher speed, low volume using higher speed equipment, prevent during application which provide reduced weed control effect.

Obtaining uniform weed coverage should be used to obtain uniform service.

Target spray movement.

Application and maximum coverage. Always use a minimum spray volume of 5 GPA for all applications using less than

Do not apply during inversion conditions, when winds are gusty or when other conditions favor poor spray coverage and/or off-target spray movement. Off-target spray movement can be minimized by increasing the spray volume per acre and not applying when winds exceed 10 mph.

SPRAY DRIFT MANAGEMENT

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment- and weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

Apply only as a medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

Apply only when the wind speed is 2–10 mph at the application site.

ADDITIONAL REQUIREMENTS FOR AERIAL APPLICATION:

The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter.

Release spray at the lowest height consistent with efficacy and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy.

When applications are made with a crosswind, the swath will be displaced downwind. The applicator must compensate for this by adjusting the path of the aircraft upwind.

Do not make applications into temperature inversions.

SPRINKLER IRRIGATION APPLICATION:

WILDCARD® XTRA Herbicide can be applied through sprinkler irrigation systems to small grains and grasses grown for seed.

Apply **WILDCARD® XTRA** Herbicide through sprinkler systems including center pivot, lateral move, side (wheel) roll, solid set or hand move irrigation systems only. If hand-moved pipe is used for chemigation, the pipe must not be handled in any way until 24 hours after chemigation has been completed and residues have been flushed from the system. When applying by chemigation, no person may enter the application site unless in an enclosed vehicle. Do not apply this product through any other type of irrigation system.

SPECIFIC REQUIREMENTS FOR APPLICATION THROUGH AUTOMATED SPRINKLER IRRIGATION SYSTEM:

- The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- Do not apply when wind speed favors drift beyond the area intended for treatment.
- Agitation is recommended in the pesticide supply tank when applying the **WILDCARD® XTRA** Herbicide.
- WILDCARD® XTRA** Herbicide should be applied continuously for the duration of the water application with center pivot and continuous lateral move systems. Application of **WILDCARD® XTRA** Herbicide should be made during the last 30–45 minutes of the irrigation set with other overhead sprinkler systems.
- For best performance, set the sprinkler system to deliver approximately 0.5 inch or less of water per acre.
- Remove scale, pesticide residues and other foreign matter from the supply tank and entire injector system. Flush with clean water.
- If **WILDCARD® XTRA** Herbicide is diluted in the supply tank, fill the tank with half of the water amount desired, add the **WILDCARD® XTRA** and then add remaining water amount with agitation. Always dilute with at least 4 parts water to 1 part **WILDCARD® XTRA**. Start the sprinklers and then inject **WILDCARD® XTRA** Herbicide into the irrigation line. **WILDCARD® XTRA** should be injected with a positive displacement pump into the main line at least 8 feet ahead of a right angle turn to ensure adequate mixing. Refer to the **WILDCARD® XTRA** Herbicide label for detailed information on application rates and timings.

CHEMIGATION USER PRECAUTIONS

Application of more than 0.5 inch/acre of irrigation water may result in decreased product performance on certain soils.

Do not apply when conditions favor drift, when system connections or fittings leak, or when nozzles do not provide uniform distribution.

Allow sufficient time for pesticide to be flushed. Crop injury, lack of effectiveness, or illegal application. Do not connect an irrigation system used for pest control. If you have questions about calibration, you should consult a person knowledgeable of the chemigation system. A person knowledgeable of the chemigation system, shall shut the system down and make

Postemergence application of **WILDCARD® XTRA**. Maximum weed stage of growth is listed under

MOST SUSCEPTIBLE BROADLEAF WEED

Annual sowthistle	(<i>Sonchus oleraceus</i>)
Black mustard	(<i>Brassica nigra</i>)
Black nightshade	(<i>Solanum nigrum</i>)
Common cocklebur	(<i>Xanthium strumarium</i>)
Common lambsquarters	(<i>Chenopodium album</i>)
Common tarweed	(<i>Hemizonia congesta</i>)
Cow cockle	(<i>Saponaria vaccaria</i>)
Cutleaf nightshade	(<i>Solanum triflorum</i>)
Eastern black nightshade	(<i>Solanum elaeagnifolium</i>)
Coast fiddleneck	(<i>Amsinckia intermedia</i>)
Field pennycress	(<i>Thlaspi arvense</i>)
Green smartweed	(<i>Polygonum scaberrimum</i>)
Hairy nightshade	(<i>Solanum sarachoides</i>)
Horned Poppy	(<i>Glaucium corniculatum</i>)
Jimsonweed	(<i>Datura stramonium</i>)
Ladysthumb	(<i>Polygonum persicaria</i>)
Lanceleaf sage	(<i>Salvia reflexa</i>)
London rocket	(<i>Sisymbrium irio</i>)
Marshelder	(<i>Iva xanthifolia</i>)
Pennsylvania smartweed	(<i>Polygonum strumarium</i>)
Pepperweed spp.	(<i>Lepidium</i> spp.)
Redroot pigweed	(<i>Amaranthus retrofractus</i>)
Russian thistle	(<i>Salsola kali</i>)
Shepherdspurse	(<i>Capsella bursa-pastoris</i>)
Silverleaf nightshade	(<i>Solanum elaeagnifolium</i>)
Smooth pigweed	(<i>Amaranthus hybridus</i>)
Spiny pigweed	(<i>Amaranthus spinosus</i>)
'Sunflower	(<i>Helianthus annuus</i>)
Tall Waterhemp	(<i>Amaranthus tuberosus</i>)
Tartary buckwheat	(<i>Fagopyrum tataricum</i>)
Tumble mustard	(<i>Sisymbrium altissimum</i>)
Wild buckwheat	(<i>Polygonum convolvulus</i>)
Wild mustard	(<i>Sinapis arvensis</i>)
Yellow rocket	(<i>Barbarea vulgaris</i>)

¹For control of sunflower, delay application until

Canada Thistle (*Cirsium arvense*)
WILDCARD® XTRA Herbicide applied at 1-1/2 gallons per acre. Do not apply when Canada thistle is 8 inches tall to the

ir spray coverage and/or off-tar-
per acre and not applying when

many equipment- and weather-
for considering all these factors

microns or greater for spinning

at a height greater than 10 feet

or must compensate for this by

grasses grown for seed.

de (wheel) roll, solid set or hand
ndled in any way until 24 hours
lying by chemigation, no person
other type of irrigation system.

IRRIGATION SYSTEM:
appropriately located on the irri-

to prevent the flow of fluid back

ated valve located on the intake
withdrawn from the supply tank

injection pump when the water

ter pump motor when the water

agm pump) effectively designed
h a system interlock.

rbicide.
ation with center pivot and con-
ig the last 30–45 minutes of the

r per acre.
r system. Flush with clean water.
amount desired, add the **WILD-**
arts water to 1 part **WILDCARD®**
: **WILDCARD® XTRA** should be
le turn to ensure adequate mix-
and timings.

nce on certain soils.
not provide uniform distribution.

Allow sufficient time for pesticide to be flushed through all the lines and nozzles before turning off irrigation water.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

Do not connect an irrigation system used for pesticide application to a public water system.

If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

A person knowledgeable of the chemigation system and responsible for its operations, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

GENERAL WEED LIST

Postemergence application of **WILDCARD® XTRA** Herbicide will control the following weeds when sprayed in the seedling stage. Maximum weed stage of growth is listed under **WILDCARD® XTRA** RECOMMENDATIONS.

MOST SUSCEPTIBLE BROADLEAF WEED SPECIES

Annual sowthistle	(<i>Sonchus oleraceus</i>)
Black mustard	(<i>Brassica nigra</i>)
Black nightshade	(<i>Solanum nigrum</i>)
Common cocklebur	(<i>Xanthium strumarium</i>)
Common lambsquarters	(<i>Chenopodium album</i>)
Common tarweed	(<i>Hemizonia congesta</i>)
Cow cockle	(<i>Saponaria vaccaria</i>)
Cutleaf nightshade	(<i>Solanum triflorum</i>)
Eastern black nightshade	(<i>Solanum ptycanthum</i>)
Coast fiddleneck	(<i>Amsinckia intermedia</i>)
Field pennycress	(<i>Thlaspi arvense</i>)
Green smartweed	(<i>Polygonum scabrum</i>)
Hairy nightshade	(<i>Solanum sarachoides</i>)
Horned Poppy	(<i>Glaucium corniculatum</i>)
Jimsonweed	(<i>Datura stramonium</i>)
Ladysthumb	(<i>Polygonum persicaria</i>)
Lanceleaf sage	(<i>Salvia reflexa</i>)
London rocket	(<i>Sisymbrium irio</i>)
Marshelder	(<i>Iva xanthifolia</i>)
Pennsylvania smartweed	(<i>Polygonum strumarium</i>)
Pepperweed spp.	(<i>Lepidium</i> spp.)
Redroot pigweed	(<i>Amaranthus retroflexus</i>)
Russian thistle	(<i>Salsola kali</i>)
Shepherdspurse	(<i>Capsella bursa-pastoris</i>)
Silverleaf nightshade	(<i>Solanum elaeagnifolium</i>)
Smooth pigweed	(<i>Amaranthus hybridus</i>)
Spiny pigweed	(<i>Amaranthus spinosus</i>)
1Sunflower	(<i>Helianthus annuus</i>)
Tall Waterhemp	(<i>Amaranthus tuberculatus</i>)
Tartary buckwheat	(<i>Fagopyrum tataricum</i>)
Tumble mustard	(<i>Sisymbrium altissimum</i>)
Wild buckwheat	(<i>Polygonum convolvulus</i>)
Wild mustard	(<i>Sinapis arvensis</i>)
Yellow rocket	(<i>Barbarea vulgaris</i>)

¹For control of sunflower, delay application until first sunflower seedlings emerging are 4 inches in height.

SUSCEPTIBLE BROADLEAF WEED SPECIES

Blue (purple) mustard	(<i>Chlorispora tenella</i>)
Common groundsel	(<i>Senecio vulgaris</i>)
Common ragweed	(<i>Ambrosia artemisiifolia</i>)
Corn chamomile	(<i>Anthemis arvensis</i>)
Corn gromwell	(<i>Lithospermum arvense</i>)
Fumitory	(<i>Fumaria officinalis</i>)
Giant ragweed	(<i>Ambrosia trifida</i>)
Hemp sesbania	(<i>Sesbania exaltata</i>)
Henbit	(<i>Lamium amplexicaule</i>)
lyleaf morningglory	(<i>Ipomoea hederacea</i>)
Knawel	(<i>Scleranthus annuus</i>)
Kochia	(<i>Kochia scoparia</i>)
Mayweed	(<i>Anthemis cotula</i>)
Prostrate knotweed	(<i>Polygonum aviculare</i>)
Puncture vine	(<i>Tribulus terrestris</i>)
Tall morningglory	(<i>Ipomoea purpurea</i>)
Tansy mustard	(<i>Descurainia pinnata</i>)
Tarweed	(<i>Hemizonia</i> spp.)
Velvetleaf	(<i>Abutilon theophrasti</i>)
Wild radish	(<i>Raphanus raphanistrum</i>)

Weeds germinating after spraying will not be controlled.

WEED SUPPRESSION

Canada Thistle (*Cirsium arvense*)

WILDCARD® XTRA Herbicide applied at 1-1/2 pints per acre provides burn down of top growth. Regrowth may occur. Make applications when Canada thistle is 8 inches tall to the bud stage.

WHE
WIL

PRODUCT	RATE	
WILDCARD® XTRA	1 pint/A	F t s l v A f c
	1-1/2 to 2 pints/A	
	2 pints/A	
	1-1/2 pints/A	S e C A f c
	1-1/2 to 2 pints/A	S e C A f c
	Chemigation Only 2 pints/A	A f b s n o f f
	Post-harvest 3/4 to 2 pints/A	M o s N C t

irrigation water.
 uniform distribution of treated water.

consult manufacturers or other experts.
 under the supervision of the responsible

sprayed in the seedling stage.

BROADLEAF WEED SPECIES

(Chlorispora tenella)
(Senecio vulgaris)
(Ambrosia artemisiifolia)
(Anthemis arvensis)
(Lithospermum arvense)
(Fumaria officinalis)
(Ambrosia trifida)
(Sesbania exaltata)
(Lamium amplexicaule)
(Ipomoea hederacea)
(Scleranthus annuus)
(Kochia scoparia)
(Anthemis cotula)
(Polygonum aviculare)
(Tribulus terrestris)
(Ipomoea purpurea)
(Descurainia pinnata)
(Hemizonia spp.)
(Abutilon theophrasti)
(Raphanus raphanistrum)

spraying will not be controlled.

height.

regrowth may occur. Make applica-

WHEAT, BARLEY, OATS AND RYE WILDCARD® XTRA RECOMMENDATIONS

APPLICATION TIMING AND SPECIFIC COMMENTS			
PRODUCT	RATE	CROP	WEEDS
WILDCARD® XTRA	1 pint/A	Fall-seeded wheat, barley, oats and rye throughout the United States and spring seeded wheat, barley, oats and rye in Idaho, Oregon, Washington, Colorado, Wyoming and Montana. Apply to wheat, barley, oats and rye from the 3-leaf stage but before the crop reaches the boot stage.	MOST SUSCEPTIBLE BROADLEAF WEEDS: Apply to weeds up to the 8-leaf stage or 4 inches in height, whichever comes first. If weed forms rosette, apply before weeds exceed 2 inches in diameter. SUSCEPTIBLE BROADLEAF WEEDS: Apply to weeds up to the 4-leaf stage or 2 inches in height, whichever comes first. If weed forms rosette, apply before weeds exceed 1 inch in diameter.
	1-1/2 to 2 pints/A		
	2 pints/A		Apply to henbit, knawel and mayweed up to the 4-leaf stage or 2 inches in height, whichever comes first. Apply to kochia and tansy mustard for improved control when these weeds exceed the recommended stage of growth or are growing under cool, dry conditions.
	1-1/2 pints/A	Spring-seeded wheat and barley except Idaho, Oregon, Washington, Colorado, Montana, and Wyoming. Apply to wheat, barley, oats and rye from the 3-leaf stage but before the crop reaches the boot stage.	MOST SUSCEPTIBLE AND SUSCEPTIBLE BROADLEAF WEEDS: Apply to weeds that do not exceed the 8-leaf stage or 4 inches in height, whichever comes first. If weed forms rosette, apply before weeds exceed 2 inches in diameter. Apply to kochia up to 2 inches in height.
	1-1/2 to 2 pints/A	Spring-seeded wheat and barley except Idaho, Oregon, Washington, Colorado, Montana, and Wyoming. Apply to wheat, barley, oats and rye from the 3-leaf stage but before the crop reaches the boot stage.	Apply to kochia that is 2–4 inches height.
	Chemigation Only 2 pints/A	Apply to wheat, barley, oats and rye from the 3-leaf stage but before the boot stage. Apply through automated sprinkler irrigation systems with mechanical transfer loading system only. See "MIXING, LOADING AND HANDLING INSTRUCTIONS" section for complete details.	Apply to MOST SUSCEPTIBLE and SUSCEPTIBLE broadleaf weeds up to the 4-leaf stage, 2 inches in height or 1 inch in diameter, whichever comes first.
	Post-harvest 3/4 to 2 pints/A	Make applications following harvest of wheat, barley, oats and rye in the states of North Dakota, South Dakota, Minnesota, and Montana. Do not plant any rotational crop until the following use season.	Apply 3/4 to 1 pint/A to MOST SUSCEPTIBLE BROADLEAF WEEDS up to the 8-leaf stage or 4 inches in height, whichever comes first. Apply 1-1/2 to 2 pints/A to SUSCEPTIBLE BROADLEAF WEEDS up to the 4-leaf stage or 2 inches in height, whichever comes first. For control of both grasses and broadleaf weeds, tank mix WILDCARD® XTRA with Roundup® or Roundup® + 2,4-D such as WEEDONE® or WEEDAR® brand herbicides.



**WHEAT, BARLEY, OATS AND RYE
WILDCARD® XTRA TANK MIXTURE RECOMMENDATIONS**

APPLICATION TIMING AND SPECIFIC COMMENTS			
PRODUCT	RATE	CROP	WEEDS
WILDCARD® XTRA + Rhonox® (MCPA Ester)	3/4 to 2 pints/A + 1/4 to 1/2 pint/A	Apply to spring-seeded wheat, barley, oats and rye from tillering stage, but before boot stage.	For control of MOST SUSCEPTIBLE and SUSCEPTIBLE weeds and improved control of redroot pigweed and Kochia. Apply to weeds up to the 8-leaf stage, 3 inches in height or 2 inches in diameter, whichever comes first. Apply to Kochia and redroot pigweed up to 2 inches in height or diameter.
WILDCARD® XTRA + Glean® + nonionic surfactant	3/4 to 1-1/2 pints/A + 1/6 to 1/3 oz./A + 1 qt./100 gals. of water	Apply to wheat and barley from the 3-leaf stage but before the crop reaches the boot stage. Refer to Glean® label for crop rotation and other restrictions.	This tank mix improves control of broadleaf weeds such as henbit, tansy mustard and chickweed. Apply to weeds up to the 8-leaf stage, 4 inches in height or 2 inches in diameter, whichever comes first.
WILDCARD® XTRA + Finesse® + nonionic surfactant	3/4 to 1-1/2 pints/A + 1/6 to 1/3 oz./A + 1 qt./100 gals. of water	Apply to wheat and barley from the 3-leaf stage but before the crop reaches the boot stage. Refer to Finesse® label for crop rotation and other restrictions.	This tank mix improves control of broadleaf weeds such as henbit, tansy mustard and chickweed. Apply to weeds up to the 8-leaf stage, 4 inches in height or 2 inches in diameter, whichever comes first.
WILDCARD® XTRA + Ally® + nonionic surfactant	3/4 to 1-1/2 pints/A + 1/10 oz./A + 1 qt./100 gals. of water	Apply to wheat and barley from the 3-leaf stage but before the crop reaches the boot stage. Refer to Ally® label for crop rotation and other restrictions.	This tank mix improves control of broadleaf weeds such as henbit, tansy mustard and chickweed. Apply to weeds up to the 8-leaf stage, 4 inches in height or 2 inches in diameter, whichever comes first.
WILDCARD® XTRA + Banvel®	3/4 to 1-1/2 pints/A + 1/8 to 1/4 pint/A	Fall-seeded wheat from the 3-leaf stage but before jointing. Spring seeded wheat from the 3- to 5-leaf stage of growth.	This tank mix improves control of broad-leaves such as prostrate knotweed and Kochia. Apply to weeds up to the 8-leaf stage, 3 inches in height or 2 inches in diameter, whichever comes first. Apply to Kochia up to 2 inches in height or diameter.
WILDCARD® XTRA + Harmony® Extra + nonionic surfactant	3/4 to 1-1/2 pints/A + 3/10 to 1/2 oz./A + 1 qt./100 gals. of water	Winter wheat. Apply from the 3-leaf stage but before the 3rd node is detectable. Refer to the Harmony® Extra label for crop rotation and other restrictions. Spring wheat and barley. Apply after the 3-leaf, but before the 1st node is detectable. Refer to the Harmony® Extra label for crop rotation and other restrictions.	This tank mix improves control of broadleaf weeds such as henbit, chickweed and redroot pigweed. Apply to weeds up to the 8-leaf stage, 4 inches in height or across, whichever comes first.
WILDCARD® XTRA + Amber® + nonionic surfactant	3/4 to 1-1/2 pints/A + 0.28 to 0.56 oz./A + 0.25 to 0.5% v/v	Apply to wheat and barley from the 3-leaf stage, but before the flag leaf is visible. Refer to the Amber® label for crop rotation and other restrictions.	This tank mix improves control of broad-leaves such as henbit, tansy mustard, and pigweed. Apply to weeds up to the 4-leaf stage, 4 inches in height or 2 inches in diameter, whichever comes first.

**WHEAT, BA
WILDCARD® XTRA TANK M**

APPLICATION		
PRODUCT	RATE	CROP
WILDCARD® XTRA + Express® + nonionic surfactant	3/4 to 1-1/2 pints/A + 1/6 to 1/3 oz./A + 1 qt./100 gals. of water	Wheat and barley stage but before Refer to the Express rotation and other
WILDCARD® XTRA + Curtail® or Curtail® M	3/4 to 1-1/2 pints/A + 2 pints/A	Apply to wheat crop begins to detectable.
WILDCARD® XTRA + metribuzin (Sencor® or Lexone®)	1 pint/A + 1/8 to 3/16 lb. ai/A	Winter wheat in Washington. Apply growth has started with a minimum been established of joints in the when crop has frost damage, or
WILDCARD® XTRA + Avenger®	1 to 2 pints/A + 2-1/2 to 4 pints/A	Winter wheat. Refer to Avenger other restriction Spring Wheat. Refer to Avenger other restriction Barley. Three-
WILDCARD® XTRA + Assert®	1 to 1-1/2 pints/A + 1 to 1-1/2 pints/A	Apply to wheat 3-leaf stage but Refer to Assert and other restriction

RESTRICTIONS AND PRECAUTIONS: WHEAT, BARLEY

- Do not graze treated field within 45 days after application.
- Do not apply when crops are under moisture stress.
- Do not apply when crop canopy covers the weeds as for
- Reduced weed control may occur when weeds are stressed.
- Refer to labels of products used in tank mixture for additional restrictions.
- **Do not apply more than 0.75 lb. of WILDCARD® XTRA**

WHEAT, BARLEY, OATS AND RYE WILDCARD® XTRA TANK MIXTURE RECOMMENDATIONS (cont.)

APPLICATION TIMING AND SPECIFIC COMMENTS			
PRODUCT	RATE	CROP	WEEDS
WILDCARD® XTRA + Express® + nonionic surfactant	3/4 to 1-1/2 pints/A + 1/6 to 1/3 oz./A + 1 qt./100 gals. of water	Wheat and barley. Apply from the 3-leaf stage but before the flag is visible. Refer to the Express® label for crop rotation and other restrictions.	This tank mix improves control of broadleaf weeds such as henbit, chickweed, redroot pigweed and suppression of Canada thistle. Apply to annual weeds up to the 8-leaf stage, 4 inches in height or across, whichever comes first and to Canada thistle 4 to 8 inches tall with 2 to 6 inches of new growth.
WILDCARD® XTRA + Curtail® or Curtail® M	3/4 to 1-1/2 pints/A + 2 pints/A	Apply to wheat and barley after the crop begins to tiller up to the 1st node detectable.	This tank mix improves control of kochia, wild buckwheat and suppression of Canada thistle. Apply to annual broadleaf weeds up to the 8-leaf stage, 4 inches in height or 2 inches in diameter and to Canada thistle in the rosette to prebud stage.
WILDCARD® XTRA + metribuzin (Sencor® or Lexone®)	1 pint/A + 1/8 to 3/16 lb. ai/A	Winter wheat in Idaho, Oregon and Washington. Apply in spring after growth has started and secondary roots with a minimum of 3 to 4 tillers have been established, but before the forming of joints in the stem. Avoid application when crop has experienced winter kill, frost damage, disease or drought.	This tank mix improves control of broadleaf weeds such as chickweed, filaree, henbit. Apply to weeds up to the 4-leaf stage, 2 inches in height or diameter, whichever comes first. A recognized authority should be consulted concerning the use of this mixture in your area.
WILDCARD® XTRA + Avenge®	1 to 2 pints/A + 2-1/2 to 4 pints/A	Winter wheat. Four-leaf to tillering stage. Refer to Avenge® label for varietal and other restrictions. Spring Wheat. Five- to 6-leaf stage. Refer to Avenge® label for varietal and other restrictions. Barley. Three- to 7-leaf stage.	This tank mix will provide wild oat control in addition to broadleaves. Apply to wild oats in the 3- to 5-leaf stage and broadleaves that do not exceed the 4-leaf stage or rosettes of 1.5 inches in diameter. Avenge® use rates per acre are 2-1/2 pints (1-10 oats per sq. ft.), 3 pints (11-25 oats per sq. ft.) or 4 pints (more than 25 oats per sq. ft.).
WILDCARD® XTRA + Assert®	1 to 1-1/2 pints/A + 1 to 1-1/2 pints/A	Apply to wheat and barley from the 3-leaf stage but before boot stage. Refer to Assert® label for crop rotation and other restrictions.	This tank mix will provide wild oat control in addition to broadleaf weeds. Apply to wild oats at the 1- to 4-leaf stage and broadleaf weeds up to the 8-leaf stage, 4 inches in height or 2 inches in diameter, whichever comes first. Use Assert® at 1-1/2 pints/A west of the Rocky Mountains or if wild oats have initiated tillering. For spray volumes in excess of 10 GPA, add 0.3 fluid oz of nonionic surfactant for each gallon in excess of 10 GPA.

RESTRICTIONS AND PRECAUTIONS: WHEAT, BARLEY, OATS AND RYE

- Do not graze treated field within 45 days after application.
- Do not apply when crops are under moisture stress.
- Do not apply when crop canopy covers the weeds as poor control will result.
- Reduced weed control may occur when weeds are stressed from lack of moisture or cold temperatures.
- Refer to labels of products used in tank mixture for additional restrictions and precautions.
- **Do not apply more than 0.75 lb. of WILDCARD® XTRA acre per year.**

CONSERVATION REEF WILDCARD®

APPLICATION		
PRODUCT	RATE	CROP
WILDCARD® XTRA	1 to 2 pints/A	Apply to grass

RESTRICTIONS AND PRECAUTIONS: CRP AREAS

- Do not allow livestock to graze in treated areas or feed
- If legumes are included in CRP area planting, severe
- Do not apply more than 1.5 lbs. of WILDCARD® XTRA
- Do not apply more than 2 applications per year with

GRASSES GROWING WILDCARD® Seedling

APPLICATION TABLE			
PRODUCT	RATE PER ACRE	RATE PER 1000 SQ. FT.	CROP
WILDCARD® XTRA	1 to 3 pints	0.375 to 1.125 fl. oz.	Grasses growing in seedling
	Chemigation Only 2 pints/A 1 to 3 pints	0.75 fl. oz. 0.375 to 1.125 fl. oz.	Grasses growing in seedling

CONSERVATION RESERVE PROGRAM AREAS (CRP) WILDCARD® XTRA RECOMMENDATIONS

APPLICATION TIMING AND SPECIFIC COMMENTS			
PRODUCT	RATE	CROP	WEEDS
WILDCARD® XTRA	1 to 2 pints/A	Apply to grasses from the 3-leaf stage.	Apply 1 pint/A to MOST SUSCEPTIBLE and 1-1/2 to 2 pints/A to SUSCEPTIBLE broadleaf weeds up to the 8-leaf stage, 4 inches in height or 2 inches in diameter, whichever comes first.

RESTRICTIONS AND PRECAUTIONS: CRP AREAS

- Do not allow livestock to graze in treated areas or feed treated grass to livestock.
- If legumes are included in CRP area planting, severe injury may occur to legumes treated with **WILDCARD® XTRA**.
- Do not apply more than 1.5 lbs. of **WILDCARD® XTRA** ae/acre per year.
- Do not apply more than 2 applications per year with a minimum retreatment interval of 21 days.

GRASSES GROWN FOR SEED PRODUCTION

WILDCARD® XTRA RECOMMENDATIONS Seedling and Established Grasses

APPLICATION TIMING AND SPECIFIC COMMENTS				
PRODUCT	RATE PER ACRE	RATE PER 1000 SQ. FT.	CROP	WEEDS
WILDCARD® XTRA	1 to 3 pints	0.375 to 1.125 fl. oz.	Apply to established and newly seeded grasses grown for seed or sod production before the boot stage. Established grasses tolerant to WILDCARD® XTRA include bentgrasses, Kentucky Bluegrass, Fescues, Ryegrass, Bermudagrass, St. Augustinegrass and Zoysiagrass. WILDCARD® XTRA may also be used on seedling grasses such as Merion, Park, Delta, or common Kentucky Bluegrasses, Pennlawn, Chewings, Illahee or Alta Fescues, Orchardgrass, Highland, Seaside or Astoria Bentgrasses, perennial Ryegrasses, Bahiagrass and Zoysiagrass.	Refer to the GENERAL WEED LIST for a listing of susceptible broadleaf weeds. Optimal control will be attained when weeds are treated in the seedling stage (less than 4-leaf stage, 2 inches in height, or 1 inch in diameter).
	Chemigation Only 2 pints/A	0.75 fl. oz.	Apply to established and newly seeded grasses grown for seed or sod production before the boot stage. Apply through automated sprinkler irrigation systems with mechanical transfer loading system only. See "MIXING, LOADING AND HANDLING INSTRUCTIONS" section for complete details. Refer to the list of established grasses that are tolerant to WILDCARD® XTRA .	
	1 to 3 pints	0.375 to 1.125 fl. oz.		

(continued)

RESTRICTIONS AND PRECAUTIONS: GRASSES

- Do not allow livestock to graze in treated areas or
- Do not apply **WILDCARD® XTRA** to grasses grown

Note: Do not apply more than 1.5 Lbs. ae/acre maximum retreatment interval of 21 days.

FLAX (L) WILDCARD

APPLICATION		
PRODUCT	RATE	CROP
WILDCARD® XTRA	0.9 pint to 1 pint/A	Apply to flax Do not apply during

RESTRICTIONS AND PRECAUTIONS: FLAX (Lin)

- Do not apply more than 0.25 lb. of **WILDCARD® XT**
- Do not apply if temperatures are expected to exceed
- Unacceptable crop injury may occur following **WILDCARD® XT**
- Application under high humidity conditions can injure
- Unless otherwise instructed, do not apply **WILDCARD® XT**
- Do not use on ornamental flax.

control of broadleaf chickweed, redroot clover, Canada thistle. Apply up to the 8-leaf stage or across, whichever comes first. Canada thistle 4 to 6 inches in new

control of Kochia, Russian thistle, Canada thistle, broadleaf weeds up to the 8-leaf stage or across, whichever comes first. Canada thistle 4 to 6 inches in new

control of broadleaf chickweed, filaree, henbit, Canada thistle, 4-leaf stage, broadleaf weeds up to the 8-leaf stage, whichever comes first. Authority should be consulted for the use of this

control of wild oat in seedling stage. Apply to wild oats and broadleaves up to the 4-leaf stage or across, whichever comes first. Oats are 2-1/2 pints to 3 pints (11-25 oats per acre) more than 25 oats per

control of wild oat in seedling stage. Apply to wild oats and broadleaves up to the 8-leaf stage, whichever comes first. Use Assert® at the Rocky Mountains. Excess of 10 GPA, ionic surfactant for of 10 GPA.



(P)

OST SUSCEPTIBLE
/A to SUSCEPTIBLE
) to the 8-leaf stage,
r 2 inches in diameter,
rst.

id® XTRA.

GENERAL WEED
isting of susceptible
veeds.
ontrol will be attained
is are treated in the
age (less than 4-leaf
ches in height,
diameter).

(continued)

RESTRICTIONS AND PRECAUTIONS: GRASSES GROWN FOR SEED OR SOD PRODUCTION

- Do not allow livestock to graze in treated areas or feed treated grasses to livestock.
- Do not apply **WILDCARD® XTRA** to grasses grown for seed production with backpack or hand-held application equipment.

Note: Do not apply more than 1.5 Lbs. ae/acre per year. Do not apply more than 2 applications per year with a minimum retreatment interval of 21 days.

**FLAX (*Linum usitatissimum* only)
WILDCARD® XTRA RECOMMENDATIONS**

APPLICATION TIMING AND SPECIFIC COMMENTS			
PRODUCT	RATE	CROP	WEEDS
WILDCARD® XTRA	0.9 pint to 1 pint/A	Apply to flax 2 to 8 inches in height. Do not apply WILDCARD® XTRA to flax during or after the bud stage.	Apply to MOST SUSCEPTIBLE weeds that do not exceed the 4-leaf stage, 2 inches in height or 1 inch in diameter, whichever comes first.

RESTRICTIONS AND PRECAUTIONS: FLAX (*Linum usitatissimum* only)

- Do not apply more than 0.25 lb. of **WILDCARD® XTRA** ae/acre per year
- Do not apply if temperatures are expected to exceed 85°F at or 3 days following application or crop injury may occur.
- Unacceptable crop injury may occur following **WILDCARD® XTRA** application to flax grown on high organic, peat-type soils.
- Application under high humidity conditions can injure flax.
- Unless otherwise instructed, do not apply **WILDCARD® XTRA** to flax with crop oil concentrate, surfactants or nitrogen solutions.
- Do not use on ornamental flax.

**CONDITIONS
AND LIMITATION**

Read the Conditions of Sale – Warranty and terms are not acceptable, return the product

The directions on this label are believed to be reliable for the crop to which the product is applied. The user is responsible to follow the label directions or good agricultural practices (the "Company") or seller. In addition, failure to follow the label directions may void the Company's warranty. The Company warrants that this product conforms to the label directions for use subject to the limitations of the label. Consistent with applicable law, the Company makes no warranty concerning the product, including no implied warranty of merchantability shall be implied by law.

To the extent consistent with applicable law, the handling or use of this product shall be limited to:

1. Refund of the purchase price paid by buyer or user
2. Replacement of the product used.

To the extent consistent with applicable law, the Company is not liable for special, indirect, incidental, or consequential damages. The Company and the seller offer this product with the limitation of warranty, liability and remedies.

© Copyright Helena Holding Company, 2012.

Assert®, Banvel®, and Avenge® are registered trademarks of Helena Holding Company. Sencor® is a registered trademark of Bayer AG, Germany. Raindrop® is a registered trademark of Delavan Corporation. Curtail® is a registered trademark of Dow AgroSciences. Ally®, Express®, Finesse®, Glean®, Harmony® and Le Amber® are registered trademarks of Syngenta Crop Protection, Inc. Rhonox®, WEEDAR® and WEEDONE® are registered trademarks of Monsanto Company. Roundup® is a registered trademark of Monsanto Company. **WILDCARD®** is a registered trademark of Helena Holding Company.



application equipment.
ns per year with a mini-

USCEPTIBLE weeds
d the 4-leaf stage,
or 1 inch in diameter,
first.

jury may occur.
janic, peat-type soils.

ints or nitrogen solutions.

CONDITIONS OF SALE – LIMITED WARRANTY AND LIMITATIONS OF LIABILITY AND REMEDIES

Read the Conditions of Sale – Warranty and Limitations of Liability and Remedies before using this product. If the terms are not acceptable, return the product, unopened, and the full purchase price will be refunded.

The directions on this label are believed to be reliable and should be followed carefully. Insufficient control of pests and/or injury to the crop to which the product is applied may result from the occurrence of extraordinary or unusual weather conditions or the failure to follow the label directions or good application practices, all of which are beyond the control of Helena Chemical Company (the “Company”) or seller. In addition, failure to follow label directions may cause injury to crops, animals, man or the environment. The Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purpose referred to in the directions for use subject to the factors noted above which are beyond the control of the Company. To the extent consistent with applicable law, the Company makes no other warranties or representations of any kind, express or implied, concerning the product, including no implied warranty of merchantability or fitness for any particular purpose, and no such warranty shall be implied by law.

To the extent consistent with applicable law, the exclusive remedy against the Company for any cause of action relating to the handling or use of this product shall be limited to, at Helena Chemical Company’s election, one of the following:

1. Refund of the purchase price paid by buyer or user for product bought, or
2. Replacement of the product used.

To the extent consistent with applicable law, the Company shall not be liable and any and all claims against the Company are waived for special, indirect, incidental, or consequential damages or expense of any nature, including, but not limited to, loss of profits or income. The Company and the seller offer this product and the buyer and user accept it, subject to the foregoing conditions of sale and limitation of warranty, liability and remedies.

© Copyright Helena Holding Company, 2012.

Assert®, Banvel®, and Avenge® are registered trademarks of BASF Corp.

Sencor® is a registered trademark of Bayer AG, Germany.

Raindrop® is a registered trademark of Delavan Corporation.

Curtail® is a registered trademark of Dow AgroSciences.

Ally®, Express®, Finesse®, Glean®, Harmony® and Lexone® are registered trademarks of E.I. DuPont DeNemours and Co.

Amber® is a registered trademark of Syngenta Crop Protection.

RhonoX®, WEEDAR® and WEEDONE® are registered trademarks of Nufarm Inc.

Roundup® is a registered trademark of Monsanto Company.

WILDCARD® is a registered trademark of Helena Holding Company.

**Y
IES**

using this product. If the
e refunded.

ontrol of pests and/or injury to
weather conditions or the fail-
f Helena Chemical Company
ials, man or the environment.
easonably fit for the purpose
f the Company. To the extent
ind, express or implied, con-
rpose, and no such warranty

ause of action relating to the
following:

inst the Company are waived
it limited to, loss of profits or
e foregoing conditions of sale

lemours and Co.



**FOR CONTROL OF CERTAIN BROAD
CONSERVATION RESERVE PROGRAM**

ACTIVE INGREDIENT:

Octanoic acid of ester of bromoxynil (3,5-dibromo-4-isooctyl ester of 2-methyl-chlorophenoxyacetic acid)

INERT INGREDIENTS

TOTAL

Contains Petroleum Distillate.

*Bromoxynil octanoate equivalent to 21.8% of

**Equivalent to 21.8% 2-methyl-chlorophenoxy.

**KEEP OUT
CAUTION**

Si usted no entiende la etiqueta
(If you do not understand the label)

**For Chemical Spill, Leaking Container
For Medical Attention**

IF SWALLOWED:	■ Call a poison control center or doctor immediately with product name in hand. ■ Do not induce vomiting unless told to do so by a poison control center or doctor. ■ Do not drink milk.
IF IN EYES:	■ Hold eye open and rinse slowly and gently with water for at least 15 minutes. ■ Remove contact lenses, if present and easy to remove. ■ Continue rinsing. ■ Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING:	■ Take off contaminated clothing. ■ Wash thoroughly with soap and water. ■ Call a poison control center or doctor for treatment advice.
IF INHALED:	■ Move person to fresh air. ■ If person is not breathing, call 911 or a poison control center or doctor immediately for treatment advice. ■ If breathing is difficult, call a poison control center or doctor immediately for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going to a hospital for emergency medical treatment information.

Note to Physician: May pose an aspiration pneumonia hazard if inhaled.

SEE INSIDE BOOKLET FOR ADDITIONAL INFORMATION

EPA Reg. No. 5905-550

EPA

**HELENA
225 SCHILLING BOULEVARD**

Job #59986



**FOR CONTROL OF CERTAIN BROADLEAF WEEDS IN SMALL GRAINS (WHEAT, BARLEY, OATS AND RYE),
CONSERVATION RESERVE PROGRAM (CRP) AREAS, GRASSES GROWN FOR SEED PRODUCTION AND FLAX**

ACTIVE INGREDIENT:

Octanoic acid of ester of bromoxynil (3,5-dibromo-4-Hydroxybenzotrile) 31.7%*
Isocetyl ester of 2-methyl-chlorophenoxyacetic acid** 34.0%**

INERT INGREDIENTS 34.3%

TOTAL 100.0%

Contains Petroleum Distillate.

Patent No. 6,232,272

*Bromoxynil octanoate equivalent to 21.8% of bromoxynil or not less than 2.0 pounds of bromoxynil per gallon.

**Equivalent to 21.8% 2-methyl-chlorophenoxyacetic acid or not less than 2.0 pounds MCPA acid per gallon.

**KEEP OUT OF REACH OF CHILDREN
CAUTION – PRECAUCIÓN**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

**For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC 1-800-424-9300
For Medical Emergencies Only, Call 1-877-325-1840**

FIRST AID

IF SWALLOWED:	<ul style="list-style-type: none"> ■ Call a poison control center or doctor immediately for treatment advice. ■ Have person sip a glass of water if able to swallow. ■ Do not induce vomiting unless told to do so by the poison control center or doctor. ■ Do not give anything by mouth to an unconscious person.
IF IN EYES:	<ul style="list-style-type: none"> ■ Hold eye open and rinse slowly and gently with water for 15-20 minutes. ■ Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. ■ Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> ■ Take off contaminated clothing. ■ Rinse skin immediately with plenty of water for 15-20 minutes. ■ Call a poison control center or doctor for treatment advice.
IF INHALED:	<ul style="list-style-type: none"> ■ Move person to fresh air. ■ If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. ■ Call a poison control center or doctor for further treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment information.

Note to Physician: May pose an aspiration pneumonia hazard. Contains petroleum distillate.

SEE INSIDE BOOKLET FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND COMPLETE DIRECTIONS FOR USE.

EPA Reg. No. 5905-550

EPA Est. No. 228-IL-1

Product of United Kingdom

AD 022712

**Manufactured For
HELENA CHEMICAL COMPANY
225 SCHILLING BOULEVARD, SUITE 300 • COLLIERVILLE, TENNESSEE 38017**