



























### Willowood Clomazone 3ME Rates and Weeds Controlled (Applied Alone)

Rate	Grasses Controlled
1 1/3 pts. (0.50 lb. AI)	Suppresses some annual grasses
2.0 pts. (0.75 lb. AI)	Barnyardgrass, Broadleaf signalgrass, Large & Smooth Crabgrass, Giant & Green Foxtail, Goosegrass, Panicum (Common, Fall, & Texas), Johnsongrass seedling, Field Sandbur (partial control)
2 2/3 pts. (1.0 lb. AI)	All grasses listed above PLUS: Cupgrass (Southwestern & Woolly), Field Sandbur, Bermudagrass (partial control), Red Rice, Itchgrass*
3 1/3 pts. (1.25 lbs. AI)	All grasses listed above PLUS: Foxtail (Purple, Robust, Yellow), Jungerice (partial control), Shattercane (partial control), Wild Proso Millet (partial control)

\*Itchgrass: Do not incorporate Willowood Clomazone 3ME. Use 3 1/3 pints for high weed pressure and heavy soil.

### Willowood Clomazone 3ME Rates and Broadleaf Weeds Controlled (Applied Alone)

Rate	Broadleaf Weeds Controlled
1 1/3 pts.* (0.50 lb. AI) 1 2/3 pts.*	Velvetleaf Spurred Anoda *Use higher rate for heavy weed pressure or heavy soil
2.0 pts. (0.75 lb. AI)	Broadleaf weeds listed above PLUS: Common Ragweed (partial control), Jimsonweed (partial control), Lambsquarter, Pennsylvania Smartweed (partial control), Prickly Sida, Purslane, Redweed, Venice Mallow
2 2/3 pts. (1.0 lb. AI)	Broadleaf weeds listed above PLUS: Cocklebur, Common Ragweed, Dayflower, Florida Beggarweed, Florida Pusley, Jimsonweed, Kochia (partial control), Redvine (partial control), Tropic Croton, Wild Poinsettia**
3 1/3 pts. (1.25 lbs. AI)	Broadleaf weeds listed above PLUS: Balloon Vine (partial control), Black nightshade (partial control), Curly Dock, Joint Vetch (partial control), Kochia, Pitted Morning Glory, Pennsylvania Smartweed, Redroot Pigweed (partial control), Smooth Pigweed (partial control)

\*\*Use in AR/LA/MS/TX/OK only - for high pressure and heavy soils use 3 1/3 pints.

Use higher specified rates (a minimum of 2 2/3 pints per acre) in fields where weed pressure is heavy, and the field has in conservation programs. Select lower to higher rates based on lighter to heavier soil types. Use a minimum application rate of 3 1/3 pints per acre for cocklebur or common ragweed control in fields that have previously been in conservation programs.

### **Willowood Clomazone 3ME Use with Other Preemergent Herbicides Registered for Use on Cotton**

Apply Willowood Clomazone 3ME in combination with other herbicides registered for use in cotton to broaden the weed control spectrum. Refer to the other herbicide label for additional information. Follow all application precautions, rotational crop restrictions and replanting instructions of each product's label when using tank mixtures. Follow the most restrictive precautions and instructions for all products used.

#### **Crop Rotation Restrictions**

Rotate to crops listed below to avoid crop injury. Refer to sections headed "Rotational Cropping Precautions".

**Note:** When using Willowood Clomazone 3ME with other registered herbicides always refer to rotational restrictions and precautions on the other product's label.

- Cover crops can be planted at any time, but stand reduction can occur in some areas. Do not harvest for food, graze, or feed cover crops planted less than 9 months after treatment with Willowood Clomazone 3ME.
- Maximum Application Rate: 1.25 lbs. active ingredient per acre per season.
- **DO NOT** apply Willowood Clomazone 3ME within 65 days of harvest.
- **DO NOT** allow livestock to graze on cotton forage or trash, feed, treated cotton forage, or trash to livestock.

### Willowood Clomazone 3ME Rotational Crop Restrictions for Cotton

Rate	Interval	Crop
1 1/3 pts. (0.50 lb. AI)	Anytime	Cabbage (transplanted), Cotton*, Peas, Peppers, Pumpkins (processing), Rice, Soybeans, Squash, Sweet Potato, Tobacco, Tuberous & Corm Vegetables
	9 Months	All crops listed above PLUS: Cabbage (direct seeded), Field Corn, Popcorn, Seed Corn, Sweet Corn, Cotton, Cucurbits, Dry Beans, Peanuts, Potatoes, Snap Beans, Sorghum, Sugar Beets, Tomatoes (Transplanted)
	12 Months	All crops
2 - 2 2/3 pts. (0.75 - 1.0 lb. AI)	Anytime	Cotton*, Peppers, Rice, Soybeans, Winter Squash (2.0 pt.), Sweet Potato, Tobacco, Tuberous & Corm Vegetables
	9 Months	All crops listed above PLUS: Cabbage, Field Corn, Popcorn, Seed Corn, Sweet Corn, Cotton, Cucurbits, Dry Beans, Peanuts, Peas, Potatoes, Pumpkins (processing), Rice, Snap Beans, Sorghum, Summer Squash, Sugar Beets, Tomatoes (Transplanted)
	12 Months	All crops
3 1/3 pts. (1.25 lbs. AI)	Anytime	Cotton*, Soybeans, Sweet Potato, Tobacco, Tuberous & Corm Vegetables
	9 Months	All crops listed above PLUS: Cabbage, Field Corn, Popcorn, Seed Corn, Sweet Corn, Cotton, Cucurbits, Dry Beans, Peanuts, Peas, Peppers, Potatoes, Pumpkins (processing), Rice, Snap Beans, Sorghum, Squash, Sugar Beets, Tomatoes (Transplanted)
	12 Months	All crops

\*Cotton: Refer to "Requirements for Planting Time Applications" and "Replanting Instructions" for additional instructions.

# RICE

Apply Willowood Clomazone 3ME as a surface broadcast treatment prior to weed emergence to control annual grass weeds in dry-seeded rice.

## Use Precautions for Rice

Off-site movement of Willowood Clomazone 3ME by spray drift or vapors can cause foliar whitening or plant yellowing. Follow all precautions and instructions in the APPLICATION PRECAUTIONS, SPRAY DRIFT PRECAUTIONS and SPRAY DRIFT MANAGEMENT sections of this label before applying this product.

## Preemergent Surface Broadcast Application

- Apply Willowood Clomazone 3ME 14 days before planting or up to 7 days after planting.
- Apply prior to weed emergence.
- Apply with ground equipment in a minimum of 10 to 40 gallons of water per acre at the rate of 2/3 to 2 1/8 pints (0.25 to 0.8 lb. AI) per acre depending upon the soil texture.
- See the table below for application rates and weeds controlled.
- Use the higher specified rate for heavy soils for maximum control.
- For preemergent broadcast treatments, refer to the directions for broadcast applications in the GROUND APPLICATION INSTRUCTIONS section.

## Early Postemergence Application

- Apply Willowood Clomazone 3ME after planting as an early postemergence treatment to rice at the 1- to 5-leaf stage.
- Use ground equipment with nozzles that produce a coarse spray with a minimum of 10-40 gallons of water per acre.
- See the table below for specific rates and weeds controlled.
- To control existing grass at the application site, include a postemergence herbicide registered to control grass species in rice. Consult postemergence herbicide label for specific directions regarding use rates and stage of weeds and crop.

## Replanting Precautions

- Rice can be replanted in fields treated with Willowood Clomazone 3ME if the crop fails to produce a uniform stand at initial planting.
- If Willowood Clomazone 3ME is tank mixed with a labeled product, refer to the replant instructions for that product.

## Replanting Restrictions

- **DO NOT** make a second application of Willowood Clomazone 3ME.
- **DO NOT** replant treated fields with any crop at intervals that are inconsistent with the ROTATIONAL CROP INSTRUCTIONS listed above. If a tank mix is used, refer to the product's label for any additional crop rotation instructions.



## Willowood Clomazone 3ME Applied Alone:

### Single Application Preemergence or Postemergence

**Rates and Weeds Controlled: Select lower to higher rates based on lighter to heavier soils**

Broadcast Rates per Acre	Soil Texture
11 - 14 fl. oz. (0.25 - 0.33 lb. Al)	Coarse (light) Soils: Loamy sand, sandy loam
17 - 21 fl. oz. (0.4 - 0.5 lb. Al)	Medium Soils: Loam, silt, silt loam, sandy clay, sandy clay loam
21 - 34 fl. oz. (0.5 - 0.8 lb. Al)	Fine (heavy) Soils: Silty clay, clay loam, silty clay loam, clay

### List of Grasses Controlled

Barnyardgrass (Watergrass) (*Echinochloa crus-galli*, *E. colonum*)

Broadleaf signalgrass (*Brachiaria platyphylla*)

Crabgrass (Large and Smooth) (*Digitaria* spp.)

Panicum (*Panicum* spp.)

—Common

—Fall

—Texas

Sprangletop (*Leptochloa* spp.)

**If levees are pulled after this product has been applied, partial weed control may result. Make additional applications of post-emerge herbicide if this occurs.**

### Split Applications

Pre-emergent surface broadcast applications and early postemergence applications (out to the 2-leaf stage) of Willowood Clomazone 3ME can be made in the same season to control the above weeds provided that the total seasonal application rate does not exceed 0.8 lb. Al/A . See the table below for rates.

**Willowood Clomazone 3ME Split Application Rate Table**

Broadcast Rates per Acre		Soil Texture
Pre-Emergent	Post-Emergent	
2/3 pt. (0.25 lb. Al)	2/3 pt. (0.25 lb. Al)	Coarse (light) Soils: Loamy sand, sandy loam
4/5 - 7/8 pt. (0.3 - 0.35 lb. Al)	2/3 - 4/5 pt. (0.25 - 0.3 lb. Al)	Medium Soils: Loam, silt, silt loam, sandy clay, sandy clay loam
1 1/8 - 1 1/3 pts. (0.4 to 0.5 lb. Al)	13 - 17 fl. oz. (0.3 to 0.4 lb. Al)	Fine (heavy) Soils: Silty clay, clay loam, silty clay loam, clay

## Split-Application Precautions

- Stand loss or other rice crop injury can occur from applications of Willowood Clomazone 3ME to fields that have been precision leveled with deep cuts.
- Contact rice specialists in your area to learn about soil amending practices designed to reduce potential for crop injury in precision leveled fields.
- Cover crops can be planted at any time, but stand reduction can occur in some areas.
- Rotate to crops as listed below to avoid crop injury. See the section “Rotational Cropping Precautions”.

**Note:** Always refer to rotational restrictions and precautions on the other product’s label if using Willowood Clomazone 3ME in combination with other registered herbicides.

## Split-Application Crop Rotation Restrictions

- **DO NOT** harvest for food, graze, or feed cover crops planted less than 9 months after treatment with Willowood Clomazone 3ME.
- **DO NOT** make split-applications in water-seeded rice.

Rate	Interval	Crop
11 - 34 fl. oz. (0.25 - 0.8 lb. AI)	Anytime	Cotton*, Peas, Peppers, Pumpkins (processing), Rice, Soybeans, Squash, Sweet Potato, Tobacco, Tuberous & Corn Vegetables
	9 Months	All crops listed above PLUS: Field Corn, Popcorn, Seed Corn, Sweet Corn, Cotton, Cucurbits, Dry Beans, Peanuts, Potatoes, Snap Beans, Sorghum, Sugar Beets, Tomatoes (Transplanted)
	12 Months	All crops

\*Cotton: Refer to “Requirements for Planting Time Applications” and “Replanting Instructions” for additional instructions.

## Crop Rotation Restrictions

- **DO NOT** apply Willowood Clomazone 3ME to rice fields where crayfish or catfish farming are concurrently included in the cultural practices.
- **DO NOT** use water containing this product’s residues from rice cultivation to irrigate food or feed crops that are not registered for use with Willowood Clomazone 3ME.
- Maximum Application Rate: 34 fl. oz. AI of product (0.8 lb. AI Willowood Clomazone 3ME) per acre per use season.

## SOYBEANS

Apply Willowood Clomazone 3ME as a preemergent soil applied treatment to control annual grasses and broadleaf weeds in soybeans. Tank mix Willowood Clomazone 3ME with other preemergent soybean herbicides for greater weed control. Willowood Clomazone 3ME can be tank mixed with Canopy®, Dual®, Lasso®, Lexone®, Lorox®, Lorox Plus®, Prowl®, Pursuit®, Scepter®, Sencor®, Sonalan®, Squadron®, or other registered herbicides. Follow all precautions, instructions, and rotational cropping restrictions of each product's label when tank mixing, including any references to potential carryover and crop injury warnings or restrictions. The most restrictive precautions and instructions must be followed for all products used.

Water or liquid fertilizer can be a carrier for Willowood Clomazone 3ME if applied alone, or if tank mixed with the herbicides listed above, unless directions for use specifically state otherwise.

### Replanting Precautions for Soybeans

- Soybeans can be replanted in fields treated with Willowood Clomazone 3ME alone (or with specified tank mixtures) if initial seeding of soybeans fails to produce a stand.
- Refer to the soybean replant instructions for that product if tank mixing with a labeled product.

### Replanting Restrictions for Soybeans

- **DO NOT** make a second application of Willowood Clomazone 3ME.
- **DO NOT** replant treated fields with any crop at intervals that are not consistent with the Rotational Crop Instructions listed on this label.

**Willowood Clomazone 3ME can be used on Soybeans in the Northern Region of the map below**  
**Northern Region**



### Use Precautions for Soybeans

Off-site movement of Willowood Clomazone 3ME by spray drift or vapors can cause foliar whitening or plant yellowing. Follow all precautions and instructions in the APPLICATION PRECAUTIONS, SPRAY DRIFT PRECAUTIONS and SPRAY DRIFT MANAGEMENT sections of this label before applying this product.

### Willowood Clomazone 3ME Rates and Grasses Controlled (Applied Alone)

Rate	Grasses Controlled
1 1/3 pts. (0.50 lb. AI)	Suppresses some annual grasses
2.0 pts. (0.75 lb. AI)	All grasses listed above PLUS: Barnyardgrass, Large & Smooth Crabgrass, Field Sandbur, Foxtail (Giant, Green, Robust, Yellow), Goosegrass, Panicum (Common, Fall, & Texas), Johnsongrass seedling (partial control)
2 2/3 pts. (1.0 lb. AI)	All grasses listed above PLUS: Bermudagrass (partial control), Broadleaf Signalgrass, Southwestern Cupgrass, Woolly Cupgrass (partial control), Seedling Johnsongrass, Shattercane (partial control), Wild Proso Millet (partial control)

### Willowood Clomazone 3ME Rates and Broadleaf Weeds Controlled (Applied Alone)

Rate	Broadleaf Weeds Controlled
1 1/3 pts. (0.50 lb. AI) 1 2/3 pts.*	Velvetleaf Spurred Anoda *Use higher rate for heavy weed pressure or heavy soil
2.0 pts. (0.75 lb. AI)	Broadleaf weeds listed above PLUS: Common Ragweed (partial control), Galinsoga, Jimsonweed (partial control), Lambsquarter, Prickly Sida, Purslane, Ladysthumb Smartweed (partial control), Pennsylvania Smartweed (partial control), Venice Mallow
2 2/3 pts. (1.0 lb. AI)	Broadleaf weeds listed above PLUS: Black Seeded Plantain, Cocklebur (partial control), Common Ragweed, Jimsonweed, Kochia, Ladysthumb Smartweed, Pennsylvania Smartweed, Tropic Croton

Use higher specified rates (a minimum of 2 2/3 pints per acre) in fields where weed pressure is heavy, and field has been in conservation programs. Select lower to higher rates based on lighter to heavier soil types.

#### Soybean - Northern Region Crop Rotation Precautions

- Cover crops can be planted at any time, but stand reduction can occur in some areas.
- Rotate to crops as listed below to avoid crop injury. See the section “Rotational Cropping Precautions”.

**Note:** Always refer to rotational restrictions and precautions on the other product’s label if using Willowood Clomazone 3ME in combination with other registered herbicides.

#### Soybean - Northern Region Crop Rotation Restrictions

- **DO NOT** harvest for food, graze, or feed cover crops planted less than 9 months after treatment with Willowood Clomazone 3ME.
- **DO NOT** allow livestock to graze on soybean vines treated with this product.
- **DO NOT** feed soybean vines or soybean vine trash treated with this product to livestock.

## Willowood Clomazone 3ME Rotational Crop Restrictions for Soybeans in the Northern Region

Rate	Interval	Crop
1 1/3 pts. (0.50 lb. AI)	Anytime	Cabbage (transplanted), Cotton*, Peas, Peppers, Pumpkins (processing), Rice, Soybeans, Squash, Sweet Potato, Tobacco, Tuberous & Corm Vegetables
	9 Months	All crops listed above PLUS: Cabbage (Direct Seeded), Field Corn, Popcorn, Seed Corn, Sweet Corn, Cotton, Cucurbits, Dry Beans, Peanuts, Potatoes, Snap Beans, Sorghum, Sugar Beets, Tomatoes (Transplanted)
	12 Months	All crops listed above PLUS: Tomatoes (all), Wheat
	16 Months	All crops
2.0 pts. (0.75 lb. AI)	Anytime	Cotton*, Peppers, Soybeans, Rice, Soybeans, Winter Squash, Sweet Potato, Tobacco, Tuberous & Corm Vegetables
	9 Months	All crops listed above PLUS: Cabbage, Field Corn, Popcorn, Seed Corn, Sweet Corn, Cotton, Cucurbits, Dry Beans, Peanuts, Peas, Potatoes, Pumpkins (processing), Rice, Snap Beans, Sorghum, Summer Squash, Sugar Beets, Tomatoes (Transplanted)
	12 Months	All crops listed above PLUS: Tomatoes (all), Wheat
	16 Months	All crops
2 2/3 pts. (1.0 lb. AI)	Anytime	Cotton*, Peppers, Soybeans, Sweet Potato, Tobacco, Tuberous & Corm Vegetables
	9 Months	All crops listed above PLUS: Cabbage, Field Corn, Popcorn, Cotton, Cucurbits, Dry Beans, Peanuts, Peas, Potatoes, Pumpkins (processing), Rice, Snap Beans, Sorghum, Squash, Sugar Beets, Tomatoes (Transplanted)
	12 Months	All crops listed above PLUS: Seed Corn, Sweet Corn, Tomatoes (all), Wheat
	16 Months	All crops

\*Cotton: Refer to "Requirements for Planting Time Applications" and "Replanting Instructions" for additional instructions.

**Willowood Clomazone 3ME can be used on Soybeans in the Southern Region of the map below**



**Southern Region**

**Additional Precautions For Soybean Applications in the Southern Region:**

Off-site movement of Willowood Clomazone 3ME by spray drift or vapors can cause foliar whitening or plant yellowing. Follow all precautions and instructions in the APPLICATION PRECAUTIONS, SPRAY DRIFT PRECAUTIONS and SPRAY DRIFT MANAGEMENT sections of this label before applying this product. Use the lowest possible boom height while maintaining a uniform spray pattern, in conjunction with nozzle type, size, operating pressure and volume that meet a droplet size classification of **coarse or greater**.

**DO NOT** spray within 300 feet of desirable plants.

**Willowood Clomazone 3ME Rates and Weeds Controlled**

<b>Rate</b>	<b>Grasses Controlled</b>
1 1/3 pts. (0.50 lb. AI)	Suppresses some annual grasses
2.0 pts. (0.75 lb. AI)	Barnyardgrass, Broadleaf signalgrass, Large & Smooth Crabgrass, Giant & Green Foxtail, Goosegrass, Panicum (Common, Fall, & Texas), Johnsongrass Seedling, Field Sandbur (partial control)
2 2/3 pts. (1.0 lb. AI)	All grasses listed above PLUS: Cupgrass (Southwestern & Woolly), Field Sandbur, Bermudagrass (partial control), Red Rice, Itchgrass*
3 1/3 pts. (1.25 lbs. AI)	All grasses listed above PLUS: Foxtail (Purple, Robust, Yellow), Junglerice (partial control), Shattercane (partial control), Wild Proso, Millet (partial control)

\*Itchgrass: Do not incorporate Willowood Clomazone 3ME. Use 3 1/3 pints for high weed pressure and heavy soil.

## Willowood Clomazone 3ME Rates and Broadleaf Weeds Controlled

Rate	Broadleaf Weeds Controlled
1 1/3 pts. (0.50 lb. AI) 1 2/3 pts.*	Velvetleaf Spurred Anoda *Use higher rate for heavy weed pressure or heavy soil
2.0 pts. (0.75 lb. AI)	Broadleaf weeds listed above PLUS: Common Ragweed (partial control), Jimsonweed (partial control), Lambsquarter, Pennsylvania Smartweed (partial control), Prickly Sida, Purslane, Redweed, Venice Mallow
2 2/3 pts. (1.0 lbs. AI)	Broadleaf weeds listed above PLUS: Cocklebur, Common Ragweed, Dayflower, Florida Beggarweed, Florida Pusley, Jimsonweed, Kochia (partial control), Redvine (partial control), Tropic Croton, Wild Poinsettia**
3 1/3 pts. (1.25 lbs. AI)	Broadleaf weeds listed above PLUS: Balloon Vine (partial control), Black nightshade (partial control), Curly Dock, Joint Vetch (partial control), Kochia, Pitted Morning Glory, Pennsylvania Smartweed, Redroot Pigweed (partial control), Smooth Pigweed (partial control), Prostrate Spurge

\*\*Use in AR/LA/MS/TX/OK only - for high pressure and heavy soils use 3 1/3 pints.

Use higher specified rates (a minimum of 2 2/3 pints per acre) in fields where weed pressure is heavy, or the field has previously been in conservation programs. Select lower to higher rates based on lighter to heavier soil types. Use a minimum application rate of 3 1/3 pints per acre for cocklebur or common ragweed control in fields that have previously been in conservation programs.

### Soybean - Southern Region Crop Rotation Precautions

- Cover crops can be planted at any time, but stand reduction can occur in some areas.
- Rotate to crops as listed below to avoid crop injury. See the section “Rotational Cropping Precautions”.

**Note:** Always refer to rotational restrictions and precautions on the other product’s label if using Willowood Clomazone 3ME in combination with other registered herbicides.

### Soybean - Southern Region Crop Rotation Restrictions

- **DO NOT** harvest for food, graze, or feed cover crops planted less than 9 months after treatment with Willowood Clomazone 3ME.
- **DO NOT** allow livestock to graze on soybean vines treated with this product.
- **DO NOT** feed soybean vines or soybean vine trash treated with this product to livestock.

## Willowood Clomazone 3ME Rotational Crop Restrictions for Soybeans in the Southern Region

Rate	Interval	Crop
1 1/3 pts. (0.50 lb. AI)	Anytime	Cabbage (transplanted), Cotton*, Peas, Peppers, Pumpkins (processing), Rice, Soybeans, Squash, Sweet Potato, Tobacco, Tuberos & Corm Vegetables
	9 Months	All crops listed above PLUS: Cabbage (Direct Seeded), Field Corn, Popcorn, Seed Corn, Sweet Corn, Cotton, Cucurbits, Dry Beans, Peanuts, Potatoes, Snap Beans, Sorghum, Sugar Beets, Tomatoes (Transplanted)
	12 Months	All crops
2.0 pts. - 2 2/3 pts. (0.75 - 1.0 lb. AI)	Anytime	Cotton*, Peppers, Rice, Soybeans, Winter Squash (2.0 pts.), Sweet Potato, Tobacco, Tuberos & Corm Vegetables
	9 Months	All crops listed above PLUS: Cabbage, Field Corn, Popcorn, Seed Corn, Sweet Corn, Cotton, Cucurbits, Dry Beans, Peanuts, Peas, Potatoes, Pumpkins (processing), Rice, Snap Beans, Sorghum, Summer Squash, Sugar Beets, Tomatoes (Transplanted)
	12 Months	All crops
3 1/3 pts. (1.25 lbs. AI)	Anytime	Cotton*, Soybeans, Sweet Potato, Tobacco, Tuberos & Corm Vegetables
	9 Months	All crops listed above PLUS: Cabbage, Field Corn, Popcorn, Seed Corn, Sweet Corn, Cotton, Cucurbits, Dry Beans, Peanuts, Peas, Peppers, Potatoes, Pumpkins (processing), Rice, Snap Beans, Sorghum, Squash, Sugar Beets, Tomatoes (Transplanted)
	12 Months	All crops

\*Cotton: Refer to "Requirements for Planting Time Applications" and "Replanting Instructions" for additional instructions.



## Application Instructions for Willowood Clomazone 3ME Applied in Combination with Other Herbicides for Use on Soybeans

Tank mix Willowood Clomazone 3ME with other soybean herbicides to broaden the weed control spectrum compared to application of this product alone. The tables below provide a combination of products and application rates to broaden weed control spectrum. Willowood, LLC makes no claims as to weed control for any combination of rates or products other than as directed in this label. Updates and additional instructions for use of this product with other herbicides not listed in the label may become available after the date of publication for this label. Contact your local Willowood, LLC representative for more recent technical bulletins and supplemental labels. Observe all precautions and limitations appearing on the labels of each product when applying mixtures of Willowood Clomazone 3ME with other herbicides. The most restrictive precautions and instructions for all products must be followed. **DO NOT** incorporate tank mixtures of Willowood Clomazone 3ME and other products. When using tank mixes, observe all application precautions, rotational crop restrictions and replanting instructions of each product's label.

### Willowood Clomazone 3ME plus Canopy® Herbicide - Northern & Southern Regions

These products tank mixed together broaden the weed control spectrum.

**DO NOT** apply this product to soils with <1/2% organic matter.

Broadcast Rates per Acre*			Soil Texture
Willowood Clomazone 3ME		Canopy	
North Region	South Region		
2.0 - 2 2/3 pts.	2 2/3 pts. - 3 1/3 pts.	4 - 6 oz.	Coarse (light) Soils: Loamy sand, sandy loam
2.0 - 2 2/3 pts.	2 2/3 pts. - 3 1/3 pts.	5 - 8 oz.	Medium Soils: Loam, silt, silt loam, sandy clay, sandy clay loam
2.0 - 2 2/3 pts.	2 2/3 pts. - 3 1/3 pts.	6 - 8 oz.	Fine (heavy) Soils: Silty clay, clay loam, silty clay loam, clay

\*Select lower to higher rates within the range to correspond to the coarser to finer soil types within the soil texture group. The rate within the range should correspond to the anticipated weed pressure.

**Willowood Clomazone 3ME plus Canopy® Herbicide - Northern & Southern Regions (continued)****Broadleaf Weeds and Grasses Controlled**

<b>Grasses</b>	<b>Broadleaf Weeds</b>
Barnyardgrass	Balloon Vine
Bermudagrass	Black nightshade
Broadleaf signalgrass	Cocklebur**
Crabgrass (Large & Smooth)	Copperleaf
Cupgrass (Southwestern & Woolly)	Curly Dock
Field Sandbur	Dayflower
Foxtail (Giant, Green, Purple, Robust, Yellow)	Florida Beggarweed
Goosegrass	Florida Pusley (partial control)
Itchgrass	Hemp sesbania
Johnsongrass seedling	Hophornbeam (partial control)
Junglerice	Jimsonweed
Panicum (Common, Fall, & Texas)	Joint Vetch
Red Rice	Kochia
Shattercane	Lambsquarter
Wild Proso Millet	Morning Glory (Annual, Ivyleaf, Entireleaf, Pitted, Tall) (partial control)**
	Pigweed (Redroot, Smooth)
	Prickly Sida
	Prostrate Spurge
	Purslane
	Ragweed (Common)
	Redroot Pigweed
	Redvine
	Redweed
	Sicklepod (partial control)**
	Smartweed (Pennsylvania)
	Smooth Pigweed
	Spurred Anoda
	Tropic Croton
	Velvetleaf
	Venice Mallow
	Wild Poinsettia

\*\*Large seeded weeds, germinating deep in the soil, such as Morning Glory, sicklepod and cocklebur or weeds with subsequent flushes may require a cultivation or an application of a postemergence herbicide for season long control.

Follow all precautions and limitations appearing on the labels of each product being mixed. Observe precautions and restrictions sections regarding planting depth, phosphate pesticide interaction, previous pesticides, and sensitive soybean varieties on the Canopy label. Follow application precautions, rotational instructions, and replanting instructions of each product's label when using tank mixing Canopy with Willowood Clomazone 3ME.

## Willowood Clomazone 3ME plus Dual® 8 E Herbicide - Northern & Southern Regions

These products tank mixed together broaden the weed control spectrum.

Broadcast Rates per Acre			Soil Texture
Willowood Clomazone 3ME		Dual® 8 E	
North Region	South Region		
2.0 - 2 2/3 pts.	2 2/3 pts. - 3 1/3 pts.	1 1/2 - 2 pts.	Coarse (light) Soils: Loamy sand, sandy loam
2.0 - 2 2/3 pts.	2 2/3 pts. - 3 1/3 pts.	2 - 2 1/2 pts.	Medium Soils: Loam, silt, silt loam, sandy clay, sandy clay loam
2.0 - 2 2/3 pts.	2 2/3 pts. - 3 1/3 pts.	2 - 2 1/2 pts.	Fine (heavy) Soils: Silty clay, clay loam, silty clay loam, clay

**Willowood Clomazone 3ME plus Dual® 8 E Herbicide - Northern & Southern Regions (continued)**

<b>Broadleaf Weeds and Grasses Controlled</b>	
<b>Grasses</b>	<b>Broadleaf Weeds</b>
Barnyardgrass	Balloon Vine
Bermudagrass	Black nightshade
Broadleaf signalgrass	Carpetweed
Crabgrass (Large & Smooth)	Cocklebur
Cupgrass (Southwestern & Woolly)	Curly Dock
Field Sandbur	Dayflower
Foxtail (Giant, Green, Purple, Robust, Yellow)	Florida Beggarweed
Goosegrass	Florida Pusley (partial control)
Itchgrass	Jimsonweed
Johnsongrass seedling	Joint Vetch
Junglerice	Kochia
Nutsedge	Lambsquarter
Panicum (Common, Fall, & Texas)	Morning Glory (Pitted)
Red Rice (Use higher rates in areas with heavy weed pressure)	Pigweed
Shattercane	Prickly Sida
Wild Proso Millet	Purslane
Witchgrass	Ragweed (Common)
	Redroot Pigweed
	Redvine
	Redweed
	Smartweed (Pennsylvania)
	Smooth Pigweed
	Spurred Anoda
	Tropic Croton
	Velvetleaf
	Venice Mallow
	Wild Poinsettia

Follow all precautions and limitations appearing on the labels of each product being mixed. Follow application precautions, rotational instructions, and replanting instructions of each product's label when using tank mixing Dual® 8 E with Willowood Clomazone 3ME. Follow the most restrictive precautions and instructions from all products mixed.

## Willowood Clomazone 3ME plus Lasso® or Lasso® Micro-Tech - Northern & Southern Regions

These products tank mixed together broaden the weed control spectrum compared to either product used alone.

Broadcast Rates per Acre			Soil Texture
Willowood Clomazone 3ME		Lasso®/Lasso® Micro-Tech	
North Region	South Region		
2.0 - 2 2/3 pts.	2 2/3 pts. - 3 1/3 pts.	2 - 2 1/2 qts.	Coarse (light) Soils: Loamy sand, sandy loam
2.0 - 2 2/3 pts.	2 2/3 pts. - 3 1/3 pts.	2 1/2 - 3 qts.	Medium Soils: Loam, silt, silt loam, sandy clay, sandy clay loam
2.0 - 2 2/3 pts.	2 2/3 pts. - 3 1/3 pts.	2 1/2 - 3 qts.	Fine (heavy) Soils: Silty clay, clay loam, silty clay loam, clay

\*Lasso: Select lower to higher rates within the specified rates provided based on lighter to heavier soil types (i.e., sand, loamy sand, sandy loam) within a textural group.

**Willowood Clomazone 3ME plus Lasso® or Lasso® Micro-Tech - Northern & Southern Regions  
(continued)**

<b>Broadleaf Weeds and Grasses Controlled</b>	
<b>Grasses</b>	<b>Broadleaf Weeds</b>
Barnyardgrass	Black nightshade
Bermudagrass	Carpetweed
Broadleaf signalgrass	Cocklebur
Crabgrass (Large & Smooth)	Curly Dock
Cupgrass (Southwestern & Woolly)	Dayflower
Field Sandbur	Florida Beggarweed
Foxtail (Giant, Green, Purple, Robust, Yellow)	Florida Pusley (partial control)
Goosegrass	Jimsonweed
Itchgrass	Joint Vetch
Johnsongrass seedling	Kochia
Junglerice	Lambsquarter
Panicum (Common, Fall, & Texas)	Morning Glory (Pitted)
Shattercane	Pigweed (Redroot, Smooth)
Wild Proso Millet	Prickly Sida
	Prostrate Spurge
	Purslane
	Ragweed (Common)
	Redroot Pigweed
	Redvine
	Redweed
	Smartweed (Pennsylvania)
	Spurred Anoda
	Tropic Croton
	Velvetleaf
	Venice Mallow
	Wild Poinsettia

Follow all precautions and limitations appearing on the labels of each product being mixed. Follow application precautions, rotational instructions, and replanting instructions of each product's label when using tank mixing Lasso® or Lasso® Tech. with Willowood Clomazone 3ME. Follow the most restrictive precautions and instructions from all products mixed.

### Willowood Clomazone 3ME plus Lorox® DF – Northern & Southern Regions

These products tank mixed together and applied preemergence will broaden the weed control spectrum compared to either product used alone. However, do not incorporate this tank mix combination.

Plant seed a minimum of 1 3/4 inches deep to avoid Lorox injury to soybeans. See Lorox label for all other planting and application precautions. One pint of Lorox L equals one pound of Lorox DF; be sure to use equivalent rates.

Do not use this combination on sand, loamy sand, gravelly soils, or exposed subsoils or soils with <1% organic matter. Use only on loamy sand with >1% organic matter.

Broadcast Rates per Acre*			Soil Texture
Willowood Clomazone 3ME		Lorox® DF	
North Region	South Region		
2.0 - 2 2/3 pts.	2 2/3 pts. - 3 1/3 pts.	2/3 - 1 lb.	Coarse (light) Soils: Loamy sand, sandy loam
2.0 - 2 2/3 pts.	2 2/3 pts. - 3 1/3 pts.	1 - 1 1/2 lbs.	Medium Soils: Loam, silt, silt loam, sandy clay, sandy clay loam
2.0 - 2 2/3 pts.	2 2/3 pts. - 3 1/3 pts.	1 1/4 - 2 lbs.	Fine (heavy) Soils: Silty clay, clay loam, silty clay loam, clay

\*Select lower to higher rates within the specified rates based on lighter to heavier soil types (i.e., loam, silt loam, sandy clay, sandy clay loam) within a textural group.

**Willowood Clomazone 3ME plus Lorox® DF – Northern & Southern Regions (continued)**

<b>Broadleaf Weeds and Grasses Controlled</b>	
<b>Grasses</b>	<b>Broadleaf Weeds</b>
Barnyardgrass	Balloon Vine
Bermudagrass	Black Nightshade
Broadleaf signalgrass	Carpetweed
Crabgrass (Large & Smooth)	Chickweed
Cupgrass (Southwestern & Woolly)	Cocklebur
Field Sandbur	Curly Dock
Foxtail (Giant, Green, Purple, Robust, Yellow)	Dayflower
Goosegrass	Florida Beggarweed
Itchgrass	Florida Pusley (partial control)
Johnsongrass seedling	Jimsonweed
Junglerice	Joint Vetch
Panicum (Common, Fall, & Texas)	Kochia
Red Rice	Lambsquarter
Shattercane	Morning Glory (Pitted)
Wild Proso Millet	Pigweed (Redroot, Smooth)
	Prickly Sida
	Prostrate Spurge
	Purslane
	Ragweed (Common)
	Redvine
	Redweed
	Smartweed (Pennsylvania)
	Spurred Anoda
	Tropic Croton
	Velvetleaf
	Venice Mallow
	Wild Mustard
	Wild Poinsettia
	Wild Radish

Follow all precautions and limitations appearing on the labels of each product being mixed. Follow application precautions, rotational instructions, and replanting instructions of each product's label when using tank mixing Lorox® DF with Willowood Clomazone 3ME. Follow the most restrictive precautions and instructions from all products mixed.



### Willowood Clomazone 3ME plus Lorox Plus® – Northern & Southern Regions

These products tank mixed together and applied preemergence will broaden the weed control spectrum compared to either product used alone. However, **do not** incorporate this tank mix combination.

Plant seed a minimum of 1 3/4 inches deep to avoid Lorox injury to soybeans. See Lorox label for all other planting and application precautions.

**DO NOT** use this combination on sand, loamy sand, gravelly soils, or exposed subsoils or soils with <1% organic matter. Use only on loamy sand with >1% organic matter.

Broadcast Rates per Acre*			Soil Texture
Willowood Clomazone 3ME		Lorox® Plus	
North Region	South Region		
2.0 - 2 2/3 pts.	2 2/3 pts. - 3 1/3 pts.	9 - 12 oz.	Coarse (light) Soils: Sand, loamy sand, sandy loam
2.0 - 2 2/3 pts.	2 2/3 pts. - 3 1/3 pts.	10 - 14 oz.	Medium Soils: Loam, silt, silt loam, sandy clay, sandy clay loam
2.0 - 2 2/3 pts.	2 2/3 pts. - 3 1/3 pts.	11 - 14 oz.	Fine (heavy) Soils: Silty clay, clay loam, silty clay loam, clay

\*Select lower to higher rates within the specified rates provided based on lighter to heavier soil types (i.e., loam, silt loam, sandy clay, sandy clay loam) within a textural group.

**Willowood Clomazone 3ME plus Lorox Plus® – Northern & Southern Regions (continued)**

<b>Broadleaf Weeds and Grasses Controlled</b>	
<b>Grasses</b>	<b>Broadleaf Weeds</b>
Barnyardgrass	Balloon Vine
Bermudagrass	Black nightshade (partial control)
Broadleaf signalgrass	Carpetweed
Crabgrass (Large & Smooth)	Cocklebur**
Cupgrass (Southwestern & Woolly)	Curly Dock
Field Sandbur	Dayflower
Foxtail (Giant, Green, Purple, Robust, Yellow)	Florida Beggarweed
Goosegrass	Florida Pusley (partial control)
Itchgrass	Jimsonweed
Johnsongrass seedling	Joint Vetch
Junglerice	Kochia
Panicum (Common, Fall, & Texas)	Lambsquarter
Red Rice	Morning Glory (Annual, Ivyleaf, Entireleaf, Pitted, Tall) (partial control)**
Shattercane	Mustards
Wild Proso Millet	Pigweed (Redroot, Smooth)
	Prickly Sida
	Prostrate Spurge
	Purslane
	Ragweed (Common)
	Ragweed (Giant) (Partial Control)
	Redvine
	Smartweed (Pennsylvania)
	Spurred Anoda
	Tropic Croton
	Velvetleaf
	Venice Mallow
	Wild Poinsettia

\*\*Loamy sand: Large seeded weeds, germinating deep in the soil, such as Morning Glory, sicklepod and cocklebur or weeds with subsequent flushes may require a cultivation or an application of a postemergence herbicide for season long control.

Follow all precautions and limitations appearing on the labels of each product being mixed. Follow application precautions, rotational instructions, and replanting instructions of each product's label when using tank mixing Lorox Plus® with Willowood Clomazone 3ME. Follow the most restrictive precautions and instructions from all products mixed.

**Willowood Clomazone 3ME plus Metribuzin (Sencor® or Lexone®) - Northern Region Only**

These products tank mixed together and applied preemergence will broaden the weed control spectrum compared to either product used alone.

**DO NOT** use this combination on soils with <0.5% organic matter.

**DO NOT** use on sand or loamy sand soils with <2.0% organic matter.

**DO NOT** use this combination on soils with a calcareous surface area or a pH >7.5 or crop injury can occur.

Metribuzin can injure soybeans in excessively moist conditions, cool weather, or in compacted soils.

<b>Broadcast Rates per Acre*</b>		<b>Soil Texture</b>
<b>Willowood Clomazone 3ME</b>	<b>Sencor® DF or Lexone® DF**</b>	
2.0 - 2 2/3 pts.	1/4 - 1/3 lb.	Coarse (light) Soils: Loamy sand, sandy loam
2.0 - 2 2/3 pts.	1/4 - 1/2 lb.	Medium Soils: Loam, silt, silt loam, sandy clay, sandy clay loam
2.0 - 2 2/3 pts.	1/3 - 1/2 lb.	Fine (heavy) Soils: Silty clay, clay loam, silty clay loam, clay

\*Select lower to higher rates within the specified rates provided based on lighter to heavier soil types (i.e., silt loam, sandy clay, sandy clay loam) within a textural group.

\*\*For pints Sencor 4 or Lexone 4L, multiply lbs. DF by 1.5

**Willowood Clomazone 3ME plus Metribuzin (Sencor® or Lexone®) - Northern Region Only  
(continued)**

<b>Broadleaf Weeds and Grasses Controlled</b>	
<b>Grasses</b>	<b>Broadleaf Weeds</b>
Barnyardgrass	Balloon Vine
Bermudagrass	Black nightshade
Broadleaf signalgrass	Carpetweed
Crabgrass (Large & Smooth)	Cocklebur
Cupgrass (Southwestern & Woolly)	Curly Dock
Field Sandbur	Dayflower
Foxtail (Giant, Green, Purple, Robust, Yellow)	Florida Beggarweed
Goosegrass	Florida Pusley (partial control)
Itchgrass	Jimsonweed
Johnsongrass seedling	Joint Vetch
Junglerice	Kochia
Shattercane	Lambsquarter
Wild Proso Millet	Morning Glory (Pitted)
	Prickly Sida
	Prostrate Spurge
	Purslane
	Ragweed (Common)
	Redroot Pigweed
	Redvine
	Redweed
	Smartweed (Pennsylvania)
	Smooth Pigweed
	Spurred Anoda
	Tropic Croton
	Velvetleaf
	Venice Mallow
	Wild Mustard
	Wild Poinsettia

Follow all precautions and limitations appearing on the labels of each product being mixed. Follow application precautions, rotational instructions, and replanting instructions of each product's label when using tank mixing Sencor® or Lexone® with Willowood Clomazone 3ME. Follow the most restrictive precautions and instructions from all products mixed.

### **Willowood Clomazone 3ME plus Metribuzin (Sencor® or Lexone®) - Southern Region Only**

These products tank mixed together and applied preemergence will broaden the weed control spectrum compared to either product used alone.

**DO NOT** use this combination on soils with <0.5% organic matter.

**DO NOT** use on sand or loamy sand soils with <2.0% organic matter.

**DO NOT** use this combination on soils with a calcareous surface area or a pH >7.5 or crop injury can occur.

Use 2/3 - 1 lb. metribuzin on silty clay and heavy clay soil of the Mississippi Delta.

Metribuzin can injure soybeans in excessively moist conditions, cool weather, or in compacted soils.

<b>Broadcast Rates per Acre*</b>		<b>Soil Texture</b>
<b>Willowood Clomazone 3ME</b>	<b>Sencor DF® or Lexone DF®**</b>	
2 2/3 pts. - 3 1/3 pts.	1/4 - 1/3 lb.	Coarse (light) Soils: Loamy sand, sandy loam
2 2/3 pts. - 3 1/3 pts.	1/4 - 1/2 lb.	Medium Soils: Loam, silt, silt loam, sandy clay, sandy clay loam
2 2/3 pts. - 3 1/3 pts.	1/3 - 2/3 lb.	Fine (heavy) Soils: Silty clay, clay loam, silty clay loam, clay

\*Select lower to higher rates within the specified rates provided based on lighter to heavier soil types (i.e., silt loam, sandy clay, sandy clay loam) within a textural group.

\*\*For pints Sencor 4 or Lexone 4L, multiply lbs. DF by 1.5

**Willowood Clomazone 3ME plus Metribuzin (Sencor® or Lexone®) – Southern Region Only  
(continued)**

<b>Broadleaf Weeds and Grasses Controlled</b>	
<b>Grasses</b>	<b>Broadleaf Weeds</b>
Barnyardgrass	Balloon Vine
Bermudagrass	Black nightshade
Broadleaf signalgrass	Carpetweed
Crabgrass (Large & Smooth)	Cocklebur
Cupgrass (Southwestern & Woolly)	Curly Dock
Field Sandbur	Dayflower
Foxtail (Giant, Green, Purple, Robust, Yellow)	Florida Beggarweed
Goosegrass	Florida Pusley (partial control)
Itchgrass	Giant Ragweed (partial control)
Johnsongrass seedling	Jimsonweed
Junglerice	Joint Vetch
Panicum (Common, Fall, & Texas)	Kochia
Red Rice	Lambsquarter
Shattercane	Morning Glory (Pitted)
Wild Proso Millet	Prickly Sida
	Prostrate Spurge
	Purslane
	Ragweed (Common)
	Redroot Pigweed
	Redvine
	Redweed
	Sicklepod (partial control & reduced competition)
	Smartweed (Pennsylvania)
	Smooth Pigweed
	Spurred Anoda
	Tropic Croton
	Velvetleaf
	Venice Mallow
	Wild Mustard
	Wild Poinsettia

Follow all precautions and limitations appearing on the labels of each product being mixed. Follow application precautions, rotational instructions, and replanting instructions of each product's label when using tank mixing Sencor® or Lexone® with Willowood Clomazone 3ME. Follow the most restrictive precautions and instructions from all products mixed.

## Willowood Clomazone 3ME plus Prowl® 4 EC Herbicide - Northern & Southern Regions

These products tank mixed together and applied preemergence will broaden the weed control spectrum compared to either product used alone.

Broadcast Rates per Acre*			Soil Texture
Willowood Clomazone 3ME		Prowl 4EC®	
North Region	South Region		
2.0 - 2 2/3 pts.	2 2/3 - 3 1/3 pts.	1 - 1 1/2 pts.	Coarse (light) Soils: Loamy sand, sandy loam
2.0 - 2 2/3 pts.	2 2/3 - 3 1/3 pts.	2 pts.	Medium Soils: Loam, silt, silt loam, sandy clay, sandy clay loam
2.0 - 2 2/3 pts.	2 2/3 - 3 1/3 pts.	2 - 2.5 pts.	Fine (heavy) Soils: Silty clay, clay loam, silty clay loam, clay

\*Select lower to higher rates within the specified rates provided based on lighter to heavier soil types (i.e., loam, silt loam, sandy clay, sandy clay loam) within a textural group.

**Willowood Clomazone 3ME plus Prowl® 4 EC Herbicide – Northern & Southern Regions (continued)**

<b>Broadleaf Weeds and Grasses Controlled</b>	
<b>Grasses</b>	<b>Broadleaf Weeds</b>
Barnyardgrass	Balloon Vine
Bermudagrass	Black nightshade
Broadleaf signalgrass	Carpetweed
Crabgrass (Large & Smooth)	Cocklebur
Cupgrass (Southwestern & Woolly)	Curly Dock
Field Sandbur	Dayflower
Foxtail (Giant, Green, Purple, Robust, Yellow)	Florida Beggarweed
Goosegrass	Florida Pusley Giant Ragweed (partial control)
Itchgrass	Hemp sesbania (Coffeeweed)
Johnsongrass seedling	Hophornbeam (partial control)
Junglerice	Jimsonweed
Panicum (Common, Fall, & Texas)	Joint Vetch
Red Rice	Kochia
Shattercane	Lambsquarter
Wild Proso Millet	Morning Glory (Pitted)
	Pigweed (Redroot, Smooth)
	Prickly Sida
	Prostrate Spurge
	Purslane
	Ragweed (Common)
	Redroot Pigweed
	Redvine
	Redweed
	Sicklepod (partial control & reduced competition)
	Smartweed (Pennsylvania)
	Smooth Pigweed
	Spurred Anoda
	Tropic Croton
	Velvetleaf
	Venice Mallow
	Wild Poinsettia

Follow all precautions and limitations appearing on the labels of each product being mixed. Follow application precautions, rotational instructions, and replanting instructions of each product's label when using tank mixing Prowl 4EC® with Willowood Clomazone 3ME. Follow the most restrictive precautions and instructions from all products mixed.



## Willowood Clomazone 3ME Plus Postemergence Applications of Pursuit® Herbicide

These products tank mixed together and applied postemergence will broaden the weed control spectrum compared to either product used alone.

## Willowood Clomazone 3ME Followed by Postemergence Applications of Pursuit® Herbicide++

Broadcast Rates per Acre		Soil Texture*
Willowood Clomazone 3ME	Pursuit	
2.0 - 2 2/3 pts. (0.25 lb. AI)	2 - 3 fl. oz.	Coarse, Medium, and Fine Soils
*Select lower to higher rates within the specified rates provided based on lighter to heavier soil types within a textural group.		

++ Postemergence application requires specific additives. Refer to current Pursuit label for specific directions.

Application of Willowood Clomazone 3ME will reduce competition from the above weeds, allowing control of light to moderate infestations with a postemergence application of Pursuit herbicide at the noted rate.

Apply to weeds less than one inch tall.

Application of Willowood Clomazone 3ME followed by a postemergence treatment with Pursuit will control all of the weeds controlled by use of Willowood Clomazone 3ME alone PLUS:

Cocklebur

Pigweed

Nightshade (Black, Eastern Black, Hairy)

Mustard Species

Pennsylvania Smartweed

The maximum weed height to control weeds is 1 inch.

Weed control may be erratic under heavy weed infestations. Improve control by cultivating approximately 14 days after the postemergence treatment.

**DO NOT** apply Pursuit herbicide when wind velocity is greater than 10 mph, or when spray may be carried to sensitive crops, including leafy vegetables and sugar beets.

Follow all precautions and limitations appearing on the labels of each product being mixed. Follow application precautions, rotational instructions, and replanting instructions of each product's label when using tank mixing Pursuit with Willowood Clomazone 3ME. Follow the most restrictive precautions and instructions from all products mixed.

### Willowood Clomazone 3ME Plus Scepter® Herbicide - Northern Region Only

These products tank mixed together and applied postemergence will broaden the weed control spectrum compared to either product used alone.

See Scepter label for specific geographic and rainfall restrictions.

Broadcast Rates per Acre		Soil Texture*
Willowood Clomazone 3ME	Scepter®	
2.0 - 2 2/3 pts.	1/3 - 1/2 pt.**	All soil types

\*Use higher rates within the specified range when weed pressure is heavy.

\*\*Rates are based on Scepter® products containing 1.5 lbs. AI Equivalent amounts of Scepter® O.T. containing 2 1/2 lbs. AI are 3.2 to 4.8 fl. oz. (1/5 to 1/3 pint).

SPECIMEN

**Willowood Clomazone 3ME Plus Scepter® Herbicide – Northern Region Only (continued)**

<b>Broadleaf Weeds and Grasses Controlled</b>	
<b>Grasses</b>	<b>Broadleaf Weeds</b>
Barnyardgrass	Balloon Vine
Bermudagrass	Black nightshade (partial control)
Broadleaf signalgrass	Cocklebur
Crabgrass (Large & Smooth)	Common Sunflower
Cupgrass (Southwestern & Woolly)	Curly Dock
Field Sandbur	Dayflower
Foxtail (Giant, Green, Purple, Robust, Yellow)	Florida Beggarweed
Goosegrass	Florida Pusley (partial control)
Itchgrass	Giant Ragweed (partial control)
Johnsongrass seedling	Jimsonweed
Junglerice	Joint Vetch
Panicum (Common, Fall, & Texas)	Kochia
Red Rice	Lambsquarter
Shattercane	Morning Glory (Pitted)
Wild Proso Millet	Pigweed spp.
	Prickly Sida
	Prostrate Spurge
	Purslane
	Ragweed (Common)
	Redroot Pigweed
	Redvine
	Redweed
	Sicklepod (partial control & reduced competition)
	Smartweed (Pennsylvania)
	Smooth Pigweed
	Spurred Anoda
	Tropic Croton
	Velvetleaf
	Venice Mallow
	Wild Poinsettia

Soybeans can be replanted if crop yield is reduced by weather conditions, but do not make additional tank mix applications with Willowood Clomazone 3ME and Scepter®.

Important: Adequate moisture in the soil is required for maximum effectiveness of this tank mix.

Follow all precautions and limitations appearing on the labels of each product being mixed. Follow application precautions, rotational instructions, and replanting instructions of each product's label when using tank mixing Scepter® with Willowood Clomazone 3ME. Follow the most restrictive precautions and instructions from all products mixed.

## Willowood Clomazone 3ME Plus Scepter® Herbicide - Southern Region Only

See Scepter label for specific geographic and rainfall restrictions.

Broadcast Rates per Acre		Soil Texture*
Willowood Clomazone 3ME	Scepter®	
2 2/3 - 3 1/3 pts.	1/2 - 2/3 pt.**	All soil types

\*Use higher rates within the specified range when weed pressure is heavy.  
\*\*Rates are based on Scepter® products containing 1.5 lbs. AI Equivalent amounts of Scepter® O.T. containing 2 1/2 lbs. AI are 4.8 to 7.2 fl. oz.

SPECIMEN

**Willowood Clomazone 3ME Plus Scepter® Herbicide – Southern Region Only (continued)**

<b>Broadleaf Weeds and Grasses Controlled</b>	
<b>Grasses</b>	<b>Broadleaf Weeds</b>
Barnyardgrass	Balloon Vine
Bermudagrass	Black nightshade (partial control)
Broadleaf signal grass	Carpetweed
Crabgrass (Large & Smooth)	Chickweed
Cupgrass (Southwestern & Woolly)	Cocklebur
Field Sandbur	Common Sunflower
Foxtail (Giant, Green, Purple, Robust, Yellow)	Copperleaf
Goosegrass	Curly Dock
Itchgrass	Dayflower
Johnsongrass seedling	Florida Beggarweed
Junglerice	Florida Pusley (partial control)
Millet	Giant Ragweed (partial control)
Panicum (Common, Fall, & Texas)	Hemp sesbania (Coffeeweed)
Shattercane	Hophornbeam (partial control)
Wild Proso	Jimsonweed
	Joint
	Kochia
	Lambsquarter
	Morning Glory (Annual, Ivyleaf, Entireleaf, Pitted, Tall) (partial control)
	Pigweed spp.
	Prickly Sida
	Purslane
	Ragweed (Common)
	Redroot Pigweed
	Redvine
	Redweed
	Sicklepod (partial control & reduced competition)
	Smartweed (Pennsylvania)
	Smooth Pigweed
	Spurred Anoda
	Tropic Croton
	Velvetleaf
	Venice Mallow
	Vetch
	Wild Mustard
	Wild Poinsettia

Soybeans can be replanted if crop yield is reduced by weather conditions, but do not make additional tank mix applications with Willowood Clomazone 3ME and Scepter®.

**Important:** Adequate moisture in the soil is required for maximum effectiveness of this tank mix.

Follow all precautions and limitations appearing on the labels of each product being mixed. Follow application precautions, rotational instructions, and replanting instructions of each product's label when using tank mixing Scepter® with Willowood Clomazone 3ME. Follow the most restrictive precautions and instructions from all products mixed.

### Willowood Clomazone 3ME Plus Sonalan® EC Herbicide - Northern Region Only

These products tank mixed together broaden the weed control spectrum compared to either product used alone.

Broadcast Rates per Acre*		Soil Texture
Willowood Clomazone 3ME	Sonalan® EC	
2.0 pts. - 2 2/3 pts.	1 1/4 - 2 pts.	Coarse (light) Soils: Loamy sand, sandy loam
2.0 pts. - 2 2/3 pts.	1 3/4 - 2 1/2 pts.	Medium Soils: Loam, silt, silt loam, sandy clay, sandy clay loam
2.0 pts. - 2 2/3 pts.	2 1/4 - 3 pts.	Fine (heavy) Soils: Silty clay, clay loam, silty clay loam, clay

\*Use lower to higher rates within the specified range to correspond to the coarser to finer soil types within the textural group. Rates within the range should correspond to weed pressure.

**Willowood Clomazone 3ME Plus Sonalan® EC Herbicide – Northern Region Only (continued)****Broadleaf Weeds and Grasses Controlled**

<b>Grasses</b>	<b>Broadleaf Weeds</b>
Barnyardgrass	Balloon Vine
Bermudagrass	Black nightshade (partial control)
Broadleaf signalgrass	Carpetweed
Crabgrass (Large & Smooth)	Chickweed
Cupgrass (Southwestern & Woolly)	Cocklebur
Field Sandbur	Curly Dock
Foxtail (Giant, Green, Purple, Robust, Yellow)	Dayflower
Goosegrass	Florida Beggarweed
Itchgrass	Florida Pusley (partial control)
Johnsongrass seedling	Hophornbeam (partial control)
Junglerice	Jimsonweed
Panicum (Common, Fall, & Texas)	Joint Vetch
Red Rice	Kochia
Shattercane	Lambsquarter
Wild Proso Millet	Morning Glory (Annual, Ivyleaf, Entireleaf, Pitted, Tall) (partial control)
	Pigweed spp.
	Prickly Sida
	Prostrate Spurge
	Purslane
	Ragweed (Common)
	Redroot Pigweed
	Redvine
	Redweed
	Smartweed (Pennsylvania)
	Smooth Pigweed
	Spurred Anoda
	Tropic Croton
	Velvetleaf
	Venice Mallow
	Wild Poinsettia

Follow all precautions and limitations appearing on the labels of each product being mixed. Follow application precautions, rotational instructions, and replanting instructions of each product's label when using tank mixing Sonalan® EC with Willowood Clomazone 3ME. Follow the most restrictive precautions and instructions from all products mixed.

## Willowood Clomazone 3ME Plus Squadron - Northern Region Only

These products tank mixed together broaden the weed control spectrum compared to either product used alone.

See Squadron label for specific geographic and rainfall restrictions.

Broadcast Rates per Acre		Soil Texture
Willowood Clomazone 3ME	Squadron®*	
2.0 - 2 2/3 pts.	1 1/2 - 2 1/4 pts.**	All soil types
*Maximum application rate in Northern Area on the Squadron label is 1 1/2 pts.		
**Use the higher rate of Squadron in areas where cocklebur infestation is heavy.		

Broadleaf Weeds and Grasses Controlled	
Grasses	Broadleaf Weeds
Barnyardgrass	Balloon Vine
Bermudagrass	Black nightshade
Broadleaf signalgrass	Cocklebur
Crabgrass (Large & Smooth)	Common Sunflower
Cupgrass (Southwestern & Woolly)	Curly Dock
Field Sandbur	Dayflower
Foxtail (Giant, Green, Purple, Robust, Yellow)	Florida Beggarweed
Goosegrass	Florida Pusley (partial control)
Itchgrass	Jimsonweed
Johnsongrass seedling	Joint Vetch
Junglerice	Kochia
Panicum (Common, Fall, & Texas)	Lambsquarter
Red Rice	Morning Glory (Pitted)
Shattercane	Pigweed spp.
Wild Proso Millet	Prickly Sida
	Prostrate Spurge
	Purslane
	Ragweed (Common)
	Redroot Pigweed
	Redvine
	Redweed
	Smartweed (Pennsylvania)
	Smooth Pigweed
	Spurred Anoda
	Tropic Croton
	Velvetleaf
	Venice Mallow

Follow all precautions and limitations appearing on the labels of each product being mixed. Follow application precautions, rotational instructions, and replanting instructions of each product's label when using tank mixing Squadron with Willowood Clomazone 3ME. Follow the most restrictive precautions and instructions from all products mixed.



### **Willowood Clomazone 3ME Plus Squadron - Southern Region Only**

These products tank mixed together broaden the weed control spectrum compared to either product used alone.

See Squadron label for specific geographic restrictions.

<b>Broadcast Rates per Acre</b>		<b>Soil Texture</b>
<b>Willowood Clomazone 3ME</b>	<b>Squadron®*</b>	
2 2/3 pts.	2 1/4 - 3 pts.*	All soil types
*Use the higher rate of Squadron in areas where cocklebur and pigweed infestations are heavy.		

SPECIMEN

**Willowood Clomazone 3ME Plus Squadron – Southern Region Only (continued)**

<b>Broadleaf Weeds and Grasses Controlled</b>	
<b>Grasses</b>	<b>Broadleaf Weeds</b>
Barnyardgrass	Balloon Vine
Bermudagrass	Black nightshade
Broadleaf signalgrass	Cocklebur
Crabgrass (Large & Smooth)	Common Sunflower
Cupgrass (Southwestern & Woolly)	Copperleaf
Field Sandbur	Curly Dock
Foxtail (Giant, Green, Purple, Robust, Yellow)	Dayflower
Goosegrass	Florida Beggarweed
Itchgrass	Florida Pusley (partial control)
Johnsongrass seedling	Jimsonweed
Junglerice	Joint Vetch
Panicum (Common, Fall, & Texas)	Kochia
Red Rice	Lambsquarter
Shattercane	Morning Glory (Pitted)
Wild Proso Millet	Pigweed spp.
	Prickly Sida
	Prostrate Spurge
	Purslane
	Ragweed (Common)
	Redroot Pigweed
	Redvine
	Redweed
	Smartweed (Pennsylvania)
	Smooth Pigweed
	Spurred Anoda
	Tropic Croton
	Velvetleaf
	Venice Mallow
	Wild Poinsettia

Follow all precautions and limitations appearing on the labels of each product being mixed. Follow application precautions, rotational instructions, and replanting instructions of each product's label when using tank mixing Squadron with Willowood Clomazone 3ME. Follow the most restrictive precautions and instructions from all products mixed.

### **Willowood Clomazone 3ME Application Followed by Postemergence Treatment**

Willowood Clomazone 3ME can be applied alone on soybeans or in a tank mix with other postemergence herbicides approved for use on soybeans to control of emerged weeds as listed on the postemergence herbicide product label. Follow all precautions and limitations appearing on the labels of each product being mixed. Follow application precautions, rotational instructions, and replanting instructions of each product's label. Follow the most restrictive precautions and instructions from all products mixed.

**DO NOT** apply Classic, Pursuit or Scepter as a postemergence treatment after Willowood Clomazone 3ME has been tank mixed with Canopy, Preview, Pursuit Plus, Scepter, or Squadron, unless their respective product label specifically provides instructions for making such a treatment. Using these products cumulatively can cause injury to soybeans or rotation crops because all of these herbicides share a common mode of action.

### **Willowood Clomazone 3ME Treatment Followed By Treatment with Acifluorfen (Ultra Blazer®) Herbicide**

A postemergence application of acifluorfen after treatment with Willowood Clomazone 3ME alone will control: Black Nightshade, Cocklebur, Hemp sesbania, Pigweed (Redroot), Pitted Morning Glory, Purple Moonflower, and Wild Mustard.

Apply 1 pint per acre of acifluorfen with .25% crop oil concentrate to actively growing weeds that have reached the 4-leaf growth stage (count **only** the fully developed true leaves). This timing correlates to soybean growth stages of the first to third trifoliate leaves.

**NOTE:** Willowood Clomazone 3ME retards weed development, thereby extending the period that acifluorfen can be used effectively. Read and follow all precautions, restrictions, and warnings on all product labels before making any application.

### **Willowood Clomazone 3ME Treatment Followed By Treatment with Scepter® Herbicide**

A postemergence application of Scepter after treatment with Willowood Clomazone 3ME alone will control: Cocklebur, Common Pigweed, and Wild Poinsettia.

Apply Scepter after crop emerges and before weeds reach 12" tall (6" for Wild Poinsettia). Make a broadcast application of 1/3 pint per acre when weeds are actively growing.

**DO NOT** apply Scepter postemergence when soybeans and weeds are stressed due to extremes in temperature and moisture.

**DO NOT** exceed 1/3 pint per acre per season of Scepter in the Northern region.

**DO NOT** exceed 2/3 pint per acre per season in the Southern region.

Squadron and Scepter contain the same active ingredient.

**Add a nonionic surfactant or crop oil concentrate to all postemergence applications. The nonionic surfactant must contain at least 80% active ingredient. Apply 2 pints per 100 gallons of spray mixture. Apply the crop oil concentrate (COC) at the rate specified on the COC label.**

Read and follow all precautions, restrictions, and warnings on all product labels before making any application.

# SUGARCANE

Willowood Clomazone 3ME is an effective preemergent soil applied treatment for controlling annual grasses and broadleaf weeds in sugarcane.

- Apply 2 2/3 to 3 1/3 pints (1.0 to 1.25 lbs. AI) per acre in 10 to 40 gallons of water per acre on newly planted or ratoon sugarcane using ground equipment.
- Apply to newly planted sugarcane in late summer or early fall.
- Apply to newly planted or ratoon sugarcane in spring through layby.

## Sugarcane Precautions

- Off-site movement of Willowood Clomazone 3ME by spray drift or vapors can cause foliar whitening or plant yellowing. Follow all precautions and instructions in the APPLICATION PRECAUTIONS, SPRAY DRIFT PRECAUTIONS and SPRAY DRIFT MANAGEMENT sections of this label before applying this product.

## Sugarcane Restrictions

- **DO NOT** apply Willowood Clomazone 3ME within 1,200 feet of: Towns and Housing Developments, Commercial Vegetable Production (except peppers, pumpkins, succulent peas, sweet corn, sweet potato, and winter squash), Commercial Fruit/Nut production, Commercial Nurseries, and Commercial Greenhouses.
- **DO NOT** apply when air currents can carry spray particles to areas where sensitive crops and plants are growing.
- **DO NOT** apply when temperature inversions exist.
- Prior to application adjacent properties must be checked. **Do not** apply within 300 feet of desirable plants.

## Preemergent Banded Treatments

Refer to the directions for banded application in the GROUND APPLICATION INSTRUCTIONS section.

## Replanting Instructions

- If the initial planting of newly planted sugarcane does not produce a uniform stand, replant sugarcane in fields treated with Willowood Clomazone 3ME alone.
- If tank mixing with an approved labeled product, refer to the replant instructions for that product.
- If a tank mix is used, refer to the product's labels for any additional replant instructions, and follow the most restrictive precautions and use directions. If replanting is required, follow the directions under REQUIREMENTS FOR PLANTING TIME APPLICATIONS section.

## Replanting Restrictions

- **DO NOT** make a second application of Willowood Clomazone 3ME if fields have been retreated.
- **DO NOT** replant treated fields with any crop at intervals that are not consistent with the ROTATIONAL CROP RESTRICTIONS on this label.

### Willowood Clomazone 3ME Rates and Weeds Controlled

Rate	Grasses Controlled
2 2/3 pts. (1.0 lb. AI)	Barnyardgrass, Broadleaf signalgrass, Large & Smooth Crabgrass, Southwestern Cupgrass, Woolly Cupgrass, Field Sandbur, Giant & Green Foxtail, Goosegrass, Itchgrass*, Panicum (Common, Fall, & Texas), Johnsongrass seedling
3 1/3 pts. (1.25 lbs. AI)	All grasses listed above PLUS: Bermudagrass**, Foxtail (Purple, Robust, Yellow), Junglerice (partial control), Shattercane (partial control), Wild Proso Millet (partial control)

\*Itchgrass: Do not incorporate Willowood Clomazone 3ME. Use 3 1/3 pints for high weed pressure and heavy soil.

\*\*Bermudagrass: Willowood Clomazone 3ME is effective in delaying Bermudagrass reestablishment when applied immediately after planting sugarcane in late summer or early fall.

### Willowood Clomazone 3ME Rates and Broadleaf Weeds Controlled

Rate	Broadleaf Weeds Controlled
2 2/3 pts. (1.0 lb. AI)	Velvetleaf, Spurred Anoda, Cocklebur, Common Ragweed, Dayflower, Florida Beggarweed, Jimsonweed, Kochia (partial control), Lambsquarter, Pennsylvania Smartweed (partial control), Prickly Sida, Purslane, Redvine (partial control), Redweed, Tropic Croton, Venice Mallow, Wild Poinsettia
3 1/3 pts. (1.25 lbs. AI)	All broadleaf weeds listed above PLUS: Balloon Vine (partial control), Black nightshade (partial control), Curly Dock, Joint Vetch (partial control), Kochia, Pitted Morning Glory, Pennsylvania Smartweed, Redroot Pigweed (partial control), Smooth Pigweed (partial control)

Willowood Clomazone 3ME can be applied in combination with other herbicides registered for use in sugarcane. If using a tank mixture, follow application precautions, rotational instructions, and replanting instructions of each product's label. Follow the most restrictive precautions and instructions from all products mixed.

### Crop Rotation Restrictions

- **DO NOT** apply more than 1.25 pounds active ingredient (3 1/3 pints) per acre during one growing season.
- **DO NOT** apply within 120 days prior to harvest on sugarcane for milling.
- **DO NOT** apply aerially or through irrigation equipment.
- **DO NOT** allow livestock to graze on treated sugarcane forage or trash, or feed treated sugarcane forage or trash to livestock.

Rate	Interval	Crop
2 2/3 pts. (1.0 lb. AI)	Anytime	Cotton*, Peppers, Soybeans, Sugarcane, Sweet Potato, Tobacco, Tuberous & Corm Vegetables
	9 Months	All crops listed above PLUS: Cabbage, Field Corn, Popcorn, Seed Corn, Sweet Corn, Cotton, Cucurbits, Dry Beans, Peas, Peanuts, Potatoes, Pumpkins (processing), Rice, Snap Beans, Sorghum, Squash, Sugar Beets, Tomatoes (Transplanted)
	12 Months	All crops
3 1/3 pts. (1.25 lbs. AI)	Anytime	Cotton*, Soybeans, Sugarcane, Sweet Potato, Tobacco, Tuberous & Corm Vegetables
	9 Months	All crops listed above PLUS: Cabbage, Field Corn, Popcorn, Seed Corn, Sweet Corn, Cotton, Cucurbits, Dry Beans, Peanuts, Peas, Peppers, Potatoes, Pumpkins (processing), Rice, Snap Beans, Sorghum, Squash, Sugar Beets, Tomatoes (Transplanted)
	12 Months	All crops

\*Cotton: Refer to "Requirements for Planting Time Applications" and "Replanting Instructions" for additional instructions.

Cover crops can be planted at any time, but stand reduction can occur in some areas. Do not harvest for food, graze, or feed cover crops planted less than 9 months after treatment with Willowood Clomazone 3ME.

## TOBACCO

Make a soil applied treatment of Willowood Clomazone 3ME prior to weed emergence to control annual grasses and broadleaf weeds.

### **DO NOT use on Tobacco Seedling Beds.**

Off-site movement of Willowood Clomazone 3ME by spray drift or vapors can cause foliar whitening or plant yellowing. Follow all precautions and instructions in the APPLICATION PRECAUTIONS, SPRAY DRIFT PRECAUTIONS and SPRAY DRIFT MANAGEMENT sections of this label before applying this product.

### **DO NOT spray Willowood Clomazone 3ME within 300 feet of desirable plants. Check for adequate buffer zone before application.**

#### **Preemergence Application (Prior to Transplant)**

- Apply 2.0 - 2 2/3 pints (0.75 - 1.0 lb. AI) per acre in a single broadcast application in a minimum of 20 gallons of water.
- Use the higher specified rate for heavy weed pressure or heavy soils for maximum control.
- Apply immediately or up to 30 days before transplanting.
- Apply as a surface treatment prior to weed emergence. If weeds emerge prior to preemergence application, cultivate before or immediately preceding treatment.
- If field conditions indicate the need for additional seedbed preparation, use equipment that will move this product no deeper than 1 1/2" - 2".

#### **Preemergence Application (Post-Transplant)**

- Apply 2.0 - 2 2/3 pints (0.75 - 1.0 lb. AI) per acre in a single broadcast application in a minimum of 20 gallons of water.
- Use the higher specified rate for heavy weed pressure or heavy soils for maximum control.
- Make one per season at maximum rate of 2 2/3 pints (1.0 lb. AI) per acre.
- Make an over-the-top treatment to plants immediately, up to 7 days after, or at transplanting, but ensure that application is made prior to emergence of weeds. If weeds emerge before application can be made, cultivate before or immediately preceding treatment.

#### **Combination Applications of Willowood Clomazone 3ME on Tobacco**

Tank mix Willowood Clomazone 3ME with other herbicides registered for use on tobacco to broaden the weed control spectrum compared to the use of this product applied alone.

Willowood Clomazone 3ME can be applied in combination with other insecticides, fungicides and/or nematocides registered for use in tobacco. If using a tank mixture, follow application precautions, rotational instructions, and replanting instructions of each product's label. Follow the most restrictive precautions and instructions from all products mixed.

Use water or liquid fertilizer as a carrier for Willowood Clomazone 3ME if applied alone, or if tank mixed with other tobacco pesticides unless use directions specifically state otherwise. Refer to the Mixing Instructions for tank mixing instructions.

## Tobacco Replanting Instructions

### Replanting Instructions

- If the initial planting of newly planted tobacco does not produce a uniform stand, replant tobacco in fields treated with Willowood Clomazone 3ME alone.
- If tank mixing with an approved labeled product, refer to the replant instructions for that product.
- If a tank mix is used, refer to the product's labels for any additional replant instructions, and follow the most restrictive precautions and use directions. If replanting is required, follow the directions under REQUIREMENTS FOR PLANTING TIME APPLICATIONS section.

### Replanting Restrictions

- **DO NOT** make a second application of Willowood Clomazone 3ME if fields have been retreated.
- **DO NOT** replant treated fields with any crop at intervals that are not consistent with the ROTATIONAL CROP RESTRICTIONS on this label.

### Willowood Clomazone 3ME Rates and Grasses Controlled in Tobacco

Rate	Grasses Controlled
1 1/3 pts. (0.50 lb. AI)	Suppresses some annual grasses
2.0 pts. (0.75 lb. AI)	All grasses listed above PLUS: Barnyardgrass, Large & Smooth Crabgrass, Field Sandbur, Foxtail (Giant, Green, Robust, Yellow), Goosegrass, Panicum (Common, Fall, & Texas), Johnsongrass seedling (partial control)
2 2/3 pts. (1.0 lb. AI)	All grasses listed above PLUS: Bermudagrass (partial control), Broadleaf Signalgrass, Southwestern Cupgrass, Woolly Cupgrass (partial control), Seedling Johnsongrass, Shattercane (partial control), Wild Proso Millet (partial control)



**Willowood Clomazone 3ME Rates and Broadleaf Weeds Controlled in Tobacco**

<b>Rate</b>	<b>Broadleaf Weeds Controlled</b>
1 1/3 pts. (0.50 lb. AI) 1 2/3 pts.*	Velvetleaf Spurred Anoda *Use higher rate for heavy weed pressure or heavy soil
2.0 pts. (0.75 lb. AI)	Broadleaf weeds listed above PLUS: Common Ragweed (partial control), Galinsoga, Jimsonweed (partial control), Lambsquarter, Prickly Sida, Purslane, Ladysthumb Smartweed (partial control), Pennsylvania Smartweed (partial control), Venice Mallow
2 2/3 pts. (1.0 lb. AI)	Broadleaf weeds listed above PLUS: Black Seeded Plantain, Cocklebur (partial control), Common Ragweed, Jimsonweed, Kochia, Ladysthumb Smartweed, Pennsylvania Smartweed, Tropic Croton

## Tobacco Rotational Crop Restrictions

Cover crops can be planted at any time, but stand reduction can occur in some areas. Do not harvest for food, graze, or feed cover crops planted less than 9 months after treatment with Willowood Clomazone 3ME.

### Willowood Clomazone 3ME Rotational Crop Restrictions for Tobacco in the Northern Region

Rate	Interval	Crop
1 1/3 pts. (0.50 lb. AI)	Anytime	Cabbage (transplanted), Cotton*, Peas, Peppers, Pumpkins (processing), Rice, Soybeans, Squash, Sweet Potato, Tobacco, Tuberous & Corm Vegetables
	9 Months	All crops listed above PLUS: Cabbage (Direct Seeded), Field Corn, Popcorn, Seed Corn, Sweet Corn, Cotton, Cucurbits, Dry Beans, Peanuts, Potatoes, Snap Beans, Sorghum, Sugar Beets, Tomatoes (Transplanted)
	12 Months	All crops listed above PLUS: Tomatoes (all), Wheat
	16 Months	All crops
2.0 pts. (0.75 lb. AI)	Anytime	Cotton*, Peppers, Rice, Soybeans, Winter Squash, Sweet Potato, Tobacco, Tuberous & Corm Vegetables
	9 Months	All crops listed above PLUS: Cabbage, Field Corn, Popcorn, Seed Corn, Sweet Corn, Cotton, Cucurbits, Dry Beans, Peanuts, Peas, Potatoes, Pumpkins (processing), Rice, Snap Beans, Sorghum, Summer Squash, Sugar Beets, Tomatoes (Transplanted)
	12 Months	All crops listed above PLUS: Tomatoes (all), Wheat
	16 Months	All crops
2 2/3 pts. (1.0 lb. AI)	Anytime	Cotton*, Peppers, Soybeans, Sweet Potato, Tobacco, Tuberous & Corm Vegetables
	9 Months	All crops listed above PLUS: Cabbage, Field Corn, Popcorn, Cotton, Cucurbits, Dry Beans, Peanuts, Peas, Potatoes, Pumpkins (processing), Rice, Snap Beans, Sorghum, Squash, Sugar Beets, Tomatoes (Transplanted)
	12 Months	All crops listed above PLUS: Seed Corn, Sweet Corn, Tomatoes (all), Wheat
	16 Months	All crops

\*Cotton: Refer to "Requirements for Planting Time Applications" and "Replanting Instructions" for additional instructions.

## Willowood Clomazone 3ME Rotational Crop Restrictions for Tobacco in the Southern Region

Rate	Interval	Crop
1 1/3 pts. (0.50 lb. AI)	Anytime	Cabbage (transplanted), Cotton*, Peas, Peppers, Pumpkins (processing), Rice, Soybeans, Squash, Sweet Potato, Tobacco, Tuberous & Corm Vegetables
	9 Months	All crops listed above PLUS: Cabbage (Direct Seeded), Field Corn, Popcorn, Seed Corn, Sweet Corn, Cotton, Cucurbits, Dry Beans, Peanuts, Potatoes, Snap Beans, Sorghum, Sugar Beets, Tomatoes (Transplanted)
	12 Months	All crops
2.0 pts. - 2 2/3 pts. (0.75 - 1.0 lb. AI)	Anytime	Cotton*, Peppers, Rice, Soybeans, Winter Squash (2.0 pts.), Sweet Potato, Tobacco, Tuberous & Corm Vegetables
	9 Months	All crops listed above PLUS: Cabbage, Field Corn, Popcorn, Seed Corn, Sweet Corn, Cotton, Cucurbits, Dry Beans, Peanuts, Peas, Potatoes, Pumpkins (processing), Rice, Snap Beans, Sorghum, Summer Squash, Sugar Beets, Tomatoes (Transplanted)
	12 Months	All crops
3 1/3 pt. (1.25 lb. AI)	Anytime	Cotton*, Soybeans, Sweet Potato, Tobacco, Tuberous & Corm Vegetables
	9 Months	All crops listed above PLUS: Cabbage, Field Corn, Popcorn, Seed Corn, Sweet Corn, Cotton, Cucurbits, Dry Beans, Peanuts, Peas, Peppers, Potatoes, Pumpkins (processing), Rice, Snap Beans, Sorghum, Squash, Sugar Beets, Tomatoes (Transplanted)
	12 Months	All crops

\*Cotton: Refer to "Requirements for Planting Time Applications" and "Replanting Instructions" for additional instructions.

# VEGETABLE CROPS

## **IMPORTANT: Read Before Use**

**Preemergence ground treatments of Willowood Clomazone 3ME can cause crop injury, or even loss to the following crops: Succulent snap beans, brassica (head and stem subgroup 5A), cucumbers, melons, succulent peas, peppermint, peppers, spearmint, squash (including processed pumpkin varieties), sweet potatoes, and tuberous and corm vegetables.**

Willowood Clomazone 3ME can be used as a soil applied treatment prior to weed emergence to suppress and/or control annual grasses and broadleaf weeds in snap beans (succulent), brassica (head and stem subgroup 5A), cucumbers, peas (succulent), peppers, squash (including processing pumpkins) and sweet potatoes.

Make a single application in a minimum of 10 gallons of water per acre at the rate(s) specified in the individual crop listings sections below. Apply to soil prior to weed emergence, prior to vegetable seeding or transplanting, or after vegetable seeding, but prior to crop emergence. Place seed or roots of the transplants below the chemical barrier when planting.

Tank mix Willowood Clomazone 3ME with other herbicides registered for use on vegetables to broaden the weed control spectrum compared to the use of this product applied alone. If using a tank mixture, follow application precautions, rotational instructions, and replanting instructions of each product's label. Follow the most restrictive precautions and instructions from all products mixed.

Use water or liquid fertilizer as a carrier for Willowood Clomazone 3ME if applied alone, or if tank mixed with other tobacco pesticides unless use directions specifically state otherwise. Refer to the Mixing Instructions for tank mixing instructions.

Place seeds or roots of the transplants below the chemical barrier when planting.

## **Vegetable Precautions**

Off-site movement of Willowood Clomazone 3ME by spray drift or vapors can cause foliar whitening or plant yellowing. Follow all precautions and instructions in the APPLICATION PRECAUTIONS, SPRAY DRIFT PRECAUTIONS and SPRAY DRIFT MANAGEMENT sections of this label before applying this product.

## **Vegetable Restrictions**

**DO NOT spray Willowood Clomazone 3ME within 300 feet of desirable plants. Check for adequate buffer zone before application.**

## **Crop Use Directions/Restrictions**

### **Snap Beans (Succulent)**

- Apply broadcast at 0.4 to 0.67 pint per acre.
- Use the lower specified rate on coarse soils and the higher specified rate on fine soils.
- Refer to the soil texture chart and the Precautions/Restrictions sections below for additional information.

### **Snap Bean Restrictions:**

- **DO NOT** apply within 45 days of harvest.
- **DO NOT** make more than one application per season.

### **Brassica (head and stem subgroup 5A)**

- **Direct Seeded Cabbage** - apply broadcast at 0.67 pints per acre.
- **Transplant Cabbage** - apply broadcast at 0.67 to 1.3 pints per acre.
- Use the lower specified rate on coarse soils and the higher specified rate on fine soils.
- Refer to the soil texture chart and the Precautions/Restrictions sections below for additional information.

### **Brassica (head and stem subgroup 5A) Restrictions:**

- **DO NOT** apply within 45 days of harvest.
- **DO NOT** make more than one application per season.

### **Cucumber**

- Apply broadcast at 0.4 to 1 pint per acre.
- Use the lower specified rate on coarse soils and the higher specified rate on fine soils.
- Refer to the soil texture chart and the Precautions/Restrictions sections below for additional information.

### **Cucumber Restrictions:**

- **DO NOT** apply within 45 days of harvest.
- **DO NOT** make more than one application per season.

### **Muskmelon & Watermelon**

- Apply broadcast at 0.4 to 0.67 pint per acre.
- Use the lower specified rate on coarse soils and the higher specified rate on fine soils.
- Refer to the soil texture chart and the Precautions/Restrictions sections below for additional information.

### **Peas (Succulent)**

- Apply broadcast at 1.3 pints per acre.
- Refer to the soil texture chart and the Precautions/Restrictions sections below for additional information.

### **Pea Restrictions:**

- **DO NOT** make more than one application per season.

### **Peppers**

- Apply broadcast at 0.67 to 2.67 pints per acre.
- Use the lower specified rate on coarse soils and the higher specified rate on fine soils.
- Refer to the soil texture chart and the Precautions/Restrictions sections below for additional information.

### **Pepper Restrictions:**

- **DO NOT** make more than one application per season.
- **Not registered for use on banana peppers.**

### **Squash (including Processing Pumpkin Varieties)**

- **Winter squash:** Apply broadcast prior to seeding or transplanting, at 0.67 to 2 pints per acre.
- **Summer squash:** Apply broadcast prior to seeding or transplanting, at 0.67 to 1.33 pints per acre.
- Use the lower specified rate on coarse soils and the higher specified rate on fine soils.
- Refer to the soil texture chart and the Precautions/Restrictions sections below for additional information.

### **Alternative Treatment for Squash:**

- Apply as a preemergent (to weeds) soil surface treatment before, or just prior to transplanting, to row middles between plastic covered beds.
- Use rates as noted for summer and winter squash above.
- **DO NOT** apply to soil that will be covered with plastic mulch.
- Refer to the soil texture chart and the Precautions/Restrictions sections below for additional information.

### **Squash Restrictions:**

- **DO NOT** apply within 45 days of harvest.
- **DO NOT** make more than one application per season.
- **DO NOT** use on Jack-O-Lantern pumpkins.
- **DO NOT** use on the following varieties and/or cultivars of squash: NK530, NK580, Turks Turban, Golden Delicious, all Banana types, and all other *Cucurbita maxima* types that have a pink or burnt orange coloration at harvest.

### **Sweet Potatoes**

- Apply broadcast at 1.3 to 4.0 pints per acre.
- Use the lower specified rate on coarse soils and the higher specified rate on fine soils.
- Refer to the soil texture chart and the Precautions/Restrictions sections below for additional information.

### **Alternative Treatment for Sweet Potatoes:**

- Make a single application post-transplant prior to weed emergence at a maximum of 4.0 pints per acre.
- Refer to the soil texture chart and the Precautions/Restrictions sections below for additional information.

### **Sweet Potato Restrictions:**

- **DO NOT** apply within 95 days of harvest (or 125 days if more than 3.3 pints per acre was applied).

## Tuberous and Corm Vegetables (Arracacha, Cassava, Tanier, Yams)

- Make a broadcast application at 2 to 3.3 pints per acre.
- Use the lower specified rate on coarse soils and the higher specified rate on fine soils.
- Refer to the soil texture chart and the Precautions/Restrictions sections below for additional information.

## Tuberous and Corm Vegetables (Arracacha, Cassava, Tanier, Yams) Restrictions:

- **DO NOT** apply within 125 days of harvest.
- **DO NOT** make more than one application per season.

## SOIL TEXTURES:

Select specified rates within the range based on light to heavy soil types within a textural group:

**Coarse (light) soils:** sand, loamy sand, sandy loam

**Medium soils:** loam, silt, silt loam, sandy clay, sandy clay loam

**Fine (heavy) soils:** silty clay, clay loam, silty clay loam, clay

## Willowood Clomazone 3ME Rates and Grasses Controlled in Vegetable Crops

Rate per Acre	Grasses Controlled
0.4 - 1 pt. (0.15 - 0.38 lb. AI)	Suppresses/Partially Controls: Barnyardgrass, Large & Smooth Crabgrass, Foxtail (Giant, Green, Robust), Goosegrass, Panicum (Common & Fall)
1 - 1.3 pts. (0.38 - 0.5 lb. AI)	Suppresses/All grasses listed above PLUS: Field Sandbur (suppression/partial control), Johnsongrass seedling (suppression/partial control)
1.3 - 4 pts. (0.5 - 1.5 lbs. AI)	All grasses listed above PLUS: Yellow Foxtail Controls seedling Johnsongrass at 1 - 1.5 pts./acre

## Willowood Clomazone 3ME Rates and Broadleaf Weeds Controlled in Vegetable Crops

Rate per Acre	Broadleaf Weeds Controlled
0.4 - 1 pt. (0.15 - 0.38 lb. AI)	Partially controls: <i>Galinsoga</i> spp., Lambsquarters, Spurred Anoda, Velvetleaf
1 - 1.3 pts. (0.38 - 0.5 lb. AI)	Suppresses/Partially Controls Broadleaf weeds listed above PLUS: Prickly Sida, Venice Mallow
1.3 - 4 pts. (0.5 - 1.5 lbs. AI)	Suppresses/Partially Controls Broadleaf weeds listed above PLUS: Common Ragweed*, Jimsonweed*, Purslane, Smartweed*

\*Controlled at 1 - 1.5 lbs. AI/A

### **Vegetable Replanting Instructions**

- If the initial planting of newly planted vegetables or seeds do not produce a uniform stand, replant vegetables in fields treated with Willowood Clomazone 3ME alone.
- If tank mixing with an approved labeled product, refer to the replant instructions for that product.
- If a tank mix is used, refer to the product's labels for any additional replant instructions, and follow the most restrictive precautions and use directions. If replanting is required, follow the directions under REQUIREMENTS FOR PLANTING TIME APPLICATIONS section.

### **Vegetable Replanting Restrictions**

- **DO NOT** retreat field with a second application of Willowood Clomazone 3ME.
- **DO NOT** replant treated fields with any crop at intervals that are not consistent with the Rotational Crop Restrictions on this label.
- If a tank mix is used, refer to the product's labels for any additional replant instructions, and follow the most restrictive precautions and use directions. If replanting is required, follow the directions under REQUIREMENTS FOR PLANTING TIME APPLICATIONS section.

### **Vegetable Rotational Crop Precautions**

- Cover crops can be planted at any time, but stand reduction can occur in some areas.
- Rotate crops as listed below to avoid crop injury.
- Refer to sections headed "Rotational Cropping Precautions" and "Product Information" for additional information.
- If using Willowood Clomazone 3ME with other herbicides registered for use on vegetables, always refer to rotational restrictions and precautions on the other product's label. The most restrictive label language applies.

### **Vegetable Rotational Crop Restrictions**

- **DO NOT** harvest for food, graze, or feed cover crops planted less than 9 months after treatment with Willowood Clomazone 3ME.



### Willowood Clomazone 3ME Rotational Crop Restrictions for Vegetables

Rate	Interval	Crop
0.4 - 0.67 pt. (0.15 - 0.25 lb. AI)	Anytime	Beans (succulent), Cabbage (all), Cotton*, Cucurbits, Peas, Peppers, Rice, Soybeans, Sweet Potato, Tobacco, Tuberous & Corm Vegetables
	9 Months	All crops listed above PLUS: Beans (dry), Field Corn, Popcorn, Seed Corn, Sweet Corn, Cotton, Peanuts, Potatoes, Sorghum, Sugar Beets, Tomatoes (Transplanted)
	12 Months	All crops listed above PLUS: Tomatoes (direct seeded), Wheat
	16 Months	All crops
1.0 pt. (0.38 lb. AI)	Anytime	Cabbage (transplant), Cotton*, Cucurbits, Peas, Peppers, Rice, Soybeans, Sweet Potatoes, Tobacco, Tuberous & Corm Vegetables
	9 Months	All crops listed above PLUS: Beans (succulent & dry), Cabbage (direct seeded), Field Corn, Popcorn, Seed Corn, Sweet Corn, Cotton, Peanuts, Potatoes, Sorghum, Tomatoes (Transplanted)
	12 Months	All crops listed above PLUS: Tomatoes (direct seeded), Wheat
	16 Months	All crops
1.3 pts. (0.5 lb. AI)	Anytime	Cabbage (transplant), Cotton*, Peas, Peppers, Rice, Soybeans, Sweet Potato, Tobacco, Tuberous & Corm Vegetables
	9 Months	All crops listed above PLUS: Beans (succulent & dry), Cabbage (dry seeded), Field Corn, Popcorn, Seed Corn, Sweet Corn, Cotton, Cucurbits, Peanuts, Potatoes, Sorghum, Sugar Beets, Tomatoes (Transplanted)
	12 Months	All crops listed above PLUS: Tomatoes (direct seeded), Wheat
	16 Months	All crops

\*Cotton: Refer to "Requirements for Planting Time Applications" and "Replanting Instructions" for additional instructions.

### Willowood Clomazone 3ME Rotational Crop Restrictions for Vegetables (continued)

Rate	Interval	Crop
2.0 pts. (0.75 lb. AI)	Anytime	Cotton*, Peppers, Rice, Soybeans, Squash (winter), Sweet Potatoes, Tobacco, Tuberous & Corm Vegetables
	9 Months	All crops listed above PLUS: Beans (succulent & dry), Cabbage (dry seeded & transplant), Field Corn, Peas, Popcorn, Seed Corn, Sweet Corn, Cotton, Cucurbits, Peanuts, Potatoes, Rice, Sorghum, Sugar Beets, Tomatoes (Transplanted)
	12 Months	All crops listed above PLUS: Tomatoes (direct seeded), Wheat
	16 Months	All crops
2.67 - 3.3 pts. (1.0 - 1.25 lbs. AI)	Anytime	Cotton*, Peppers (2.67 pt.), Soybeans, Sweet Potatoes, Tobacco, Tuberous & Corm Vegetables
	9 Months	All crops listed above PLUS: Beans (succulent & dry), Cabbage (dry seeded), Field Corn, Popcorn, Cotton, Cucurbits, Peanuts, Peppers, Potatoes, Soybeans, Sorghum, Sweet Potatoes, Sugar Beets, Tobacco, Tomatoes (Transplanted), Tuberous & Corm Vegetables
	12 Months	All crops listed above PLUS: Seed Corn, Sweet Corn, Tomatoes (direct seeded), Wheat
	16 Months	All crops
4 pts. (1.5 lbs. AI)	Anytime	Sweet Potatoes
	9 Months	All crops listed above PLUS: Beans (succulent & dry), Cabbage (dry seeded), Field Corn, Popcorn, Cotton, Cucurbits, Peanuts, Peppers, Potatoes, Sorghum, Soybeans, Sugar Beets, Tobacco, Tomatoes (Transplanted), Tuberous & Corm Vegetables
	12 Months	All crops listed above PLUS: Seed Corn, Sweet Corn, Tomatoes (direct seeded)
	16 Months	All crops

\*Cotton: Refer to "Requirements for Planting Time Applications" and "Replanting Instructions" for additional instructions.

## PEPPERMINT AND SPEARMINT

Review the disclaimer located directly under the heading “Vegetable Crops”.

Make a preemergent application of Willowood Clomazone 3ME at 1.3 pints per acre (0.5 lb. AI/A) to the soil surface to control annual grasses and broadleaf weeds.

### Grasses Controlled in Peppermint and Spearmint

Grasses Controlled
Suppresses/Partially Controls: Barnyardgrass, Large & Smooth Crabgrass, Foxtail (Giant, Green, Ro-bust), Goosegrass, Panicum (Common & Fall)

### Broadleaf Weeds Controlled in Peppermint and Spearmint

Broadleaf Weeds Controlled
Controls: Velvetleaf
Suppresses/Partially controls: <i>Galinsoga</i> spp., Lambsquarters, Spurred Anoda, Venice Mallow

### Peppermint and Spearmint Precautions

- Cover crops can be planted at any time, but stand reduction can occur in some areas.

### Peppermint and Spearmint Restrictions

- **DO NOT** apply within 84 days of harvest.
- **DO NOT** make more than one application per season.
- **DO NOT** apply more than 0.5 lb. AI per acre clomazone (1.3 pts. per acre Willowood Clomazone 3ME) per application.

### Peppermint and Spearmint Rotational Crop Restrictions

- Rotate crops as listed below to avoid crop injury.
- Refer to sections headed “Rotational Cropping Precautions” and “Product Information” for additional information.
- If using Willowood Clomazone 3ME with other herbicides registered for use on vegetables, always refer to rotational restrictions and precautions on the other product’s label. The most restrictive label language applies.
- **DO NOT** harvest for food, graze, or feed cover crops planted less than 9 months after treatment with Willowood Clomazone 3ME.

### Willowood Clomazone 3ME Rotational Crop Restrictions for Peppermint and Spearmint

Interval	Crop
Anytime	Cabbage (transplant), Cotton*, Peas, Peppers, Rice, Soybeans, Squash, Sweet Potatoes, Tobacco, Tuberous & Corm Vegetables
9 Months	All crops listed above PLUS: Beans (succulent & dry), Cabbage (Direct Seeded), Field Corn, Popcorn, Seed Corn, Sweet Corn, Cotton, Cucurbits, Peanuts, Potatoes, Sorghum, Sugar Beets, Tomatoes (Transplanted)
12 Months	All crops listed above PLUS: Tomatoes (direct seeded), Wheat
16 Months	All crops

\*Cotton: Refer to "Requirements for Planting Time Applications" and "Replanting Instructions" for additional instructions.

SPECIMEN

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

**PESTICIDE STORAGE:** STORE ABOVE -4°F TO KEEP PRODUCT FROM FREEZING. If frozen, thaw before use. Keep out of reach of children and animals. Store in original containers only. Store in a dry place. Carefully open containers. After partial use, replace lids and close tightly. Do not put concentrate or dilute material into food or drink containers. In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills. To confine spill: Dike surrounding area or absorb with sand, cat litter or commercial clay. Place damaged package in a holding container. Identify contents.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

### CONTAINER DISPOSAL

**Plastic: Non-refillable Containers:** Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

## IMPORTANT: READ BEFORE USE

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the unopened product container at once.

By using the product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

**CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Willowood, LLC. To the extent consistent with applicable law, such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, WILLOWOOD, LLC MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. To the extent consistent with applicable law, no agent of Willowood, LLC is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, WILLOWOOD, LLC DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

**LIMITATIONS OF LIABILITY:** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID OR AT WILLOWOOD, LLC'S ELECTION, THE REPLACEMENT OF PRODUCT.

CANOPY, GLEAN, LEXONE, LOROX, and LOROX PLUS are registered trademarks of E.I. duPont de Nemours and Co., Inc.

SONALAN and TREFLAN are registered Trademarks of Dow AgroSciences, LLC

LASSO and ROUNDUP are registered trademarks of Monsanto Company

SENCOR is a registered trademark of Bayer AG

PROWL, PURSUIT, SCEPTER, SQUADRON, and TRI-SCEPT are registered trademarks of BASF

DUAL is a registered trademark of Syngenta Group Corporation

BLAZER is a registered trademark of United Phosphorus, Inc.

SPECIMEN

SPECIMEN



# Willowood CLOMAZONE 3ME

Agricultural and Commercial Herbicide Used to Control Weeds on Snap Beans (Succulent), Brassica (head and stem subgroup 5A), Cucumbers, Cotton, Melons (Muskmelons and Watermelons), Peas (Succulent), Peppermint, Peppers, Rice, Soybeans, Spearmint, Squash, Sugarcane, Sweet Potatoes, Tobacco, and Tuberosus and Corn Vegetables

For formulation use only in California.

#### ACTIVE INGREDIENT:

Clomazone: 2-(2-Chlorophenyl)methyl-4,4-dimethyl-3-isoxazolidinone ..... 31.1%

OTHER INGREDIENTS: ..... 68.9%

TOTAL: ..... 100.0%

Contains 3.0 lbs. active ingredient per gallon

EPA Reg. No. 87290-55

**KEEP OUT OF REACH OF  
CHILDREN  
CAUTION**

#### FIRST AID

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

**IF SWALLOWED:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency medical information concerning this product, call the poison control center at 1-800-222-1222.

#### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Causes moderate eye irritation. Avoid contact with eyes or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

#### ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from the area treated. Do not apply where runoff is likely to occur. Do not contaminate water when disposing of equipment washwaters. Apply this product only as specified on this label.

#### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

**PESTICIDE STORAGE:** STORE ABOVE -4°F TO KEEP PRODUCT FROM FREEZING. If frozen, thaw before use. Keep out of reach of children and animals. Store in original containers only. Store in a dry place. Carefully open containers. After partial use, replace lids and close tightly. Do not put concentrate or dilute material into food or drink containers. In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills. To confine spill: Dike surrounding area or absorb with sand, cat litter or commercial clay. Place damaged package in a holding container. Identify contents.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

#### CONTAINER DISPOSAL

**Plastic: Non-refillable Containers:** Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

See inside booklet for complete First Aid, Precautionary Statements, and Directions for Use.

Manufactured For:

Willowood, LLC

1887 Whitney Mesa Drive #9740 Henderson, NV 89014-2069

20141112